

HEART

Editorials

- 537** Should percutaneous devices be used to close a patent foramen ovale after cerebral infarction or TIA?
J Chambers
- 539** Warfarin in atrial fibrillation: underused in the elderly, often inappropriately used in the young *R G Hart*
- 541** Changes in cognition following cardiac surgery *S Newman, J Stygall*
- 543** Lifestyle changes after a myocardial infarction *D W Johnston*
- 545** How valuable is cardiac rehabilitation and who should get it? *D R Thompson, D P de Bono*

Review

- 547** Left atrial appendage: structure, function, and role in thromboembolism *N M Al-Saady, O A Obel, A J Camm*

Papers

- 555** Left atrial appendage anatomy and function: short term response to sustained atrial fibrillation *M J Weigner, S E Katz, P S Douglas, W J Manning*
- 559** Transcatheter closure of atrial septal defects in adults with the Amplatzer septal occluder *R Dhillon, B Thanopoulos, G Tsaousis, F Triposkiadis, M Kyriakidis, A Redington*
- 563** Stroke and atrial fibrillation: is stroke prevention treatment appropriate beforehand? *D Deplanque, F Corea, C Arquizan, L Parnetti, J L Mas, V Gallai, D Leys, and the SAFE I Study Investigators*
- 570** Use of antithrombotic measures for stroke prevention in atrial fibrillation *I Perez, A Melbourn, L Kalra*
- 575** Left ventricular beat to beat performance in atrial fibrillation: dependence on contractility, preload, and afterload *H J Muntinga, A T M Gosselink, P K Blanksma, P J De Kam, E E Van Der Wall, H J G M Crijns*
- 581** Risk of myocardial infarction in young female smokers *N R Dunn, B Faragher, M Thorogood, L de Caestecker, T M MacDonald, C McCollum, S Thomas, R Mann*
- 584** Non-adherence with ACE inhibitor treatment is common in heart failure and can be detected by routine serum ACE activity assays *A D Struthers, G Anderson, R J MacFadyen, C Fraser, T M MacDonald*
- 589** Impact of concurrent amiodarone treatment on the tolerability and efficacy of carvedilol in patients with chronic heart failure *P S Macdonald, A M Keogh, C Aboyoun, M Lund, R Amor, D McCaffrey*
- 594** Evolution and long term outcome in cases with fetal diagnosis of congenital heart disease: Italian multicentre study *V Fesslova', S Nava, L Villa, and the Fetal Cardiology Study Group of the Italian Society of Pediatric Cardiology*
- 600** Stent implantation for aortic coarctation and recoarctation *A G Magee, G Brzezinska-Rajszyz, S A Qureshi, E Rosenthal, M Zubrzycka, J Ksiazyk, M Tynan*
- 607** Influence of left ventricular relaxation on the pressure half time of aortic regurgitation *S F de Marchi, S Windecker, B C Aeschbacher, C Seiler*
- 614** Utility of cardiac troponin I, creatine kinase-MB_{mass}, myosin light chain 1, and myoglobin in the early in-hospital triage of "high risk" patients with chest pain *G S Hillis, N Zhao, P Taggart, W C Dalsey, A Mangione*
- 621** A new mutation of the cardiac troponin T gene causing familial hypertrophic cardiomyopathy without left ventricular hypertrophy *A Varnava, C Baboonian, F Davison, L de Cruz, P M Elliott, M J Davies, W J McKenna*
- 625** Malalignment of the sarcomeric filaments in hypertrophic cardiomyopathy with cardiac myosin heavy chain gene mutation *A Muraishi, H Kai, K Adachi, H Nishi, T Imaizumi*
- 630** N-terminal proatrial natriuretic peptide correlates with systolic dysfunction and left ventricular filling pattern in patients with idiopathic dilated cardiomyopathy *F M Fruhwald, A Fahrleitner, N Watzinger, H Dobnig, M Schumacher, R Maier, R Zweiker, G Leb, W Klein*

BMJ
Publishing
Group



BRITISH CARDIAC SOCIETY

contents continued inside