Isovolumic contraction time of right ventricle in d-transposition of great arteries*

JEAN-CLAUDE FOURON, FRANÇOIS VALLOT, FRANÇOIS BOURLON, MICHEL LOMBAERT, GILLES DUCHARME, ANDRÉ DAVIGNON

From the section of Cardiology, Departments of Pediatrics and Pediatric Research, Ste-Justine Hospital and University of Montreal, Canada

SUMMARY The pre-ejection period of the right ventricle in d-transposition of the great arteries is known to be prolonged, compared with the same interval of the left ventricle of normal subjects. In the present study, the echocardiographic measurement of the components of the pre-ejection period of the right ventricle of 14 patients with d-transposition of the great arteries shows that the isometric contraction time of the right ventricle in d-transposition of the great arteries is similar to the same interval calculated on the left ventricle of 76 normal children of comparable age. On the other hand, the electromechanical delay was significantly greater for the right ventricle of d-transposition of the great arteries than for the left ventricle of the normal subjects. It is concluded that the prolonged pre-ejection period of the right ventricle in d-transposition of the great arteries is not the result of right ventricular dysfunction but solely of a longer electromechanical delay.

In d-transposition of the great arteries (d-TGA), the right ventricle must maintain a normal output against the high resistance of the systemic circulation. Its ability to sustain this load throughout life has been seriously questioned since angiographic studies have shown that right ventricular end-diastolic volume is abnormally increased and the ejection fraction decreased, both before and after intra-atrial repair by the Mustard procedure.^{1 2}

A recent publication on the assessment of myocardial function after Mustard's operation showed that the ratio of systolic time intervals for the right and left ventricles was appropriate for normal systemic and pulmonary ventricles, respectively.³ On the other hand most studies on myocardial function using systolic time intervals seem to be in agreement with volume measurements and have shown a prolonged right ventricular pre-ejection period and a shortened right ventricular ejection time in d-TGA suggesting that right ventricular function was depressed.^{4 5} However, these reports did not specify which part of the pre-ejection period, that is the electromechanical delay or the isovolumic contraction time, was responsible for prolonging the right ventricular pre-ejection period. Clarifica-

tion of this point is essential before any conclusion can be drawn about the performance of the right ventricle. Indeed, isovolumic contraction time is the component of the pre-ejection period widely recognised as a reliable index of myocardial contractility.^{6 7}

The frequent clinical observation that post-operative patients with d-TGA achieve normal physical performance without any sign of cardiac failure, and the fact that a prolonged right ventricular pre-ejection period has been found right after birth in this disease,⁸ prompted us to investigate the possibility that in d-TGA, the right ventricular pre-ejection period was prolonged because of the lengthening of the right ventricular electromechanical delay.

Subjects and methods

Fourteen patients with d-TGA, surgically repaired by Mustard procedure, were included in this study. In all but two an atrial septostomy had been done previously. Their ages varied from 3½ to 16 years, with a mean of 7 years. Duration of follow-up after corrective surgery averaged three years and eight months (range: one to six years). They were all clinically asymptomatic. Associated lesions included mild pulmonary stenosis in two and a small ventricular septal defect in one. In all patients, the

* Work supported in part by the Canadian Heart Foundations, La Fondation Justine-Lacoste-Beaubien and the MacDonald Stewart Foundation.

Received for publication 29 October 1979

standard electrocardiogram showed right ventricular hypertrophy, without evidence of abnormal intra-ventricular conduction delay. Another group of 76 healthy children, aged 2½ to 16 years, with a mean of 8 years, was chosen for comparison.

In all cases echocardiograms were taken with an Echoline 20 A ultrasonoscope, coupled with a Cambridge fibreoptic recorder. An electrocardiographic lead with a well-identified Q wave was simultaneously recorded. Systolic time intervals were measured according to techniques already described.⁵⁻⁹ The pre-ejection period of the systemic ventricle was calculated from the beginning of the Q wave on the electrocardiogram to the opening point of the aortic valve leaflets and identified as right ventricular pre-ejection period (RPEP) for the d-TGA group and left ventricular pre-ejection period (LPEP) for the normals. The corresponding isovolumic contraction time is, for the d-TGA, the difference between Q to aortic valve opening and Q to tricuspid valve closure (RICT) (Fig), and for the normals, the difference between Q to aortic valve opening and Q to mitral valve closure (LICT). The electromechanical delay was calculated as RPEP less RICT (REMD), and LPEP less LICT (LEMD). In the group of normal subjects, the pre-

Fig. Composite of aortic and tricuspid echograms of a patient with d-TGA showing the technique of measurement of Q to aortic valve opening (Q Ao) and Q to tricuspid valve closure (Q Tc). Isovolumic contraction time = Q Ao - Q Tc.

ejection period of the right ventricle was measured from the beginning of the Q wave to the opening point of the pulmonary valves; isovolumic contraction time was calculated as the difference of Q to pulmonary valve opening less Q to tricuspid valve closure, and electromechanical delay was the difference of the pre-ejection period less isovolumic contraction time. The paper speed was set as 100 mm/s, allowing measurement to the nearest 5 ms.¹⁰ The average of at least five cardiac cycles was used in the calculations. Measurements were made at similar RR intervals.

In order to select the proper parametric tests for the statistical analysis of our materials, normality of the data was assessed by Lilliefors's test,¹¹ and equality of variances by Bartlett's test.¹² The Student's t test was then used to evaluate the difference between right ventricular pre-ejection period of the d-TGA group and left ventricular pre-ejection period of the normal subjects. A one-way analysis of variance with fixed effects was applied to compare the right ventricular isovolumic contraction time and electromechanical delay of patients with d-TGA, to the same intervals calculated on the left and right ventricular echocardiograms of the group of normal children. A contrast analysis using Scheffe's technique¹³ was applied to determine the means that were significantly different.

In the group of normal subjects, the regression equations of isovolumic contraction time and electromechanical delay versus heart rate yielded weak correlation coefficients ($r=0.44$ and 0.22 , respectively). This is in accordance with previous publications on the subject.⁹⁻¹⁴ For this reason, comparisons were made between values uncorrected for heart rate.

Results

Values found for the components of the pre-ejection period of the systemic right ventricle of the 14 children with d-TGA are given in the Table, as well as the mean values for the left and right ventricles of 76 normal subjects. The mean pre-ejection period of the right ventricle of patients with d-TGA was significantly greater (98 ms, SD=13) than the same interval for the left ventricle of normal children (79 ms, SD=10) ($p<0.001$). The right ventricular isovolumic contraction time of the d-TGA was not prolonged (26 ms, SD=5) when compared with values obtained in normal left ventricles (33 ms, SD=8). On the other hand, the electromechanical delay of the right ventricle in d-TGA averaged 72 ms (SD=11), while the same interval measured on the left ventricle of normal children was 46 ms (SD=8). The difference

between these two values is highly significant ($p < 0.001$).

The average value of 72 ms found for the electromechanical delay of the right ventricle of patients with d-TGA was higher ($p < 0.01$) than that observed in the right ventricle of our group of normals (60 ms, SD=9). The same observation can be made for the right ventricular isovolumic contraction time in d-TGA (26 ms, SD=5) and in normal subjects (13 ms, SD=8) ($p < 0.01$).

Table 1 Pre-ejection period and its components in systemic ventricle of patients with d-TGA and in left and right ventricles of 76 normal children

Case no.	Age (y)	PEP (ms)	ICT (ms)	EMD (ms)
Patients with d-TGA				
1	3½	100	25	75
2	4½	80	27	53
3	5	100	20	80
4	5	92	20	72
5	5	105	25	80
6	5	80	22	58
7	5	78	28	50
8	5	103	25	78
9	7	105	30	75
10	8	85	25	60
11	9	118	38	80
12	9	110	28	82
13	10	105	25	80
14	16	110	30	80
Mean	7	98 (SD 13)	26 (SD 5)	72 (SD 11)
Range: 3½ to 16				
Normal children:				
Mean				
LV	8	79 (SD 10)	33 (SD 8)	46 (SD 8)
Range: 2½ to 16				
Mean				
RV	—	73 (SD 9)	13 (SD 8)	60 (SD 9)

Abbreviations: PEP, pre-ejection period; ICT, isovolumic contraction time; EMD, electromechanical delay; SD, standard deviation; LV, left ventricle; RV, right ventricle.

Comments

The ability of the right ventricle to perform adequately as a systemic pump is crucial for the long-term prognosis of patients with d-TGA having undergone intra-atrial rerouting of the venous return. Previous angiographic studies have compared volumetric parameters of the right ventricle of patients with d-TGA with those of the right ventricle of normal individuals. Abnormal values were always found.^{1-2,15} In fact, the question is how the right ventricle in d-TGA compares with the left ventricle of normal individuals. Such an approach has been applied in echocardiographic studies, where right ventricular systolic time intervals of d-TGA were compared with left ventricular intervals of normal subjects. In most of these studies, the conclusion that the right ventricular function was depressed in d-TGA, was based on

finding a prolonged right ventricular pre-ejection period.^{4,5} The present study demonstrates that this conclusion is unwarranted. Indeed, the part of the pre-ejection period which closely correlates with measures of contractility, that is the isovolumic contraction time, is, in our group of d-TGA, similar to the same interval measured on the left ventricle of normal subjects. This finding is in full agreement with the normal pressure-velocity indices (dP/dt and derived velocity variables) previously reported for the right ventricle of d-TGA.^{16,17} The apparent contradiction between our results and the data presented in the angiographic reports may be explained by the work of Peterson and co-workers,¹⁸ who showed that changes in ejection fraction volumes and modification of the contractile state could occur independently of each other.

According to the present study, it is the prolongation of the right ventricular electromechanical delay which is responsible for the longer right ventricular pre-ejection period described in d-TGA when compared with the normal left ventricle. A previous report on time relation of dynamic events in the cardiac chambers, has documented a longer electromechanical delay for the right ventricle than for the left ventricle in normal hearts.¹⁹ Moreover, in our study, the right ventricular electromechanical delay in d-TGA has been found to be longer than that of normal right ventricles. This could be because of the greater myocardial mass of the hypertrophied right ventricle in d-TGA, resulting in a longer ventricular activation time.²⁰

We believe that serial evaluation by systolic time intervals of right ventricular function in d-TGA should be based on the measurement of the isovolumic contraction time in preference to that of the complete pre-ejection period.

References

- Graham TP, Atwood GF, Boucek RJ, Boerth RC, Nelson JH. Right heart volume characteristics in transposition of the great arteries. *Circulation* 1975; **51**: 881-9.
- Graham TP, Atwood GF, Boucek RJ, Boerth RC, Bender HW. Abnormalities of right ventricular function following Mustard's operation for transposition of the great arteries. *Circulation* 1975; **52**: 678-84.
- Silverman NH, Payot M, Stanger P, Rudolph AM. The echocardiographic profile of patients after Mustard's operation. *Circulation* 1978; **58**: 1083-93.
- Korfhagen J, Meyer R, Kaplan S. Non-invasive assessment of right ventricular performance in transposition of the great arteries (abstract). *Am J Cardiol* 1977; **39**: 310.
- Hirschfeld S, Meyer R, Schwartz DC, Korfhagen J, Kaplan S. Measurement of right and left ventricular

- systolic time intervals by echocardiography. *Circulation* 1975; **51**: 304-9.
- 6 Ahmed SS, Levinson GE, Schwartz CJ, Ettinger PO. Systolic time intervals as measures of the contractile state of the left ventricular myocardium in man. *Circulation* 1972; **46**: 559-71.
 - 7 Mason DT. Usefulness and limitations of the rate of rise of intraventricular pressure (dP/dt) in the evaluation of myocardial contractility in man. *Am J Cardiol* 1969; **23**: 516-27.
 - 8 Fouron JC, Lombaert M, Payot M, Davignon A. Systolic time intervals during the first hours of life in D-transposition of the great arteries (abstract). Can Cardiovasc Soc, 30th Annual Meeting, Toronto.
 - 9 Hirschfeld S, Meyer R, Korfhagen J, Kaplan S, Liebman J. The isovolumic contraction time of the left ventricle. An echographic study. *Circulation* 1976; **54**: 751-6.
 - 10 Spodick DH, Ball HG, Pigott VM. Effects of recording speed on precision of time-based polycardiographic measurements. Optimal paper speeds for measuring points and intervals. *Br Heart J* 1978; **40**: 1344-8.
 - 11 Lilliefors HW. On the Kolmogorov-Smirnov test for normality with mean and variance unknown. *J Am Stat Assoc* 1967; **62**: 399-402.
 - 12 Winer BJ. *Statistical principles in experimental design*. 2nd ed. New York: McGraw-Hill, 1971: 207.
 - 13 Scheffe H. *The analysis of variance*. New York: John Wiley, 1959: 68.
 - 14 Spitaels S, Arbogast R, Fouron JC, Davignon A. The influence of heart rate and age on the systolic and diastolic time intervals in children. *Circulation* 1974; **49**: 1107-15.
 - 15 Jarmakani JM, Canent RV. Preoperative and postoperative right ventricular function in children with transposition of the great vessels. *Circulation* 1974; **49** and **50**, suppl II: 39-45.
 - 16 Godman MJ, Friedli B, Pasternac A, Kidd BSL, Trusler GA, Mustard WT. Hemodynamic studies in children four to ten years after the Mustard operation for transposition of the great arteries. *Circulation* 1976; **53**: 532-8.
 - 17 Pasternac A, Hawker RE, Kidd L. Right and left ventricular function in complete transposition of the great arteries (abstract). *Am J Cardiol* 1973; **31**: 152.
 - 18 Peterson KL, Skloven D, Ludbrook P, Uther JB, Ross J Jr. Comparison of isovolumic and ejection phase indices of myocardial performance in man. *Circulation* 1974; **49**: 1088-101.
 - 19 Braunwald E, Fishman AP, Cournand A. Time relationship of dynamic events in the cardiac chambers, pulmonary artery and aorta in man. *Circ Res* 1956; **4**: 100-7.
 - 20 Scott RC. The correlation between the electrocardiographic patterns of ventricular hypertrophy and the anatomic findings. *Circulation* 1960; **21**: 256-91.
- Requests for reprints to Dr Jean-Claude Fouron, Ste-Justine Hospital, 3175 Ste-Catherine Road, Montreal, Quebec H3T 1C5, Canada.