- 1990;336:129–33.
- 79 Blankenhorn D, Nessim S, Johnson R, San Marco M, Azen A, Cashin-Hemphill L. Beneficial effects of combined colestipol-niacin therapy on coronary atherosclerosis and coronary venous bypass grafts. *JAMA* 1987;257:3233–40.
 - 80 Buchwald H, Varco RL, Matts JP, *et al.* Effect of partial ileal bypass surgery on mortality and morbidity from coronary heart disease in patients with hypercholesterolemia. *N Engl J Med* 1990;323:946–55.
 - 81 Brown G, Albers JJ, Fisher LD, *et al.* Regression of coronary artery disease as a result of intensive lipid-lowering therapy in men with high levels of apolipoprotein B. *N Engl J Med* 1990;323:1289–98.
 - 82 Lichtlen P, Hugenholtz P, Rafflenbeul W, Hecker H, Jost S, Deckers J. Retardation of angiographic progression of coronary artery disease by nifedipine. *Lancet* 1990; 335:1109–13.
 - 83 Gottlieb S, Brinker J, Mellits E, *et al.* Effect of nifedipine on the development of coronary bypass graft stenoses in high risk patients. A randomized double blind, placebo-controlled trial [Abstract]. *Circulation* 1989;80(suppl II):A909.
 - 84 Loaldi A, Polese A, Montorsi P, *et al.* Comparison of nifedipine, propranolol and isosorbide dinitrate on angiographic progression and regression of coronary arterial narrowings in angina pectoris. *Am J Cardiol* 1989;64:433–9.
 - 85 Waters D, Lesperance J, Theroux P, *et al.* A controlled clinical trial to assess the effect of a calcium channel blocker on the progression of coronary atherosclerosis. *Circulation* 1990;82:1940–53.
 - 86 Strong J, Solberg L, Restrepo C. Atherosclerosis in persons with coronary heart disease. *Lab Invest* 1968;18:527–37.
 - 87 Holme I. An analysis of randomized trials evaluating the effect of cholesterol reduction on total mortality and coronary heart disease incidence. *Circulation* 1990;82:1916–24.
 - 88 Lipids Research Clinics Program. The Lipid Research Clinics Coronary Primary Prevention Trial Results: II The relationship of reduction in incidence of coronary heart disease to cholesterol lowering. *JAMA* 1984;251:365–74.

PLANTS IN CARDIOLOGY

Coffea arabica L. From Bentley R, Trimmen H. *Medicinal plants*. London: J and A Churchill, 1880:144.

Theobroma cacao L. From Kohler FE. *Medizinal-Pflanzen, Atlas* 1887; Band II: plate 183.

Camellia sinensis (L.) Kuntze. From Curtis W. *Botanical magazine* 1807;25 and 26: Tab 998.

Methyl xanthine diuretics

Caffeine, theobromine, and theophylline were first studied in 1886 and although they are now obsolete they were important before the discovery of the synthetic diuretics in 1951 and especially before organic mercurials were introduced in 1920. They act by depression of renal tubular reabsorption. Caffeine is only a weak diuretic. Theophylline was shown to be better than theobromine by H M Marvin in an excellent early clinical trial (*Journal of the American Medical Association* 1926;87: 2043–6). It was effective in two thirds of patients who were still oedematous after treatment with digitalis. Theophylline also has a direct stimulant effect on the myocardium which was demonstrated at cardiac catheterisation by McMichael and colleagues at Hammersmith Hospital (*Clinical Science* 1946–48;6:125–35) and shown by them to be greater than that of digoxin in hypertensive right heart failure.

Theophylline is found in the tea plant *Camellia sinensis* (Theaceae) a native of China and India but the amount, 0.1%, is small for clinical use so it has to be synthesised. Theobromine is present in the cocoa tree *Theobroma cacao* (Sterculiaceae) which comes

from the tropical forests of South America. Caffeine is present in both tea and cocoa and in other traditional beverages. In Africa these are coffee *Coffea arabica* (Rubiaceae) and the cola nut *Sterculia acuminata* (Sterculiaceae). In South America the ancient drink maté is made from a species of holly *Ilex paraguariensis* (Aquifoliaceae) while yoco comes from a species of *Paullinia* (Sapindaceae). Thus the methyl xanthines come from six plant families in the Old and New World that have no obvious botanical similarities. As has so often happened with medicinal plants, their therapeutic value was discovered by chance.

Purgatives were often used to treat heart failure before the xanthine diuretics were introduced and several of them were plants—elaterium, senna, aloes, cascara, croton, rhubarb, and podophyllum. Presumably they produced their effect because of the coincidental sodium loss that accompanied diarrhoea.

Theophylline seems to be making an exciting therapeutic comeback. It is now used to control erythrocytosis in patients with renal transplants. It reduces erythropoietin production by adenosine antagonism (*New England Journal of Medicine* 1990;323:86–90.

A HOLLMAN