Editorial

Hypoxia and the heart

The heart can be exposed to hypoxia in several circumstances including pulmonary disease, pulmonary embolism, aspiration, during cardiac arrest, and at high altitude. In patients with chronic pulmonary diseases there may be severe hypoxia on exercise and during sleep.^{1,2} The mechanisms of nocturnal hypoxia are complex, and include reductions in alveolar ventilation and mismatching of ventilation and perfusion.^{3,4} Hypoxia in these patients leads to secondary pulmonary hypertension and cor pulmonale.^{5,6} Patients with sleep apnoea syndromes, either obstructive or central, also have episodes of nocturnal hypoxia.

There have been few studies of the effects on the heart of arterial hypoxaemia as distinct from myocardial ischaemia. It is likely that the effects differ: first, arterial hypoxaemia is generalised whereas myocardial ischaemia is usually localised; and secondly, the cellular metabolism of ischaemic myocardium differs from that of perfused but hypoxic myocardium.^{7,8}

The effects of hypoxia on myocytes are similar to those in other cells. There is a decline in ATP concentrations and increased glucose uptake and anaerobic glycolysis, the resulting pyruvate being largely converted into lactate and alanine.⁸ During hypoxia without ischaemia there is a sustained increase in glycolytic flux with less intracellular acidosis than occurs during ischaemia.⁸ Hypoxia also causes less mitochondrial damage, and the rate of ATP synthesis recovers more rapidly after a period of non-ischaemic hypoxia than after ischaemia.^{9,10}

Unlike localised myocardial ischaemia, hypoxaemia will also affect the heart via direct and reflex effects on other sites in the circulation. Hypoxia acts directly as a vasodilator especially in the coronary and cerebral vessels, and stimulation of the carotid body chemoreceptors causes reflex vasoconstriction of the splanchnic, muscle, pulmonary, and cutaneous beds, with either vasodilation or no net effect in the coronary and cerebral beds.¹¹

Direct effects of hypoxaemia on pump function

Although there have been many studies on the effects of hypoxia on isolated hearts and muscle preparations *in vitro* there are few reports in humans or intact animals. Observations on those who travel to high altitude are therefore of interest: PO_2 is 150 mm Hg at sea level but only 80 mm Hg at 14 000 feet. Typically the resting heart rate increases acutely, but returns to slightly above pre-ascent values in one to two days.¹² Stroke volume gradually decreases over 5–10 days to 20% below pre-ascent values.¹³ Echocardiography shows a decrease in left ventricular end diastolic and end systolic diameters, while the ejection fraction and other indices of contractility remain normal.¹⁴ Cardiac output increases acutely and then decreases over the next 2–5 days to a steady state some 20% below resting cardiac output at sea level.¹² During exercise at a given submaximal work rate, cardiac output is approximately 20% below that during exercise at the same work rate at sea level.¹⁵

From the observation that echocardiographic indices of contractility remain normal, it has been argued that the decrease in stroke volume represents a physiological adaptation rather than impaired myocardial function.¹⁴ The reduction in cardiac output may be secondary to improved

oxygen delivery to the tissues caused by an increased haematocrit and adaptive changes in haemoglobin-oxygen binding.

In chronic hypoxic lung disease the function of left heart is usually normal as judged by left ventricular ejection fraction, pulmonary wedge pressure, and cardiac output.^{16–18} When coronary disease coexists, however, hypoxia together with carbon dioxide retention and acidosis may precipitate left ventricular failure.¹⁹ In postmortem studies of patients dying of chronic cor pulmonale, thickening and patchy fibrosis of the left ventricular myocardium are common.^{20,21} Left ventricular function may also be affected in chronic airflow limitation by the wide swings in intrathoracic pressure and by hypertrophy of the right ventricle.^{22,23}

In patients with sleep-disordered breathing there may be episodes of acute hypoxaemia, during which cardiac output is reduced²⁴ and pulmonary wedge pressure may increase.²⁵ In obstructive sleep apnoea these changes may be the result of the changes in intrathoracic pressure rather than hypoxaemia.²⁶ Acute severe hypoxaemia also occurs in adult respiratory distress syndrome, fat embolism, and pulmonary embolism, and is often associated with severe depression of cardiac output, but in these conditions hypoxaemia is only one of many pathophysiological changes.

Effects of hypoxaemia on coronary blood flow

Experiments in healthy individuals showed that breathing hypoxic gas mixtures considerably increased coronary blood flow: a reduction of the inspired concentration of oxygen to 10% led to a doubling of coronary flow as the coronary arterial oxygen content was reduced.²⁷ After ascent to high altitude, coronary blood flow shows a small acute increase, and then decreases over 7–10 days to 25% below pre-ascent values, while the coronary arteriovenous difference in oxygen content increases.²⁸ It is thought that this results from increased unloading of oxygen in the capillaries caused by a rightwards shift in the haemoglobin-oxygen binding curve which occurs as an adaptation to high altitude. Myocardial oxygen consumption remains unchanged or decreases slightly.²⁸

Changes in coronary flow have not been reported in relation to hypoxaemia in other clinical situations. In animal experiments it seems that hypoxia causes dilatation of coronary microvessels that is mediated by prostaglandins rather than by an endothelium-derived relaxant factor.²⁹

Effects of hypoxaemia on electrophysiology

Hypoxia in healthy individuals does not seem to cause arrhythmias either when a hypoxic gas mixture is breathed²⁷ or at high altitude,¹² unless ascent to high altitude results in acute mountain sickness with pulmonary oedema. In contrast, brief episodes of hypoxaemia during sleep apnoea are associated with various arrhythmias including prolonged sinus pauses, atrial fibrillation, transient atrioventricular conduction block, multifocal ventricular extrasystoles, and non-sustained ventricular tachycardia.^{30–33} The mechanisms of arrhythmogenesis are

not clear,³⁴ and although there is a close correlation between the level of hypoxaemia and the occurrence of ventricular extrasystoles,³⁵ other factors including hypercapnia and autonomic changes may contribute to arrhythmias.³⁶ Indeed, ventricular arrhythmias may occur during sleep in subjects without apnoea,³⁷ and may be related to the nocturnal reduction in mean heart rate.³⁸ The arrhythmias associated with sleep apnoea are improved or abolished when patients with obstructive sleep apnoea are treated by tracheostomy³⁹ or when obese patients lose weight.^{40,41} Patients with severe chronic left ventricular failure may also show episodes of oxygen desaturation during sleep. Although the severity of hypoxaemia is less than in hypoxic lung disease, some of these episodes are associated with arrhythmias^{42,43} and might relate to the increased risk of sudden death in this condition.⁴²

Hypoxic pulmonary vasoconstriction

Hypoxia is a potent stimulus to pulmonary vasoconstriction, but the relation of pulmonary vascular resistance to hypoxia is complex. First it is non-linear: studies of healthy individuals showed that pulmonary vascular resistance increased particularly rapidly when P_{O_2} was ≤ 8 kPa.⁴⁴ Second, the response to hypoxia varies widely between individual subjects,⁴⁵ and an individual's pulmonary vascular reactivity may determine his risk of developing cor pulmonale at altitude or with chronic lung disease.⁴⁶ Some of those who reside at altitudes of 10 000 feet or above develop pulmonary hypertension and features of right ventricular strain, but many do not.⁴⁷

In chronic hypoxic lung disease other factors besides hypoxic pulmonary vasoconstriction may be important in the development of pulmonary hypertension. The rate of rise of pulmonary artery pressure in these patients is variable⁶ and long-term oxygen therapy has only a small effect on pulmonary hypertension.^{48,49} Recent evidence from necropsy studies also suggests that hypoxic pulmonary vasoconstriction may be less important: abnormalities of the intima and development of longitudinal muscle in the media were noted in these patients.^{50,51} These changes may be caused by repeated stretching and distortion of the small pulmonary vessels; whereas hypoxia causes circular muscle hypertrophy. Emphysematous destruction of the pulmonary vascular bed can also increase pulmonary artery pressure.⁵² Hypoxia causes secondary polycythaemia, further increasing pulmonary vascular resistance to a small extent.^{53,54}

The mechanisms of cor pulmonale in severe chronic hypoxic lung disease are probably multifactorial. As well as increased pulmonary vascular resistance owing to a combination of vasoconstriction and destruction of the pulmonary vasculature, hypoxia increase renin, aldosterone, and vasopressin (anti-diuretic hormone) concentrations.⁵⁴ Together with reduced renal blood flow these lead to salt and water retention, and these changes may be more important than any direct effects of hypoxaemia on the heart in this condition.^{56,57}

Conclusions

It is perhaps surprising how little is known about the effects on the heart of hypoxia as opposed to ischaemia. The cardiovascular responses of healthy individuals to hypoxia have been studied in the laboratory and during ascent to high altitudes, but the effects of hypoxia on diseased hearts are not well understood. Chronic hypoxic lung disease and sleep apnoea have many effects on the heart and circulation, and it is not known whether hypoxia itself or other mechanisms are the more important. Potentially serious

arrhythmias during sleep apnoeas and in heart failure may be triggered by hypoxaemia. It is important to be aware of these effects because episodes of hypoxia occur in various conditions, and are probably underdiagnosed. It is possible that oxygen therapy will be beneficial in some circumstances but much work remains to be done.

S W DAVIES
J A WEDZICHA

London Chest Hospital, London E2

- Wynne JW, Block AJ, Hemenway J, Hunt LA, Flick MR. Disordered breathing and oxygen desaturation during sleep in patients with chronic obstructive lung disease. *Am J Med* 1979;66:573-9.
- Douglas NJ, Calverley PMA, Leggett RJE, Brash HM, Flenley DC, Brezinova V. Transient hypoxaemia during sleep in chronic bronchitis and emphysema. *Lancet* 1979;i:1-4.
- Littner MR, McGinty DJ, Arand DL. Determinants of oxygen desaturation in the course of ventilation during sleep in chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1980;122:849-57.
- Catterall JR, Douglas NJ, Calverley PMA, Shapiro C, Brezinova V, Flenley DC. Transient hypoxemia during sleep in chronic obstructive pulmonary disease is not a sleep apnea syndrome. *Am Rev Respir Dis* 1983;128:24-9.
- Flenley DC. Clinical hypoxia: causes, consequences and correction. *Lancet* 1978;i:542-6.
- Weitzenblum E, Sautegau A, Ehrhart M, Mammosser M, Hirth C, Roegel E. Long-term course of pulmonary pressure in chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1984;130:993-8.
- Rovetto MJ, Whitmer JT, Neely JR. Comparison of the effects of anoxia and whole heart ischemia on carbohydrate utilisation in isolated working rat hearts. *Circ Res* 1973;32:699-711.
- Neely JR, Morgan HE. Relationship between carbohydrate and lipid metabolism and the energy balance of heart muscle. *Annu Rev Physiol* 1974;36:413-59.
- Mela L, Bacalzo LV Jr, Miller LD. Defective oxidative metabolism of rat liver mitochondria in hemorrhagic and endotoxin shock. *Am J Physiol* 1971;220:571-7.
- Trump BF, Mergner WJ, Kahng MW, Saladino AJ. Studies on the subcellular pathophysiology of ischemia. *Circulation* 1976;53(suppl 1):17-26.
- Abbond FM, Heistad DD, Mark AL, Schmid PG. Reflex control of the peripheral circulation. *Progr Cardiovasc Dis* 1976;18:371-403.
- Vogel JA, Harris CW. Cardiopulmonary responses of resting man during early exposure to high altitude. *J Appl Physiol* 1967;22:1124-8.
- Alexander JK, Hartley LH, Modelski M, Grover RF. Reduction in stroke volume during exercise in man following ascent to 3100m altitude. *J Appl Physiol* 1967;23:849-58.
- Alexander JK, Grover RF. Mechanism of reduced stroke volume at high altitude. *Clin Cardiol* 1983;6:301-3.
- Wolfel EE, Groves BM, Brooks GA, Butterfield GE, Mazzeo RS, Moore LG, et al. Oxygen transport during steady-state submaximal exercise in chronic hypoxia. *J Appl Physiol* 1991;70:1129-36.
- Fishman AP. The left ventricle in "chronic bronchitis and emphysema". *N Engl J Med* 1971;285:402-4.
- Steele P, Ellis JH Jr, Van Dyke D, Sutton F, Creagh E, Davies H. Left ventricular ejection fraction in severe chronic obstructive airways disease. *Am J Med* 1975;59:21-8.
- Christianson LC, Shah A, Fisher VJ. Quantitative left ventricular cineangiography in patients with chronic obstructive pulmonary disease. *Am J Med* 1979;66:399-404.
- Murphy ML, Adamson J, Hutcheson F. Left ventricular hypertrophy in patients with chronic bronchitis and emphysema. *Ann Intern Med* 1974;81:307-13.
- Michelson N. Bilateral ventricular hypertrophy due to pulmonary disease. *Br J Dis Chest* 1980;38:435-7.
- Kohama A, Tanouchi J, Hori M, Kitabatake A, Kamada T. Pathologic involvement of the left ventricle in chronic cor pulmonale. *Chest* 1990;98:794-800.
- Bemis CE, Serur JR, Borkenhagen D, Sonnenblick EH, Urschel CW. Influence of right ventricular filling pressure on left ventricular pressure and dimension. *Circ Res* 1974;34:498-504.
- Buda AJ, Pinsky MR, Ingels NB Jr, Daughters GT II, Stinson EB, Alderman EL. Effect of intrathoracic pressure on left ventricular performance. *N Engl J Med* 1979;301:453-9.
- Guilleminault C, Motta J, Mihn F, Melvin K. Obstructive sleep apnoea and cardiac index. *Chest* 1986;89:331-4.
- Buda AJ, Schroeder JS, Guilleminault C. Abnormalities of pulmonary wedge pressures in sleep-induced apnea. *Int J Cardiol* 1981;1:67-74.
- Tolle FA, Judy WV, Yu P-L, Markand ON. Reduced stroke volume related to pleural pressure in obstructive sleep apnoea. *J Appl Physiol* 1983;55:1718-24.
- Hellem H, Ord JW, Talmers FN, Christensen RC. Effects of hypoxia on coronary blood flow and myocardial metabolism in normal human subjects. [Abstract]. *Circulation* 1957;16:893.
- Grover RF, Lufschanowski R, Alexander JK. Alterations in the coronary circulation of man following ascent to 3,100 m altitude. *J Appl Physiol* 1976;41:832-8.
- Myers PR, Muller JM, Tanner MA. Effects of oxygen tension on endothelium dependent responses in canine coronary microvessels. *Cardiovasc Res* 1991;25:885-94.
- Imaizumi T. Arrhythmias in sleep apnoea. *Am Heart J* 1990;100:513-6.
- Miller WP. Cardiac arrhythmias and conduction disturbances in the sleep apnea syndrome: prevalence and significance. *Am J Med* 1982;73:317-21.
- Guilleminault C, Connolly SJ, Winkle RA. Cardiac arrhythmias and conduction disturbances during sleep in 400 patients with sleep apnea syndrome. *Am J Cardiol* 1983;52:490-4.
- Orr WC. Editorial: sleep apnoea, hypoxaemia, and cardiac arrhythmias. *Chest* 1986;89:1-2.

- 34 Zwillich C, Delvin T, White D, Douglas NJ, Weil J, Martin R. Bradycardia during sleep apnoea; characteristics and mechanisms. *J Clin Invest* 1982;69:1286-92.
- 35 Shepard JW Jr, Garrison MW, Grither DA, Dolan GF. Relationship of ventricular ectopia to oxyhaemoglobin desaturation in patients with obstructive sleep apnoea. *Chest* 1985;88:335-40.
- 36 Kawakami T, Otsuka K, Saito H, et al. Arrhythmogenesis during sleep-induced apnoea (letter). *Am Heart J* 1984;108:1591-3.
- 37 Lown B, Tykocinski M, Garfein A, Brooks P. Sleep and ventricular premature beats. *Circulation* 1973;48:691-701.
- 38 Winkle RA. The relationship between ventricular ectopic beat frequency and heart rate. *Circulation* 1982;66:439-46.
- 39 Tilkian A, Motta J, Guilleminault C. Cardiac arrhythmias in sleep apnea syndromes. New York: Alan Liss, 1978.
- 40 Estes EH Jr, Sieker HO, McIntosh HD, Kelser GA. Reversible cardiopulmonary syndrome with extreme obesity. *Circulation* 1957;16:179-87.
- 41 Peiser J, Ovnat A, Uwytyed K, Lavie P, Charuzi I. Cardiac arrhythmias during sleep in morbidly obese sleep-apnoeic patients before and after gastric bypass surgery. *Clin Cardiol* 1985;8:519-21.
- 42 Davies SW, John LM, Wedzicha JA, Lipkin DP. Overnight studies in severe chronic left heart failure: arrhythmias and oxygen desaturation. *Br Heart J* 1991;65:77-83.
- 43 Cripps T, Rucker G, Stradling J. Nocturnal hypoxia and arrhythmias in patients with impaired left ventricular function. *Br Heart J* 1992;68:382-6.
- 44 Grover RF, Wagner WW Jr, McMurty IF, Reeves JT. Pulmonary circulation. In: Shepherd JT, Abboud FM, Geiger SR, eds. *Handbook of physiology, section 2: The cardiovascular system, vol 3*. Bethesda: American Physiological Society, 1983:103.
- 45 Fowler KT, Read J. Effect of alveolar hypoxia and zonal distribution of pulmonary blood flow. *J Appl Physiol* 1963;18:244-50.
- 46 Lindsay DA, Read J. Pulmonary vascular responsiveness in the prognosis of chronic obstructive lung disease. *Am Rev Respir Dis* 1972;105:242-50.
- 47 Vogel JHK, Weaver WF, Rose RL, Blount SG Jr, Grover RF. Pulmonary hypertension on exertion in normal man living at 10,150 feet (Leadville, Colorado). In: Grover RF, ed. *Normal and abnormal pulmonary circulation*. Basel: S Karger, 1963:269-85.
- 48 Nocturnal Oxygen Therapy Trial Group. Continuous or nocturnal oxygen therapy in hypoxemic chronic obstructive lung disease. *Ann Intern Med* 1980;93:391-8.
- 49 M.R.C. Working Party. Long term domiciliary oxygen therapy in chronic hypoxic cor pulmonale complicating chronic bronchitis and emphysema. *Lancet* 1981;i:681-6.
- 50 Wilkinson M, Langhorne CA, Heath D, Barer GR, Howard P. A pathophysiological study of 10 cases of hypoxic cor pulmonale. *Q J Med* 1988;66:65-87.
- 51 Magee F, Wright JL, Wiggs BR, Pare PD, Hogg JC. Pulmonary vascular structure and function in chronic obstructive pulmonary disease. *Thorax* 1988;43:183-9.
- 52 Williams IP, Boyd MJ, Humberstone AM, Wilson AG, Millard FJC. Pulmonary arterial hypertension and emphysema. *Br J Dis Chest* 1984;78:211-6.
- 53 Segel NM, Bishop J. The circulation in patients with chronic bronchitis and emphysema at rest and during exercise with special reference to the influence of changes in blood viscosity and blood volume on the pulmonary circulation. *J Clin Invest* 1966;45:1555-68.
- 54 Wallis PJW, Skehan JD, Newland AC, Wedzicha JA, Mills PG, Empey DW. Effects of erythrapheresis on pulmonary haemodynamics and oxygen transport in patients with secondary polycythaemia and cor pulmonale. *Clin Sci* 1986;70:91-8.
- 55 Farber MD, Roberts LR, Weinberger MH, Robertson GL, Fineberg MS, Manfredi F. Abnormalities of sodium and H₂O handling in chronic obstructive lung disease. *Arch Intern Med* 1982;142:1326-30.
- 56 Richens JM, Howard P. Oedema in cor pulmonale. *Clin Sci* 1982;62:255-9.
- 57 Berbel LN, Miro RE. Pulmonary hypertension in the pathogenesis of cor pulmonale. *Cardiovasc Rev* 1983;4:359-63.