LETTERS TO THE EDITOR

● *The British Heart Journal welcomes letters commenting on papers that it has published within the past six months.*

● *All letters must be typed with double spacing and signed by all authors.*

● *No letter should be more than 600 words.*

● *In general, no letter should contain more than six references (also typed with double spacing).*

Acronym aggravation

SIR,—I enjoyed reading the editorial on insulin and atherosclerosis by Savage and Saad.¹ Unfortunately this otherwise excellent article was marred by the fact that the authors used the acronym CARDIA without explaining it.

Your Notice to Contributors clearly says that abbreviations should not be used in the text. The same rule should apply to acronyms. According to the International

Committee of Medical Journal Editors² the full term for which an abbreviation or an acronym stands should precede its first use in the text.

Perhaps Savage and Saad justified not explaining the acronym CARDIA by referring to an earlier publication, their reference 20. Unfortunately there was no explanation of CARDIA in that reference either.³ The authors of reference 3 referred to another article⁴ in which I finally found the derivation of the acronym. CARDIA is an approximate acronym for Coronary Artery (Disease) Risk Development In (Young) Adults.⁴

Physicians, especially cardiologists, like to use or invent acronyms. But unless they are explained they lead to frustration and sometimes confusion. Acronyms are often necessary but can sometimes be quite frustrating if you do not know what they stand for.⁵ That was why I recently prepared a list of acronyms of major cardiological trials (table),⁶ which is currently being updated. Acronyms can sometimes cause confusion, because several trials share the same acronyms—for example CATS, PACT, and TIPE.⁶

TSUNG O CHENG
Division of Cardiology,
George Washington University Medical Center,
2150 Pennsylvania Avenue NW,
Washington DC, USA

1 Savage PJ, Saad MF. Insulin and atherosclerosis: villain, accomplice, or innocent bystander? *Br Heart J* 1993;69:473-5.

- 2 International Committee of Medical Journal Editors: Uniform requirements for manuscripts submitted to biomedical journals. *N Engl J Med* 1991;324:424-8.
- 3 Manolio TA, Savage PJ, Burke GL, Liu K, Wagenknecht LE, Sidney S, Jacobs DR Jr, Roseman JM, Donahue RP, Oberman A. Association of fasting insulin with blood pressure and lipids in young adults. The CARDIA study. *Arteriosclerosis* 1990;10:430-6.
- 4 Friedman GD, Butter GR, Donahue RP, Hughes GH, Hulley SB, Jacobs DR Jr, Liu K, Savage PJ. CARDIA: Study design, recruitment, and some characteristics of the examined subject. *J Clin Epidemiol* 1988;41:1105-16.
- 5 Brown JD. Acronymy. *Med J Aust* 1992;157:356.
- 6 Cheng TO. Acronyms of major cardiological trials. *Am J Cardiol* 1992;70:1512-4.

We apologise for not following the recommended practice of spelling out all acronyms at the first mention in Savage and Saad's editorial. Many of the articles that are submitted to the British Heart Journal for publication use acronyms without explanation. Like the authors of these articles and many speakers at meetings we share a tendency to drift away from best practice. We thank Dr Cheng for drawing attention to our lapse and for supplying overwhelming evidence (in his list reproduced with permission of the American Journal of Cardiology 1992;70:1512-4) why authors and speakers must spell out all acronyms at the first mention.—EDITOR

Acronyms of Major Cardiological Trials

ACME (Angioplasty Compared to Medicine)
AFASAK (Atrial Fibrillation, Aspirin, Antikoagulation)
AFTER (Anistreplase Following Thrombolysis Effect on Reocclusion)
AICLA (Accidents Ischémiques Cérébraux Liés à l'Atherosclérose)
AIMS (APSAC Intervention Mortality Study)
AIRE (Acute Infarction Remipril Efficacy)
AITIA (Aspirin in Transient Ischemic Attacks)
AMIS (Aspirin in Myocardial Infarction Study)
AMPI (APSAC in Acute Myocardial Infarction Placebo-Controlled Investigation)
ANBP (Australian National Blood Pressure Trial)
APRICOT (Antithrombotics in Prevention of Reocclusion in Coronary Thrombolysis)
APRICOT (Aspirin vs Coumadin Trial)
APSI (Acebutalol Prevention Secondaire de l'Infarctus)
APSIM (APSAC dans l'Infarctus du Myocarde)
APSI (Angina Prognosis Study with Isoptin and Seloken Trial)
ARIC (Atherosclerosis Risk in Communities Study)
ARIS (Anturane Reinfarction Italian Study)
ARMS (APSAC Reocclusion Multicenter Study)
ART (Anturane Reinfarction Trial)
ASK (Australian SK Trial in Stroke)
ASPECT (Anticoagulants in Secondary Prevention of Events in Coronary Thrombolysis)
ASPS (Australian Swedish Pirodolo Study)
ASSET (Anglo-Scandinavian Study of Early Thrombolysis)
ATACS (Antithrombotic Therapy in Acute Coronary Syndromes)
ATEST (Atenolol and Streptokinase Trial)
ATIAIS (Anturane Transient Ischemic Attack Italian Study)
BAATF (Boston Area Anticoagulation Trial for Atrial Fibrillation)
BARI (Bypass Angioplasty Revascularization Investigation)
BEST (Beta-Blocker Stroke Trial)
BHAT (Beta-Blocker Heart Attack Trial)
BIRNH (Belgian Interuniversity Research on Nutrition and Health)
BRHS (British Regional Heart Study)
BWIS (Baltimore Washington Infant Study)
CABRI (Coronary Artery Bypass Revascularization Investigation)
CAFA (Canadian Atrial Fibrillation Anticoagulation Study)
CAMIAT (Canadian Amiodarone Myocardial Infarction Arrhythmia Trial)
CAPPHY (Captopril Primary Prevention in Hypertension)
CAPRIE (Clopidogrel vs Aspirin in Patients at Risk of Ischemic Events)
CAPS (Cardiac Arrhythmia Pilot Study)

CARDIA (Coronary Artery Risk Development in Young Adults)
CARPORT (Coronary Artery Restenosis Prevention On Repeated Thromboxane-Antagonism)
CASH (Cardiac Arrest Study Hamburg)
CASH (Canadian Amlodipine-Atenolol in Silent Ischemia Study)
CASS (Coronary Artery Surgery Study)
CAST (Cardiac Arrhythmia Suppression Trial)
CATS (Canadian American Ticlopidine Study)
CATS (Captopril and Thrombolysis in Myocardial Infarction Study)
CDP (Coronary Drug Project)
CECCC (Confidential Enquiry into Cardiac Catheterization Complications)
CEDIM (Italian Study on L-Carnitine and Digital Echocardiography in Myocardial Infarction)
CITO (Collaborazione Italiana per la Trombosi in Ortopedia)
CLAS (Cholesterol Lowering Atherosclerosis Study)
CLIP (Cholesterol Lowering Intervention Program)
CONSENSUS (Cooperative North Scandinavian Enalapril Survival Study)
CORIS (Coronary Risk Factor Study)
CRAFT (Catheterization Rescue Angioplasty Following Thrombolysis)
DART (Diet and Reinfarction Trial)
DAVIT (Danish Verapamil in Myocardial Infarction Trial)
DIMT (Dutch Iloperidine Multicenter Trial)
DRS (Diltiazem Reinfarction Study)
DUCCS (Duke University Clinical Cardiology Studies)
Dutch IRS (Dutch Invasive Reperfusion Study)
DVT (Danish Verapamil Trial)
EARS (European Atherosclerosis Research Study)
EAST (Emory Angioplasty Surgery Trial)
ECCOMAC (European Concerted CoMmunity Action Programmes)
ECAT (European Concerted Action on Thrombosis and Disabilities)
ECATAP Trial (ECAT Angina Pectoris Trial)
ECTIM (Etude Cas-Témoins sur l'Infarctus du Myocarde)
ECSS (European Cooperative Study Group)
ECSS (European Coronary Surgery Study)
ECST (European Carotid Surgery Trial)
EIS (European Infarction Study)
ELCA Registry (Excimer Laser Coronary Angioplasty Registry)
EMERAS (Estudio Multicenter Estreptoquinasa Republicas Americas Sud)
EMIAT (European Myocardial Infarction Amiodarone Trial)
EMIP (European Myocardial Infarction Project)
EMPAP (Enoxaparine Maxipa Prevention of Angioplasty Restenosis)
EMS (European Multicenter Study)
EPPI (Etude de Prescription Post Infarctus)
EPSIM (Enquete de Prevention Secondaire de l'Infarctus du Myocarde)

- ERICA (European Risk and Incidence, a Coordinated Analysis)
 ESPS (European Stroke Prevention Study)
 ESVM (Electrophysiologic Study vs Electrocardiographic Monitoring)
 ETDRS (Early Treatment Diabetic Retinopathy Study)
 EVA (European Vascular Agency)
 EVAS-IM (Etude Vaudoise APSAC vs Streptokinase dans l'infarctus du Myo-
 carde)
 EXCEL (Expanded Clinical Evaluation of Lovastatin)
 ÉWPHE (European Working Party on Hypertension in the Elderly)
 FACET (Flosequinan ACE-Inhibitor Trial)
 FATS (Familial Atherosclerosis Treatment Study)
 FHS (Family Heart Study)
 FIPS (Frankfurt Isoptin Progression Study)
 FRESH (Food Reeducation Elementary School Health Study)
 GABI (German Angioplasty Bypass Investigation)
 GAMIS (German Austrian Myocardial Infarction Study)
 GAUS (German Activator Urokinase Study)
 GCP (German Cardiovascular Prevention Study)
 GEMT (German Eminase Multicenter Trial)
 GISSI (Gruppo Italiano per lo Studio della Streptochinasi nell'Infarto Mio-
 cardico)
 GMT (Göteborg Metoprolol Trial)
 GPP (Göteborg Primary Prevention Trial)
 GRASP (Glaxo Restenosis and Symptoms Project)
 GREAT (Grampian Region Early Antistreptase Trial)
 GRECO (German Study with Recombinant t-PA in Coronary Conclusion)
 GUIDE Trial (Guidance by Ultrasound Imaging for Decision Endpoints
 Trial)
 GUSTO (Global Utilization of Streptokinase and t-PA for Occluded Arter-
 ies)
 HALS (Heart Attacks in London Study)
 HAPPHY (Heart Attack Prevention in Primary Hypertension)
 HART (Heparin Aspirin Reperfusion Trial)
 HDPF (Hypertension Detection and Follow-Up Program)
 HHS (Helsinki Heart Study)
 HINT (Holland Interuniversity Nifedipine/Metoprolol Trial)
 HYNON (Hypertension Non-Drug Treatment Cooperative Study)
 ICIN (Intracoronary Streptokinase Trial of the Interuniversity Cardiology
 Institute of the Netherlands)
 IMAGE (International Metoprolol/Nifedipine Angina Exercise Trial)
 IMPACT (International Mexiletine & Placebo Antiarrhythmic Coronary
 Trial)
 INCLIN (International Clinical Epidemiology Network)
 INTACT (International Nifedipine Trial on Antiatherosclerotic Therapy)
 INTERSALT (International Studies of Salt and Blood Pressure)
 IPPPSH (International Prospective Primary Prevention Study on Hyper-
 tension)
 IRS (Invasive Reperfusion Study)
 ISAM (Intravenous Streptokinase in Acute Myocardial Infarction)
 ISIS (International Study of Infarct Survival)
 KAMIT (Kentucky Acute Myocardial Infarction Trial)
 LAPIS (Late Potentials in Myocardial Infarction Study)
 LATE (Late Assessment of Thrombolytic Efficacy)
 LIMIT (Leicester Intravenous Magnesium Intervention Trial)
 LIMITS (Liquemin in Myocardial infarction During Thrombolysis with Sa-
 rurapase)
 LIT (Lopressor Intervention Trial)
 LRC-CPPT (Lipid Research Clinics-Coronary Primary Prevention Trial)
 MAPHY (Metoprolol Atherosclerosis Prevention in Hypertension)
 MARCATOR (Multicenter American Research Trial with Cilazapril After
 Angioplasty to Prevent Transluminal Coronary Obstruction & Resteno-
 sis)
 MARS (Monitored Atherosclerosis Regression Study)
 MAST-I (Multicenter Acute Stroke Trial—Italy)
 MDPI (Multicenter Diltiazem Post Infarction Trial)
 MEHP (Metoprolol in Elderly Hypertensive Patients)
 MELODHY (Metoprolol Low Dose in Hypertension)
 MERCATOR (Multicenter European Research Trial with Cilazapril After
 Angioplasty to Prevent Transluminal Coronary Obstruction & Resteno-
 sis)
 M-HEART (Multi-Hospital Eastern Atlantic Restenosis Trial)
 MIAMI (Metoprolol in Acute Myocardial Infarction)
 MIDAS (Myocardial Infarction Data Acquisition System)
 MILESTONE (Multicenter iloprost European Study on Endangitis)
 MILIS (Multicenter Investigation of the Limitation of Infarct Size)
 MITI (Myocardial Infarction Triage and Intervention)
 MONICA (Monitoring Trends and Determinants in Cardiovascular Dis-
 ease)
 MPRG (Multicenter Postinfarction Research Group)
 MRFIT (Multiple Risk Factor Intervention Trial)
 NACI Registry (New Approaches to Coronary Intervention Registry)
 NAMIS (Nifedipine Angina Myocardial Infarction Study)
 NASCET (North American Symptomatic Carotid Endarterectomy Trial)
 NCEP (National Cholesterol Education Program)
 NGHS (NHLBI Growth and Health Study)
 NHANES (National Health and Nutrition Examination Survey)
 OCSF (Oxfordshire Community Stroke Project)
 OSIRIS (Optimization Study of Infarct Reperfusion Investigated by ST-
 Monitoring)
 PACK (Prevention of Atherosclerotic Complications with Ketanserin)
 PACT (Pre-Hospital Application of Coronary Thrombolysis)
 PACT (Pro-Urokinase in Acute Coronary Thrombolysis)
 PACTE (Prevention des Accidents Thrombo-Emboliques Chez les Por-
 teurs de Protheses Valvulaires Cardiaques)
 PAIMS (Plasminogen Activator Italian Multicenter Study)
 PAMI (Primary Angioplasty Myocardial Infarction Trial)
 PARIS (Persantine Aspirin Reinfarction Study)
 PARK (Prevention of Angioplasty Reocclusion with Ketanserin)
 PARTNER (Peripheral Arterial Disease Response to Taprostene with New
 Established Response Criteria)
 PASS (Practical Applicability of Saruplase Study)
 PATS (Prehospital Administration of t-PA Study)
 PDAY (Pathobiological Determinants of Atherosclerosis in Youth)
 PIOPED (Prospective Investigation of Pulmonary Embolism Diagnosis)
 PLAC (Pravastatin Limitation of Atherosclerosis in Coronary Arteries)
 POSCH (Program on Surgical Control of Hyperlipidemia)
 PREMIS (Prehospital Myocardial Infarction Study)
 PRIMI (Prourokinase in Myocardial Infarction)
 PROCAM Study (Prospective Cardiovascular Münster Study)
 PROFILE (Prospective Randomized Flosequinan Longevity Evaluation)
 PROMISE (Prospective Randomized Milrinone Survival Evaluation)
 QUIET (Quinapril Ischemic Events Trial)
 RADIANCE (Randomized Assessment of Digoxin and Inhibitors of Angio-
 tensin Converting Enzyme)
 RAAMI (Rapid Administration of Alteplase in Myocardial Infarction)
 REPAIR (Reperfusion in Acute Infarction, Rotterdam)
 RESCUE (Randomized Evaluation of Salvage Angioplasty with Combined
 Utilization of Endpoints)
 RISK (Regional Study av Instabil Kranskärslssjukdom)
 RITA (Randomized Intervention Treatment of Angina)
 ROBUST (Recanalization of Occluded Bypass Graft, Urokinase Study
 Trial)
 ROCKET (Regionally Organized Cardiac Key European Trial)
 SAFE (Safety After Fifty Evaluation)
 SALT (Swedish Aspirin Low-Dose Trial)
 SAMIT (Streptokinase Angioplasty Myocardial Infarction Trial)
 SAVE (Survival and Ventricular Enlargement)
 SCATI (Studio sulla Calciparina nell'Angina e nella Trombosi Ventricolare
 nell'Infarto)
 SCRIP (Stanford Coronary Risk Intervention Project)
 SEPIVAC (Studio Epidemiologica sull'Incidenza delle Vasculopatie Acute
 Cerebrali)
 SESAM (Study in Europe of Saruplase and Alteplase in Myocardial Infarc-
 tion)
 SHAVE (Steerable Housing for Atherovascular Excision)
 SHEP (Systolic Hypertension in the Elderly Program)
 SIAM (Streptokinase in Acute Myocardial Infarction)
 SMISS (Silent Myocardial Ischemia Stress Study)
 SMT (Stockholm Metoprolol Trial)
 SOLVD (Studies of Left Ventricular Dysfunction)
 SPAF (Stroke Prevention in Atrial Fibrillation)
 SPINAF (Stroke Prevention in Nonrheumatic Atrial Fibrillation)
 SPRINT (Secondary Prevention Reinfarction Israeli Nifedipine Trial)
 SRT (Sorbini Retinopathy Trial)
 SSSS (Scandinavian Simvastatin Survival Study)
 STAI (Study Ticlopidine in Angor Instabile)
 STAMP (Systemic Thrombolysis in Acute Myocardial Infarction with
 Prourokinase)
 STARS (St. Thomas' Atherosclerosis Regression Study)
 STEP (Study of Taprostene in Elective PTCA)
 STIMS (Swedish Ticlopidine Multicentre Study in Patients with Intermit-
 tent Claudication)
 STOP Hypertension (Swedish Trial in Old Patients with Hypertension)
 SWIFT (Should We Intervene Following Thrombolysis)
 TACS (Thrombolysis and Angioplasty in Cardiogenic Shock)
 TACT (Ticlopidine vs Placebo for Prevention of Acute Closure Trial)
 TAMI (Thrombolysis and Angioplasty in Myocardial Infarction)
 TAPS (t-PA APSAC Patency Study)
 TASS (Ticlopidine Aspirin Stroke Study)
 TAUSA (Thrombolysis and Angioplasty in Unstable Angina)
 TEAHAT (Thrombolysis Early in Acute Heart Attack Trial)
 TEAM (Trial of Eminase vs Alteplase in Myocardial Infarction)
 TIARA (Timolol en Infarto Agudo, Republica Argentina)
 TIBET (Total Ischemic Burden European Trial)

TICO (Thrombolysis in Coronary Occlusion)
 TIMAD (Ticlopidine in Micro-Angiopathy of Diabetes)
 TIMI (Thrombolysis in Myocardial Infarction)
 TIPE (Thrombolysis in Pulmonary Embolism Study)
 TIPE (Thrombolysis in Peripheral Embolism Patient Study)
 TOHMS (Trial of Hypertensive Medications Study)
 TOHP (Trials of Hypertension Prevention)
 TOMHS (Treatment of Mild Hypertension Study)
 TOP (Thrombolysis in Old Patients)
 TPAT (Tissue Plasminogen Activator, Toronto)
 TPT (Thrombosis Prevention Trial)
 TRENT (Trial of Early Nifedipine Treatment in Acute Myocardial Infarction)

TRUST (Trial of United Kingdom for Stroke Treatment)
 UNASEM (Unstable Angina Study Using Eminase)
 UNSA (Unstable Angina Study)
 UPET (Urokinase Pulmonary Embolism Trial)
 URALMI (Urokinase and Alteplase in Myocardial Infarction)
 USIM (Urokinasi per via Sistemica nell'Infarto Miocardico)
 VACA Registry (Valvuloplasty and Angioplasty of Congenital Anomalies Registry)
 V-HeFT (Veterans Heart Failure Trial)
 WARIS (Warfarin Reinfarction Study)
 WHA Study (Worcester Heart Attack Study)
 WWICT (Western Washington Intracoronary Streptokinase Trial)
 WWIST (Western Washington Intravenous Streptokinase Trial)

Monitoring myocardial damage in cardiac surgery by troponin T detection

SIR,—Perioperative myocardial injury remains the most common cause of death in cardiac surgery. The need for new diagnostic criteria to assess the comparative efficacy of different myocardial protection techniques prompted us to identify reliable markers of myocardial necrosis. Katus *et al* reported that the serum concentration troponin T, a cardiospecific protein, reliably detects myocardial cell necrosis in patients undergoing myocardial revascularisation.¹ We assayed troponin T (ELISA, Troponin T, Boehringer Mannheim) in 40 consecutive patients of whom 34 underwent coronary artery bypass grafting (CABG), four mitral valve replacement (MVR), and two aortic valve replacement (AVR). Myocardial protection was accomplished by antegrade-retrograde blood cardioplegia according to the method described by Buckberg.² No perioperative deaths occurred. Using the same electrocardiographic (ECG) criteria for perioperative myocardial infarction described by Katus *et al* we found two cases among the CABG patients and one among the AVR patients. Troponin T concentrations were lower than 0.1 µg/l in all preoperative samples and rose after surgery in the three patients with perioperative myocardial infarction to a peak value of 6.06 µg/l, 11.67 µg/l, and 28.75 µg/l respectively. These results accord with those of Katus *et al* who reported a troponin T median peak value of 11 µg/l (range 6–31 µg/l) in patients in whom ECG signs of perioperative myocardial infarction developed after CABG. In patients with no evidence of myocardial infarction after surgery troponin T release was significantly lower (mean (SD) 0.96 (0.98) µg/l, median 0.68 µg/l, range 0.26–4.6 µg/l) than that in patients with perioperative myocardial infarction. Surprisingly, these values were also lower than that reported by Katus *et al* in a similar subgroup (median 5 µg/l, range 1.3–11 µg/l) who were cardioprotected by a Bretschneider HTK cardioplegic solution.³ Because there were no apparent differences in the duration of cardiopulmonary bypass and aortic cross-clamping or the number of diseased and grafted coronary vessels we suggest that the reduced troponin T release seen in our patients was due to the different myocardial protection protocol. Therefore troponin T seems to be a highly specific and sensitive marker for myocardial cell necro-

sis, that is also useful for assessing the efficiency of myocardial protection techniques.

MICHELE TRIGGIANI

ALBERTO DOLCI*

FRANCESCO DONATELLI

ADALBERTO GROSSI

*Institute for Cardiovascular and Respiratory Diseases, University of Milan and *Protein Unit, Department of Medicine Laboratory, Scientific Institute HS Raffaele, Via Olgettina 60, 20132 Milan, Italy*

- 1 Katus HA, Schoepenthan M, Tanzeem A, Bauer HG, Saggau W, Diederich KW, *et al*. Non-invasive assessment of perioperative myocardial cell damage by circulating cardiac troponin T. *Br Heart J* 1991;65: 259–64.
- 2 Buckberg GD. Antegrade-retrograde blood cardioplegia to ensure cardioplegic distribution: operative techniques and objectives. *J Card Surg* 1989;4:216–38.
- 3 Gebhard MM, Bretschneider HJ, Gersing E, Preusse CJ, Schnabel PA, Ulbricht LJ. Calcium-free cardioplegia. *Eur Heart J* 1983;4:151–60.

BRITISH CARDIAC SOCIETY NEWSLETTER

The challenges that lie ahead in staffing cardiac units into the twenty first century, which were the subject of a meeting last November, require continuous debate. Both the Specialist Advisory Committee and the Manpower and Training Committee are developing strategies to improve our training programmes and ensure that staffing levels are appropriate.

The working party on cardiology in district general hospitals, chaired by Andrew McLeod, is contributing to this debate. All the indications are that nearly double the number of consultants will be required over the next decade and it remains to be seen how the funding will be provided. The formal government response to the Calman report after a period of consultation is awaited and will no doubt precipitate vigorous debate. Harmonising our training programmes with the rest of Europe is not going to be easy and cannot be achieved overnight.

The Specialist Advisory Committee is reviewing the content of training, which will

become more structured with formal guidelines and a formal assessment of training of trainees being introduced.

Archives

Arthur Holman, who was appointed by Council, writes: "The main task of the archivist is the preservation and proper arrangement of the Society's records. These will include: minutes of Council meetings; financial accounts; minutes of annual general meetings; records of the scientific meetings and programme books; membership records; minutes of Officers' meetings; records and meetings of associated groups; and correspondence. Advice is being sought from a professional archivist on how these records should be kept and indexed, with special reference to computer management.

Our important need to have a complete set of the *British Heart Journal* from its foundation in 1939 has been met by a most welcome gift from Richard Emanuel. He has given us the bound volumes of the journal that belonged to his father, Professor J G Emanuel, and we are deeply grateful to him for his generosity. We now have to obtain a set of *Cardiovascular Research*.

In addition, I intend to establish a small library of books that will cover the development of cardiology from the mid-nineteenth century to the present day. If possible we would also like to have a small collection of historical instruments both diagnostic and therapeutic—for example, the Mackenzie polygraph and the mitral valve dilator.

If members have books, instruments, or other items of historical interest that they would like to donate to the Society I will be most grateful if they will get in touch with me either at the Society's office or at my home: Seabank, Chick Hill, Pett, Hastings, East Sussex TN35 4EQ (tel: 0424 813228)."

Data Management Committee: progress on the Read Clinical Terms

Malcolm Towers writes: "Our lists of diagnostic (clinical) terms in congenital heart disease and "adult" heart disease and terms for special investigations—such as electrocardiography, electrophysiology, and nuclear cardiology—have been completed. The list of terms for echocardiography is almost complete. The Centre for Coding and Classification now has the