



Editor's choice
Scan to access more
free content

New insights on diabetes mellitus and obesity in Africa—Part 1: prevalence, pathogenesis and comorbidities

Andre Pascal Kengne,^{1,2,3} Justin-Basile Echouffo-Tcheugui,⁴ Eugene Sobngwi,^{5,6} Jean-Claude Mbanya^{5,7}

¹Department of Medicine, University of Cape Town and South African Medical Research Council, Cape Town, South Africa

²The George Institute for Global Health, Sydney, Australia

³Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht, The Netherlands

⁴Hubert Department of Global Health, Rollins School of Public Health, Emory University, Atlanta, Georgia, USA

⁵Yaounde Central Hospital and Department of Internal Medicine and Specialties, Faculty of Medicine and Biomedical Sciences, National Obesity Centre, University of Yaoundé 1, Yaoundé, Cameroon

⁶Institute of Health and Society, Newcastle University, Newcastle, UK

⁷University of Technology, Kingston, Jamaica

Correspondence to

Dr Andre Pascal Kengne, National Collaborative Programme on Cardiovascular and Metabolic Disease, South African Medical Research Council, PO Box 19070 Tygerberg, Cape Town 7505, South Africa; andre.kengne@mrc.ac.za

Bongani Mayosi is the Guest Editor of this Cardiology in Africa review series.

Received 5 November 2012

Revised 4 February 2013

Accepted 25 February 2013

Published Online First

16 May 2013



► <http://dx.doi.org/10.1136/heartjnl-2013-304130>

To cite: Kengne AP, Echouffo-Tcheugui J-B, Sobngwi E, et al. *Heart* 2013;**99**:979–983.

ABSTRACT

Evidence continues to accumulate on the rising burden of diabetes mellitus at a higher pace in Africa. In a series of two papers, we sought to summarise recent evidence on diabetes and obesity in Africa based on a systematic review of studies published between January 2002 and October 2012. This first paper on the prevalence, pathogenesis and comorbidities shows that the increase in diabetes prevalence has paralleled that of obesity in Africa. Recent surveys on diabetes and obesity have been largely suboptimal. Hence, the need for more representative and robust continent-wide prevalence figures, which may be somehow achieved through pooling of existing data. Prospective studies linking environmental risk factors to disease occurrence and outcomes remain scarce, and genetic factors for diabetes or obesity have not been extensively assessed. The health consequences of diabetes are manifold, and include a complex interaction with other conditions like HIV infection and sickle cell disease/trait.

INTRODUCTION

The importance of diabetes mellitus as a major threat to the health of populations in Africa is increasingly being recognised, partly as the result of the growing global interest for chronic diseases in general, but also of the increasing availability of evidence to support the case of diabetes in Africa.¹ Estimates from the International Diabetes Federation (IDF) for the year 2012 indicate that people with diabetes comprise about 371 million individuals.² About 7.5% (27.5 million) of this population reside in Africa including 14.7 million in sub-Saharan Africa (SSA) and 12.8 million in North Africa and Sudan. IDF's projections suggest that the global number of people with diabetes will increase by 51% to 552 million by 2030. The largest expected relative growth will occur in Africa, which by then will be home to about 49.7 million people with diabetes (81% relative increase), translating into about 9% of the global population with diabetes.²

The rapid increase in diabetes figures in Africa is largely attributed to accelerated environmental changes characterised by rapid urbanisation and modernisation, as well as ageing of the population. The underlying diabetes-promoting behavioural changes include the adoption of sedentary lifestyles and unhealthy eating habits. Together with the ensuing overweight and obesity, these result in many people developing diabetes and other risk factors for cardiovascular diseases. Globally, overweight and obesity have been recognised as the main

drivers of the growing epidemic of diabetes. Between 1980 and 2008, the body mass index (BMI) of the adult population in the world increased by about 0.4 kg/m² per 10 years in men and 0.5 kg/m² in women.³ During this same time interval, the rates of change in BMI in men and women across SSA and North Africa were always higher than the global average.³ The average glucose levels of the population in Africa have mostly remained low within this period, likely reflecting the time lag needed before the increasing BMI can have a sizable effect on the prevalence of diabetes mellitus.⁴

In this series of two papers, we provide a critical review of recent evidence pertaining to diabetes mellitus (type 2 diabetes in particular) and obesity in Africa. More than a simple update of existing reviews that have addressed the subject matters from a SSA or a Northern African perspective,^{5–7} we use a broader African perspective to summarise recently published evidence. Where relevant, we contrast the evidence by region, and identify the research gaps, remaining challenges, and existing opportunities to improve diabetes and obesity prevention and control in Africa. The first article of the series focuses on the prevalence, pathogenesis and comorbidities of obesity and diabetes, using the most relevant evidence published over the last 10 years. The sources of the evidence and approaches for gathering them are presented in detail in the second paper.⁸

PREVALENCE AND INCIDENCE OF DIABETES AND OBESITY

Prevalence studies of diabetes

A total of 13 community-based prevalence studies have been published over the last 10 years from nine SSA countries and Tunisia (table 1). The overall prevalence of diabetes (age-standardised only in two studies^{9 10}) ranged from 2.8% in rural Angola¹¹ to 28.2% among urban mixed-ancestry South Africans.⁹ The proportion of screen-detected cases (reported in five studies) among those with diabetes ranged from 21.3% in rural Congo (Democratic Republic) to 90% in urban Senegal.¹² Sex-specific prevalence was reported in seven studies, showing higher rates in men than women in three studies^{11 13 14} and the reverse in four others.^{10 12 15 16} Two studies provided urban versus rural comparisons in the same countries, both showing urban prevalence of diabetes about twice that observed in rural settings.^{17 18} Three of the studies used self-selected sample of participants,^{15 19 20} two studies from South Africa^{9 10} and one from Angola¹¹ used oral glucose tolerance

Table 1 Selected population-based prevalence studies on diabetes in Africa, 2002–2012

Study	Country	Sample	Sampling	Setting	Age (years)	Diagnosis	Prevalence of diabetes (%)			
							Overall	Men	Women	Undiagnosed
Evaristo-Neto ¹¹	Angola	421	Random	Rural	30–69	OGTT	2.8	3.2	2.7	–
Echouffo-Tcheugui ¹⁵	Cameroon	1591	Self-selected	Urban	43.7	FBG	15.3	13.7	17.0	6.3
Katchunga ¹⁷	DRC	424	Random	Urban	53.3	RBG	4.7	–	–	1.0
Katchunga ¹⁷	DRC	245	Random	Rural	58.5	RBG	2.9	–	–	1.6
Ploubidis ¹⁸	Kenya	2959	Random	Rural	64.7	RBG	5.1	–	–	–
Ploubidis ¹⁸	Kenya	1437	Random	Urban	60.8	RBG	10.1	–	–	–
Ejim ¹³	Nigeria	858	Random	Rural	59.8	FBG	4.4	7.3	3.3	–
Duboz ¹²	Senegal	600	Random	Urban	>20	FBG	17.9	14.0	21.8	16.2
Peer ¹⁰	South Africa	1099	Random	Urban	43.3	OGTT	12.1 (13.1)	10.2 (11.3)	13.8 (14.7)	4.9
Erasmus ⁹	South Africa	642	Random	Urban	50.9	OGTT	28.2 (26.3)	–	–	18.1
Baragou ¹⁶	Togo	2000	Random	Urban	39	2 FBG	7.3	6.9	7.3	–
Nsakashalo-Senkwe ⁵⁴	Zambia	1928	Random	Urban	≥25	FBG	2.7	–	–	–
Hammami ¹⁴	Tunisia	598	Random	Urban and rural	72.3	RBG	27.4	29.2	26.5	–
Chamie ²⁰	Uganda	2283	Self-selected	Urban	35	RBG	3.5	–	–	0.8

Figures within parenthesis are age-adjusted.

FBG, fasting blood glucose; OGTT, oral glucose tolerance test; RBG, random blood glucose.

test to diagnose diabetes, while the remaining used random or fasting capillary glucose measurement (table 1).

In general, the quality of the available prevalence studies of diabetes remains suboptimal, such that in many settings these studies do not provide accurate estimates of the magnitude of the condition. Over the last 10 years, several African countries have made significant progress by embracing the WHO STEPwise approach to chronic disease surveillance (STEPS), which involves repeated population-based data collection on the levels of risk factors.²¹ At least 20 countries in Africa have conducted one such survey.²² Some surveys have collected data to assess diabetes status, usually based on self-reports and fasting capillary glucose in a subsample of the surveyed population. Very few of the STEPS survey data have been appropriately analysed and published to allow regional comparisons.²² However, these surveys used similar methodological approaches to sampling and data collection, which offers an opportunity for refining and updating prevalence data for diabetes and other risk factors, through pooling of individual countries' data to generate more reliable regional and continent-wide figures. Such an approach has been used for instance for physical activity.²² Making the existing STEPS data publicly available may foster collaboration within Africa and beyond to maximise its exploitation.

Studies of incident diabetes and risk factors

Risk factors for diabetes in Africa have been investigated mostly using the cross-sectional design, which has the limitation of not establishing the sequence of events.⁶ A recent overview of longitudinal studies conducted in SSA,²³ with relevance to cardiovascular disease and diabetes, showed that the few available prospective studies found the main determinants of incident diabetes to be baseline glucose tolerance status and BMI.²³ In a more recent cohort study among people aged >65 years in Nigeria, urban residence and high socioeconomic status were found to be the main risk factors of self-reported incident diabetes.²⁴ A few cohort studies are currently underway in Africa; these include the Prospective Urban Rural Epidemiology Study (African sites)²⁵ and the Bellville-South cohort in South Africa.⁹ Preliminary data from the Bellville-South cohort show rates of abnormal glucose tolerance to be very high among

mixed-ancestry populations, and determined essentially by the baseline cardiometabolic risk profile.²⁶

Prevalence studies of obesity

About 20 prevalence studies on obesity in African populations were published over the last decade, with nearly half of them originating from Nigeria (table 2). The age bracket of the included population suggests an increasing interest for childhood and adolescent obesity in Africa, with over half of the studies conducted in the paediatric population. Childhood obesity on its own is a broad and complex topic and is not discussed further in the current review. Available studies are also characterised by a low representation of rural populations. These studies have reported prevalence figures for obesity in adults ranging from 5% in rural Uganda²⁷ to 30% in Rural Nigeria.¹³ The frequency of obesity is almost always higher in female subjects than in male subjects (table 2). When reported, the prevalence of overweight across studies was most often higher than that for obesity, with differentially higher figures for women in some studies, but not all. Very few studies reported the prevalence of abdominal obesity, which ranged from 7.5%²⁸ to 66.5%²⁹ among adults in urban Nigeria. There was no North-Africa versus SSA pattern in the profile of published studies, partly due to the small number of studies from Northern Africa. Cross-sectional surveys that used WHO STEPS approach have also reported a variable crude prevalence of obesity ranging from 2.2% in a nationwide survey in Madagascar to 25.1% in Seychelles.

PATHOGENESIS OF DIABETES AND ROLE OF OBESITY

The increase in the prevalence of type 2 diabetes appears to be closely linked to the upsurge in obesity in Africa. While the role of obesity in the pathogenesis through insulin resistance is not disputed, the relative contributions of insulin resistance and β -cell dysfunction to the pathophysiology of type 2 diabetes in African populations are not completely clear. Data on the changes in the β -cell function and insulin resistance in the early stages of the disease process in African populations are scarce.³⁰ Although there is accumulating evidence to indicate that the pathogenic processes in type 2 diabetes are triggered by environmental factors, namely, physical inactivity and dietary changes, which mainly exacerbate insulin resistance, the genetic basis for

Table 2 Population-based prevalence studies on overweight and obesity in Africa

Study	Country	Sample	Sampling	Setting	Age (years)	Obesity (%)			Overweight (%)			Abdominal obesity (%)		
						Overall	Men	Women	Overall	Men	Women	Overall	Men	Women
Fouda ⁵⁵	Cameroon	552	Random	Urban	34.5	23.4	17.8	36.1	49.1	51.7	43.2	35.2	19.8	69.8
Sani ⁵⁶	Nigeria	300	Convenient	Urban	37.6	21.3	10.9	29.2	32	–	–	43.7	12.4	67.3
Ejike ²⁸	Nigeria	1584	Random	Urban	21.8	1.3	0.4	2.5	19.4	17.1	22.3	7.5	1.3	16.1
Ejim ¹³	Nigeria	858	Random	Rural	59.8	30	21.1	33.6	–	–	–	31	2.4	42.6
Wahab ⁵⁷	Nigeria	300	Convenient	Urban	–	21	9.3	29.8	53.3	41.9	62	–	–	–
Okafor ²⁹	Nigeria	898	Random	Urban	48.7	21.2	–	–	40.4	–	–	66.5	–	–
Mkhonto ⁵⁸	South Africa	532	Convenient	Rural	45.9	24.4	9.6	29.6	26.1	17.7	29.0	–	–	–
Malaza ⁴¹	South Africa	14 198	Convenient	Rural	≥15	45.7	15.9	45.7	23.3	31.3	4.9	–	–	–
Baragou ¹⁶	Togo	2000	Random	Urban	39	25.2	16.7	32.2	–	–	–	–	–	–
Mayega ²⁷	Uganda	1656	Random	Rural	44	5.3	2.2	8.2	12.3	7.5	16.9	13.8	1.4	24.5
Rudatsikira ⁵⁹	Zambia	1928	Random	Urban	–	14.3	5.1	18.6	–	–	–	–	–	–

insulin resistance and β -cell dysfunction has not been widely investigated in African populations. While environmental factors are very important, genetic factors probably have a role. Some of the diabetes genes discovered through genome-wide association studies elsewhere have recently been confirmed in African populations. However, this has been restricted to a smaller number of genes loci studied among West African populations from Nigeria and Ghana,³¹ and Northern African populations from Tunisia and Morocco.^{32–35} Although some of the studies on diabetes have also investigated obesity, overall, studies of the genetics of obesity in African populations remain rare.^{36–38} Given the significant interethnic differences in risk allele frequency and location in studies conducted out of the continent, it is logical to speculate that there may be some specificities of African populations, especially given the potential variability that may exist from the North to South of the continent.

CONCURRENT AND COMORBIDITIES OF DIABETES IN AFRICA

Diabetes, obesity and HIV infection

There is increasing evidence linking HIV infection with diabetes mellitus and impaired glucose tolerance, resulting from the natural history of HIV infection, comorbidities and treatments.^{39–40} The improved survival among people with HIV in Africa, following the improved access to highly active antiretroviral therapy (HAART) and treatment of comorbidities in many countries in the region in the recent years, will likely affect the prevalence of diabetes and dysglycaemia in Africa. However, the true magnitude of these effects is yet to be determined, and the mechanisms involved still to be investigated.³⁹

There have been suggestions that the burden of HIV infection could mask the emerging epidemic of obesity in Africa,³⁹ due to HIV related weight loss and wasting. Although access to HAART has significantly increased in SSA in recent years, population-based measures of effects, if any, on the occurrence of obesity have been lacking. In a cross-sectional study conducted in 2010 on a large sample from rural South Africa, with a high prevalence of HIV infection (24.1%), HIV infection was still associated with low BMI, even in the context of high access to HAART.⁴¹ In this study, while the prevalence of overweight was not different by HIV status, obesity was significantly more prevalent among HIV-negative individuals than HIV-positive participants (24.5% vs 20.0%). Being on HAART was associated with low BMI, a finding difficult to interpret in the absence of adjustment for the severity of HIV infection and the duration on HAART.⁴¹ Furthermore, whether some of the observed

effects could be ascribed to the fact that HIV patients on HAART are more likely to be health conscious than untreated patients or HIV negative remains unknown.

Diabetes and SCD

A recent study from Cameroon has provided findings suggesting similar frequencies of sickle cell trait (SCT) in people with diabetes and in the general population,⁴² and that the presence of SCT did not affect the metabolic control of diabetes. A striking fact in this study, however, was the older age of patients with SCT, for a comparable duration of diagnosed diabetes with the non-trait patients, indicating a possible late onset of type 2 diabetes in SCT carriers. Clarifying a possible link between SCT and diabetes onset or early attrition in diabetes is important in Africa, where the largest number of people with SCT and sickle cell disease (SCD) are found. The growing population with diabetes in Africa will likely include more people with both diabetes and SCT/SCD, and may result in differential features of diabetes across the continent, since the majority of those with SCT/SCD are typically found in Equatorial Africa, the epicentre of the disease.⁴² Resolving this uncertainty may invite efforts beyond what a single country can achieve. The health consequences of the co-occurrence of diabetes and SCT on the outcomes of both conditions are yet to be clarified.

Diabetes and MetS

Early studies of cardiovascular risk in people with diabetes were based on single risk factor approach.^{43–44} In recent studies, investigators have increasingly embraced the multiple risk factors approach to cardiovascular risk evaluation in diabetes. About 13 such studies published over the last 10 years from six SSA and one Northern Africa countries have characterised the cardiovascular disease risk in diabetes in clinical setting through screening for metabolic syndrome (MetS, table 3). These studies have used various criteria to define MetS, which makes comparison across studies difficult, but studies have been consistent in showing higher prevalence of MetS in women with diabetes than in men. Among the studies that used the IDF criteria to diagnose MetS, the prevalence ranged from 27.1% in urban Congo (Democratic Republic)⁴⁵ to >80% in a nationwide sample in Seychelles.⁴⁶ The variable prevalence across studies is mostly due to the use of variable thresholds, not supported by local evidence, to screen for different components of MetS. This is reflected by the non-perfect agreement between diagnostic criteria in those studies that have compared the yield of different criteria in the same population in Africa.^{46–48}

Table 3 Studies of metabolic syndrome in diabetes in hospital settings across Africa

Study	Country	Year published	Sample	Setting	Age	Duration	Criteria	Metabolic syndrome (%)		
								Overall	Men	Women
Longo-Mbenza ⁴⁵	DRC	2010	1266	Urban	50	–	IDF	27.1	–	–
Kengne ⁴⁷	Cameroon	2012	308	Urban	55.8	3.0	IDF	71.7	58	86.1
	Cameroon	2012	308	Urban	55.8	3.0	NCEP	60.4	42.7	78.8
Alshkri ⁶⁰	Libya	2008	99	Urban	56	9.4	NCEP	92	82	97
	Libya	2008	99	Urban	56	9.4	IDF	81	68	88
	Libya	2008	99	Urban	56	9.4	NCEP	60.4	42.7	78.8
Ogbera ⁵⁰	Nigeria	2010	973	Urban	58.6	6.9	JIS	86	83	87
Unadike ⁶¹	Nigeria	2009	240	Urban	50.8	–	NCEP	62.5	41.5	79.1
Adediran ⁶²	Nigeria	2007	408	Urban	56	–	WHO	51	44	56
Isezuo ⁶³	Nigeria	2005	254	Urban	52	6.1	WHO	59	–	–
Alebiosu ⁶⁴	Nigeria	2004	218	Urban	53.4	8.5	WHO	25.2	25	25.6
Puepet ⁶⁵	Nigeria	2009	634	Urban	54.2	–	IDF	63.6	74.5	54.9
Kelliny ⁴⁶	Seychelles	2008	130	Nationwide	25–64	–	NCEP	–	66.8	87.1
	Seychelles	2008	130	Nationwide	25–64	–	WHO	–	85.5	79.7
	Seychelles	2008	130	Nationwide	25–64	–	IDF	–	74	93
Kalk ⁶⁶	South Africa (black)	2008	500	Urban	48.3	2.0	IDF	50.2	46.5	74.1
	South Africa (white)	2008	254	Urban	54.1	6.0	IDF	73.6	52.9	73.1
Makuyana ⁶⁷	Zimbabwe	2004	109	Urban	–	–	WHO	43	–	–

IDF, International Diabetes Federation; JIS, Joint Interim Statement; NCEP, National Cholesterol Education Program, Adult Treatment Panel III.

The Joint Interim Statement (JIS) criteria recently published as part of the efforts to harmonise the definition criteria for MetS has not been recommended for use in people with diabetes as virtually all those with diabetes would qualify for MetS based on the JIS criteria.⁴⁹ Studies of MetS in people with diabetes in Africa using the JIS criteria have rather found variable prevalence rates not reaching 100% in all circumstances.^{47 50}

HEALTH CONSEQUENCES OF DIABETES

The burden and likely determinants of the well characterised microvascular and macrovascular complications of diabetes, and related morbidity and mortality in Africa have been recently described.^{6 23 39 43 51 52} Recent global efforts have focused on establishing a connection among diabetes mellitus, treatment and cancer risk. This issue was recently examined in a cohort of 9559 participants followed for a median of 15.1 years in Mauritius.⁵³ The study revealed a high risk of all-cause cancer mortality with increasing 2-h baseline glucose in men, but not in women. In an analysis stratified by gender and ethnicity, a significant association between baseline diabetes and all-cause cancer mortality was found only among South Asian men.⁵³ Although this study suggested that the increasingly reported association between diabetes and cancer risk may also apply to populations in Africa, the study was statistically underpowered to provide a definitive answer to the question in this setting.

CONCLUSIONS

Diabetes mellitus and obesity are important health challenges faced by countries in Africa, which are currently experiencing rapid transitions. Despite the uncertainties about the true magnitude of the conditions, diabetes and obesity figures will increase in Africa over the next few decades at a faster pace than elsewhere in the world. There is a hope that our knowledge base on the burden of diabetes and obesity in Africa can be substantially improved at little additional cost through careful exploitation of data already collected in many countries in recent years. However, new research is ultimately needed to

better characterise the interaction of genetic and environmental factors on the occurrence of diabetes and obesity in Africa.

Contributors Study conception: APK, JCM. Literature search and review: APK, JBE. Drafting the manuscript: APK, JBE. Revision of the draft: APK, JBE, ES, JCM.

Competing interests None.

Provenance and peer review Commissioned; externally peer reviewed.

Funding None

Data sharing No data to share

Ethics approval and Patients consent Not applicable

REFERENCES

- 1 Echouffo-Tcheugui JB, Kengne AP. A United Nation high level meeting on chronic non-communicable diseases: utility for Africa? *Pan Afr Med J* 2012;11:71.
- 2 International Diabetes Federation. In: Unwin N, Whiting D, Guariguata L, Ghyoot G, Gan D, Eds. *Updated diabetes atlas 2011*. 5th ed. Brussels, 2011.
- 3 Finucane MM, Stevens GA, Cowan MJ, et al. National, regional, and global trends in body-mass index since 1980: systematic analysis of health examination surveys and epidemiological studies with 960 country-years and 9.1 million participants. *Lancet* 2011;377:557–67.
- 4 Danaei G, Finucane MM, Lu Y, et al. National, regional, and global trends in fasting plasma glucose and diabetes prevalence since 1980: systematic analysis of health examination surveys and epidemiological studies with 370 country-years and 2.7 million participants. *Lancet* 2011;378:31–40.
- 5 Mbanya JC, Kengne AP, Assah F. Diabetes care in Africa. *Lancet* 2006;368:1628–9.
- 6 Mbanya JC, Motala AA, Sobngwi E, et al. Diabetes in sub-Saharan Africa. *Lancet* 2010;375:2254–66.
- 7 Badran M, Laher I. Type II Diabetes Mellitus in Arabic-Speaking Countries. *Int J Endocrinol* 2012;2012:902873.
- 8 Kengne AP, Echouffo-Tcheugui J, Sobngwi E, et al. New Insights on Diabetes Mellitus and Obesity in Africa—Part 2: Prevention, Screening and Economic burden. *Heart* 2013. doi: 10.1136/heartjnl-2013-303773
- 9 Erasmus RT, Soita DJ, Hassan MS, et al. High prevalence of diabetes mellitus and metabolic syndrome in a South African mixed ancestry population: the Bellville-South Africa study—Baseline data. *S Afr Med J* 2012;102:841–4.
- 10 Peer N, Steyn K, Lombard C, et al. Rising diabetes prevalence among urban-dwelling black South Africans. *PLoS One* 2012;7:e33336.
- 11 Evaristo-Neto AD, Foss-Freitas MC, Foss MC. Prevalence of diabetes mellitus and impaired glucose tolerance in a rural community of Angola. *Diabetol Metab Syndr* 2010;2:63.

- 12 Duboz P, Chapuis-Lucciani N, Boetsch G, *et al.* Prevalence of diabetes and associated risk factors in a Senegalese urban (Dakar) population. *Diabetes Metab* 2012;38:332–6.
- 13 Ejim EC, Okafor CI, Emehel A, *et al.* Prevalence of cardiovascular risk factors in the middle-aged and elderly population of a Nigerian rural community. *J Trop Med* 2011;21:308687.
- 14 Hammami S, Mehri S, Hajem S, *et al.* Prevalence of diabetes mellitus among non institutionalized elderly in Monastir City. *BMC Endocr Disord* 2012;12:15.
- 15 Echouffo-Tcheugui JB, Dzudie A, Epacka ME, *et al.* Prevalence and determinants of undiagnosed diabetes in an urban sub-Saharan African population. *Prim Care Diabetes* 2012;6:229–34.
- 16 Baragou S, Djibril M, Atta B, *et al.* Prevalence of cardiovascular risk factors in an urban area of Togo: a WHO STEPS-wise approach in Lome, Togo. *Cardiovasc J Africa* 2012;23:309–12.
- 17 Katchunga P, Masumbuko B, Belma M, *et al.* Age and living in an urban environment are major determinants of diabetes among South Kivu Congolese adults. *Diabetes Metab* 2012;38:324–31.
- 18 Ploubidis GB, Mathenge W, De Stavola B, *et al.* Socioeconomic position and later life prevalence of hypertension, diabetes and visual impairment in Nakuru, Kenya. *Int J Public Health* 2013;58:133–41.
- 19 Gessler N, Labhard ND, Stolt P, *et al.* The lesson of Monsieur Nouma: effects of a culturally sensitive communication tool to improve health-seeking behavior in rural Cameroon. *Patient Educ Couns* 2012;87:343–50.
- 20 Chamie G, Kwarisiima D, Clark TD, *et al.* Leveraging rapid community-based HIV testing campaigns for non-communicable diseases in rural Uganda. *PLoS One* 2012;7:e43400.
- 21 WHO Regional Office for Africa. STEPS survey on Chronic Disease Risk Factors. <http://www.afro.who.int/en/clusters-a-programmes/hpr/health-risk-factors/diseases-surveillance/surveillance-country-profiles/step-survey-on-noncommunicable-disease-risk-factors.html>. (accessed 19 Sep 2012)
- 22 Guthold R, Louazani SA, Riley LM, *et al.* Physical activity in 22 African countries: results from the World Health Organization STEPwise approach to chronic disease risk factor surveillance. *Am J Prev Med* 2011;41:52–60.
- 23 Kengne AP, Ntyintyane LM, Mayosi BM. A systematic overview of prospective cohort studies of cardiovascular disease in sub-Saharan Africa. *Cardiovasc J Africa* 2012;23:103–12.
- 24 Balogun WO, Gureje O. Self-reported incident type 2 diabetes in the Ibadan study of ageing: relationship with urban residence and socioeconomic status. *Gerontology* 2013;59:3–7.
- 25 Teo K, Chow CK, Vaz M, *et al.* The Prospective Urban Rural Epidemiology (PURE) study: examining the impact of societal influences on chronic noncommunicable diseases in low-, middle-, and high-income countries. *Am Heart J* 2009;158:1–7 e1.
- 26 Matsha T, Soita DJ, Hassan MS, *et al.* Three-year's changes in glucose tolerance status in the Bellville South cohort: Rates and phenotypes associated with progression. *Diabetes Res Clin Pract* 2012. doi: 10.1016/j.diabres.2012.10.018.
- 27 Mayega RW, Makumbi F, Rutebemberwa E, *et al.* Modifiable socio-behavioural factors associated with overweight and hypertension among persons aged 35 to 60 years in Eastern Uganda. *PLoS One* 2012;7:e47632.
- 28 Eijke CE, Ijeh, II. Obesity in young-adult Nigerians: variations in prevalence determined by anthropometry and bioelectrical impedance analysis, and the development of % body fat prediction equations. *Int Arch Med* 2012;5:22.
- 29 Okafor CI, Fasanmade O, Ofogebu E, *et al.* Comparison of the performance of two measures of central adiposity among apparently healthy Nigerians using the receiver operating characteristic analysis. *Indian J Endocrinol Metab* 2011;15:320–6.
- 30 Amoah AG, Schuster DP, Gaillard T, *et al.* Insulin resistance, beta cell function and cardiovascular risk factors in Ghanaians with varying degrees of glucose tolerance. *Ethn Dis* 2002;12:S3–10–17.
- 31 Chen G, Adeyemo A, Zhou J, *et al.* Genome-wide search for susceptibility genes to type 2 diabetes in West Africans: potential role of C-peptide. *Diabetes Res Clin Pract* 2007;78:e1–6.
- 32 Bouhaha R, Baroudi T, Ennafaa H, *et al.* Study of TNFalpha -308G/A and IL6–174G/C polymorphisms in type 2 diabetes and obesity risk in the Tunisian population. *Clin Biochem* 2010;43:549–52.
- 33 Ezzidi I, Mtraoui N, Nemr R, *et al.* Variants within the calpain-10 gene and relationships with type 2 diabetes (T2DM) and T2DM-related traits among Tunisian Arabs. *Diabetes Metab* 2010;36:357–62.
- 34 Ezzidi I, Turki A, Messaoudi S, *et al.* Common polymorphisms of calpain-10 and the risk of type 2 diabetes in a Tunisian Arab population: a case-control study. *BMC Med Genet* 2010;11:75.
- 35 Bouhaha R, Choquet H, Meyre D, *et al.* TCF7L2 is associated with type 2 diabetes in nonobese individuals from Tunisia. *Pathol Biol (Paris)* 2010;58:426–9.
- 36 Adeyemo A, Chen G, Zhou J, *et al.* FTO genetic variation and association with obesity in West Africans and African Americans. *Diabetes* 2010;59:1549–54.
- 37 Chen G, Adeyemo AA, Johnson T, *et al.* A genome-wide scan for quantitative trait loci linked to obesity phenotypes among West Africans. *Int J Obes (Lond)* 2005;29:255–9.
- 38 Yako YY, Fanampe BL, Hassan MS, *et al.* Association of cocaine- and amphetamine-related transcript, leptin and leptin receptor gene polymorphisms with anthropometric obesity phenotype indicators in South African learners. *J Nutrigenet Nutrigenomics* 2011;4:210–21.
- 39 Levitt NS. Diabetes in Africa: epidemiology, management and healthcare challenges. *Heart* 2008;94:1376–82.
- 40 Paik IJ, Kotler DP. The prevalence and pathogenesis of diabetes mellitus in treated HIV-infection. *Best Pract Res Clin Endocrinol Metab* 2011;25:469–78.
- 41 Malaza A, Mossong J, Barnighausen T, *et al.* Hypertension and obesity in adults living in a high HIV prevalence rural area in South Africa. *PLoS One* 2012;7:e47761.
- 42 Ama V, Kengne AP, Nansseu NJ, *et al.* Would sickle cell trait influence the metabolic control in sub-Saharan individuals with type 2 diabetes? *Diabet Med* 2012;29:e334–e7.
- 43 Kengne AP, Amoah AG, Mbanya JC. Cardiovascular complications of diabetes mellitus in sub-Saharan Africa. *Circulation* 2005;112:3592–601.
- 44 Kengne AP, Njamnshi AK, Mbanya JC. Cardiovascular risk reduction in diabetes in Sub-Saharan-Africa: what should the priorities be in the absence of global risk evaluation tools? *Clin Med: Cardiol* 2008;2:25–31.
- 45 Longo-Mbenza B, Kasiam Lasi On'kin JB, Nge Okwe A, *et al.* The metabolic syndrome in a Congolese population and its implications for metabolic syndrome definitions. *Diabetes Metab Syndr* 2011;5:17–24.
- 46 Kelliny C, William J, Riesen W, *et al.* Metabolic syndrome according to different definitions in a rapidly developing country of the African region. *Cardiovasc Diabetol* 2008;7:27.
- 47 Kengne AP, Limen SN, Sobngwi E, *et al.* Metabolic syndrome in type 2 diabetes: comparative prevalence according to two sets of diagnostic criteria in sub-Saharan Africans. *Diabetol Metab Syndr* 2012;4:22.
- 48 Motala AA, Esterhuizen T, Pirie FJ, *et al.* The prevalence of metabolic syndrome and determination of the optimal waist circumference cutoff points in a rural South African community. *Diabetes Care* 2011;34:1032–7.
- 49 Alberti KG, Eckel RH, Grundy SM, *et al.* Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation* 2009;120:1640–5.
- 50 Ogbera AO. Prevalence and gender distribution of the metabolic syndrome. *Diabetol Metab Syndr* 2010;2:1.
- 51 Kengne AP, Dzudie A, Sobngwi E. Heart failure in sub-Saharan Africa: a literature review with emphasis on individuals with diabetes. *Vasc Health Risk Manag* 2008;4:123–30.
- 52 Burgess PL, McCormick IJ, Harding SP, *et al.* Epidemiology of diabetic retinopathy and maculopathy in Africa: a systematic review. *Diabet Med* 2012; doi: 10.1111/j.1464-5491.2012.03756.x.
- 53 Harding JL, Soderberg S, Shaw JE, *et al.* All-cause cancer mortality over 15 years in multi-ethnic Mauritius: The impact of diabetes and intermediate forms of glucose tolerance. *Int J Cancer* 2012;131:2385–93.
- 54 Nsakshalo-Senkwe M, Siziya S, Goma FM, *et al.* Combined prevalence of impaired glucose level or diabetes and its correlates in Lusaka urban district, Zambia: a population based survey. *Int Arch Med* 2011;4:2.
- 55 Fouda AA, Lemougoum D, Manga JO, *et al.* [Epidemiology of obesity in the work milieu, Douala, Cameroon]. *Rev Med Brux* 2012;33:131–7.
- 56 Sani MU, Wahab KW, Yusuf BO, *et al.* Modifiable cardiovascular risk factors among apparently healthy adult Nigerian population—a cross sectional study. *BMC Res Notes* 2010;3:11.
- 57 Wahab KW, Sani MU, Yusuf BO, *et al.* Prevalence and determinants of obesity—a cross-sectional study of an adult Northern Nigerian population. *Int Arch Med* 2011;4:10.
- 58 Mkhonto SS, Labadarios D, Mabaso ML. Association of body weight and physical activity with blood pressure in a rural population in the Dikgale village of Limpopo Province in South Africa. *BMC Res Notes* 2012;5:118.
- 59 Rudatsikira E, Muula AS, Mulenga D, *et al.* Prevalence and correlates of obesity among Lusaka residents, Zambia: a population-based survey. *Int Arch Med* 2012;5:14.
- 60 Alshkri M, Elmehdawi R. Metabolic syndrome among type-2 diabetic patients in Benghazi-Libya: a pilot study. *Libyan J Med* 2008;3:177–80.
- 61 Unadike BC, Akpan NA, Peters EJ, *et al.* Prevalence of the metabolic syndrome among patients with type 2 diabetes mellitus in Uyo, Nigeria. *Afr J Endocrinol Metab* 2009;8:7–9.
- 62 Adediran OS, Edo AE, Jimoh AK, *et al.* Prevalence of the metabolic syndrome among Nigerians with type 2 diabetes. *Diabetes Int* 2007;15:13–14.
- 63 Isezuo SA, Ezunu E. Demographic and clinical correlates of metabolic syndrome in Native African type-2 diabetic patients. *J Natl Med Assoc* 2005;97:557–63.
- 64 Alebiosu CO, Odusan BO. Metabolic syndrome in subjects with type-2 diabetes mellitus. *J Natl Med Assoc* 2004;96:817–21.
- 65 Puepet FH, Uloko A, Akogu IY, *et al.* Prevalence of the metabolic syndrome among patients with type 2 diabetes mellitus in urban North-central Nigeria. *Afr J Endocrinol Metab* 2009;8:10–12.
- 66 Kalk WJ, Joffe BI. The metabolic syndrome, insulin resistance, and its surrogates in African and white subjects with type 2 diabetes in South Africa. *Metab Syndr Relat Disord* 2008;6:247–55.
- 67 Makuyana D, Gomo Z, Munyombwe T, *et al.* Metabolic syndrome disorders in urban black Zimbabweans with type 2 Diabetes mellitus. *Cent Afr J Med* 2004;50:24–9.