

Supplemental file

eTable 1 – Search strategy for PubMed

<p>“atrial fibrillation”</p>	<p>Delivery of Health Care, Integrated [mh:noexp] OR “integrated health care” OR multidisciplinary OR “ambulatory care” OR “ambulatory monitoring” OR Outpatient* OR “Interdisciplinary Communication” OR “Interdisciplinary Communications”</p>	<p>Outcome* OR “treatment failure” OR “treatment failures” OR ((cardiac OR cardiovascular OR heart) AND (death* OR mortalit*)) OR Fatal* OR Hospitalization [mh] OR “hospital admission” OR “hospital admissions” OR “quality of life” OR HRQL OR “life quality” OR “quality adjusted life year” OR qaly OR “quality adjusted life years” OR “short form 12” OR SF-12 OR SF12 OR “short form 20” OR SF-20 OR SF20 OR “short form 36” OR SF-36 OR SF36 OR “short form 8” OR SF-8 OR SF8 OR “symptom burden” OR “burden of symptoms”</p>
------------------------------	---	--

eTable 2 – Assessment of risk of bias in included studies

	Selection bias – random sequence generation	Selection bias – allocation concealment	Performance bias - Blinding of participant and personnel	Detection bias – blinding of outcome assessment	Attrition bias – incomplete outcome data	Reporting bias – selective reporting
Hendriks et al	Low	N/A	N/A	Low	Low	Low
Stewart et al	Low	N/A	N/A	Low	Low	Low
Carter et al	High	N/A	N/A	Low	Low	Low