

was the evaluation of the short-term success and safety after PV stenting of severe stenosis.

**Methods and results** 5 patients (pts) with 11 PVS were prospectively evaluated. All 5 pts experienced onset of dyspnoea on exertion some weeks or several months before, 3 with dyspnoea at rest, 5 with cough, and 2 with chest pain. On multirow spiral CT examination for each case, the narrowest lumen of the affected PVs measured  $2\pm 1.1$  mm compared with  $10\pm 1.3$  mm at baseline ( $p\leq 0.001$ ). The length of narrow segment was  $10\pm 2.9$ mm. All of 11 veins were treated with stent implantation with Palmaz Genesis stents, which size were 7- or 8-mm-diameter and 12-, 15-, 18 -mm-length. At percutaneous intervention, these veins showed  $81\pm 13\%$  stenosis, with a mean gradient of  $20\pm 5$  mm Hg. This was significantly reduced to a residual stenosis of  $8\pm 2\%$  ( $p\leq 0.001$ ) and a residual gradient of  $3\pm 2$  mm Hg ( $p\leq 0.001$ ) immediately. The mean pulmonary pressure was  $46\pm 5$  mm Hg pre-intervention, which decreased to  $28\pm 4$  mm Hg post-intervention in our protocols. The procedure was safety and successful, no complications were occurred in our pts. The symptomatic response was improved immediately and significantly.

**Conclusions** Percutaneous intervention produces rapid and dramatic symptom relief in patients with highly symptomatic PV stenosis after radiofrequency ablation for AF. Nevertheless, mid-term and long-term follow-up study with flow determination in the affected vein is still need to assess.

#### e0490 SAFETY AND FEASIBILITY OF TRANSRADIAL CORONARY ANGIOGRAPHY AT THE OUTPATIENT CLINIC

doi:10.1136/hrt.2010.208967.490

Yang Shiwei, Zhou Yujie, Hu Dayi, Shi Dongmei, Yang Qing, Wang Jianlong, Cheng Wanjuan, Nie Bin, Wang Zhijian, Li Yunzhi, Liu Xiaoli. *Beijing Anzhen Hospital Affiliated to Capital Medical University*

**Objective** To evaluate the safety and feasibility of transradial coronary angiography at the outpatient clinic.

**Methods** From February, 2008 to June, 2008, 100 outpatients who received transradial coronary angiography in Anzhen Hospital were included in this analysis, and 100 inpatients who underwent coronary angiography were selected as control group. Primary endpoints included success rate, percentage of angiographic catheter use with different diameters, adverse events during the procedure (such as death, malignant arrhythmia, acute myocardial infarction, coronary artery spasm, coronary artery dissection, perforation or occlusion, etc.) and after the procedure (such as death, acute myocardial infarction, upper limb haematoma, osteofascial compartment syndrome, radial artery pseudoaneurysm or occlusion, etc.).

**Results** The success rate (100% vs 100%), procedure duration time ( $12.5\pm 3.4$ ) min vs ( $10.8\pm 3.6$ ) min,  $p=0.517$ ) and exposition time ( $4.3\pm 1.0$ ) min vs ( $4.1\pm 1.0$ ) min,  $p=0.629$ ) were similar between the outpatient and inpatient groups. Radial and coronary artery spasm were the main adverse events during the angiography, and haematoma was the main adverse event after the angiography. There were no significant differences of adverse events between the 2 groups. the total cost of the outpatient group was significantly lower than the inpatient control group ( $4012\pm 238$ ) yuan vs ( $5329\pm 371$ ) yuan,  $p<0.01$ ). Expenditure including chemical tests, medicine, nursing care, Room & Board all decreased significantly.

**Conclusion** Transradial coronary angiography application at the outpatient clinic was safe and feasible for stable patients, and this procedure could decrease the medical expenditure and shorten the admission time.

#### e0491 ASSOCIATION OF CORONARY HEART DISEASE WITH CAROTID ARTERY INTIMA-MEDIA THICKNESS AND BAPWV

doi:10.1136/hrt.2010.208967.491

Liu Junming, Huang Menjun, Wang Mingjian, Xie Wei, Dou Qinghui, Zou Qi. *Department of Cardiovascular Internal Medicine, Bingtuan Hospital of Production and Construction, China*

**Objective** To investigate the association of coronary heart disease with carotid artery intima-media thickness and baPWV, by measuring carotid intima - media thickness and baPWV.

**Methods** 160 patients who had been examined by coronary angiography were divided into normal group and coronary artery disease group (this group was subdivided into 1 vessel group and multi-vessel group). Carotid ultrasonography and baPWV were performed on them.

**Results** Compared to the patients in normal group, The carotid artery IMT of the coronary artery disease group increased significantly with the aggravation of coronary artery stenosis (the normal group:  $0.83\pm 0.08$  mm, the 1 vessel group:  $0.91\pm 0.11$  mm, the multi-vessel group:  $1.08\pm 0.15$  mm,  $p<0.01$ ). The baPWV increased too (the normal group  $1411.20\pm 197.71$ , the 1 vessel group  $1742\pm 200.89$ , the multi-vessel group  $2839.40\pm 319.03$ ,  $p<0.01$ ). The carotid artery IMT and baPWV were significantly higher in multi-vessel group than those in 1 vessel group and normal group ( $p<0.01$ ).

**Conclusion** Carotid atherosclerosis and baPWV have great value in predicting coronary artery disease. For the coronary heart disease high risk group, routine carotid ultrasound examine and the measure of baPWV can be used to estimate the state of coronary artery atherosclerosis.

#### e0492 CAROTID INTIMIA-MEDIA THICKNESS AND CORONARY ARTERY DISEASE

doi:10.1136/hrt.2010.208967.492

Liu Junming, Huang Wenjun, Wang Mingjian, Xie Wei, Dou Qinghui, Zou Qi. *Department of Cardiovascular Internal Medicine, Bingtuan Hospital of Production and Construction, Xinjiang, China*

**Objective** To analyse the relationship of carotid atherosclerotic and coronary artery disease (CAD), by measuring Carotid intima - media thickness.

**Methods** The plaque index and intima - media thickness (IMT) were detected by high-resolution ultrasound in patients with coronary artery angiography. The extent of difference of IMT were compared between patients with and without coronary artery disease.

**Results** Compared to the patients with no CAD, the IMT of patients with CAD significantly increased ( $0.83\pm 0.08$  mm vs  $0.91\pm 0.11$  mm;  $0.83\pm 0.08$  mm MS  $1.08\pm 0.15$  mm;  $0.91\pm 0.11$  mm MS  $1.08\pm 0.15$  mm;  $p<0.05$ ). The incidence of plaque in CAD patients was higher than that of patients with no CAD. The IMT was significantly higher in multi-vessel group than those in uni-vessel group  $p<0.05$ ).

**Conclusion** In CAD patients, the widened inner diameter and lower blood flow rate of carotid artery result in a maximal shearing force significantly lower than that of the control group. The increasing intima-media thickness is associated greatly with carotid atherosclerosis. The haemodynamics change plays an important role in the development and progression of atherosclerosis in CAD patients. Carotid atherosclerosis has great value in predicting coronary artery disease. The IMT can be the surrogate markers of the extensive of coronary artery lesion.

**Heart**

## e0492 Carotid intima-media thickness and coronary artery disease

Liu Junming, Huang Wenjun, Wang Mingjian, Xie Wei, Dou Qinghui and Zou Qi

*Heart* 2010 96: A153

doi: 10.1136/hrt.2010.208967.492

---

Updated information and services can be found at:

[http://heart.bmj.com/content/96/Suppl\\_3/A153.3](http://heart.bmj.com/content/96/Suppl_3/A153.3)

---

*These include:*

### **Email alerting service**

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

---

### **Topic Collections**

Articles on similar topics can be found in the following collections

[Drugs: cardiovascular system](#) (8842)

[Clinical diagnostic tests](#) (4779)

---

### **Notes**

---

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>