

[gw22-e0792]

THE STUDY ON HUMAN UMBILICAL VEIN ENDOTHELIAL CELLS (HUVECS) PROLIFERATION AND THE BEST ENERGY IN SHOCK WAVES TREATMENT

Hongyan Cai¹, Tao Guo², Yu Wang³, Lin Li², Lin Zhao³, Xiyun Yang² ¹Department of Cardiology, The First Affiliated Hospital Kunming Medical College; ²The First Affiliated Hospital; ³Kunming Medical College

10.1136/heartjnl-2011-300867.106

Purpose To explore how different intensity of ultrasonic shock energy influences the human umbilical vein endothelial cells (HUVECs) proliferation, differentiation and the related cytokines, and to ascertain the best energy of CSWT.

Methods The HUVECs cell lines were performed using different levels of energy (0, 0.03, 0.09, 0.18, 0.24 mJ/mm²) shock wave treatment in vitro, Utilising CCK8 colorimetric method to detect HUVECs proliferation, real time PCR, Western blotting and Flow Cytometry were utilised to detect the changes in mRNA and protein of VEGF, IL-8, ICAM-1 before and after CSWT treatment.

Results The 0.09 mJ/mm² shock energy significantly promoted the HUVECs proliferation ($p < 0.05$). The results from real time PCR revealed that 0.09 mJ/mm² treatment markedly increased the expression of VEGF, IL-8, ICAM-1 ($p < 0.001$, respectively) compared with the non-treated control, the expression of ICAM-1 0.03 mJ/mm² group were increased compared with the control group ($p < 0.01$) While the expression of VEGF, IL-8 showed no significant changes ($p > 0.05$, respectively), 0.18 mJ/mm² treatment markedly increased the expression of VEGF ($p < 0.001$) compared with the non-treated control While the expression of ICAM-1, IL-8 showed no significant changes ($p > 0.05$, respectively), 0.24 mJ/mm² treatment showed no significant increase in the expression of VEGF, IL-8, ICAM-1 ($p > 0.05$, respectively) compared with the non-treated control. Western blot analysis and Flow Cytometry showed that the expression of VEGF, IL-8, ICAM-1 in 0.09 mJ/mm² group were

significantly higher than the control group ($p < 0.05$, respectively); the expression of VEGF in 0.03 mJ/mm² group was significantly higher than the control group ($p < 0.05$) While the expression of ICAM-1, IL-8 was not significantly higher than the control group ($p > 0.05$, respectively); the expression of VEGF in 0.18, 0.24 mJ/mm² group was significantly higher than the control group ($p < 0.05$) While the expression of ICAM-1, IL-8 was not significantly higher than the control group ($p > 0.05$, respectively).

Conclusions (1) The different extracorporeal shock wave energy promoted the HUVECs proliferations to different extent, the effect of 0.09 mJ/mm² energy is the most evident. (2) The 0.09 mJ/mm² energy increase the expression of IL-8, ICAM-1 mRNA and protein most significantly. (3) The 0.09 mJ/mm² energy increased the expression of VEGF mRNA and protein most remarkably. (4) The 0.03~0.24 mJ/mm² energies also have some effects in facilitating the secretion of VEGF, IL-8 and ICAM-1.