Editor's choice
Scan to access more
free content

ORIGINAL ARTICLE

Seasonality of cardiovascular risk factors: an analysis including over 230 000 participants in 15 countries

Helena Marti-Soler,¹ Cédric Gubelmann,¹ Stefanie Aeschbacher,² Luis Alves,^{3,4} Martin Bobak,⁵ Vanina Bongard,⁶ Els Clays,⁷ Giovanni de Gaetano,⁸ Augusto Di Castelnuovo,⁸ Roberto Elosua,⁹ Jean Ferrieres,⁶ Idris Guessous,^{1,10} Jannicke Igland,¹¹ Torben Jørgensen,^{12,13} Yuri Nikitin,¹⁴ Mark G O'Doherty,¹⁵ Luigi Palmieri,¹⁶ Rafael Ramos,^{17,18} Judith Simons,¹⁹ Gerhard Sulo,¹¹ Diego Vanuzzo,²⁰ Joan Vila,⁹ Henrique Barros,^{3,4} Anders Borglykke,¹² David Conen,² Dirk De Bacquer,⁷ Chiara Donfrancesco,¹⁶ Jean-Michel Gaspoz,¹⁰ Simona Giampaoli,¹⁶ Graham G Giles,²¹ Licia Iacoviello,⁸ Frank Kee,¹⁵ Ruzena Kubinova,²² Sofia Malyutina,^{14,23} Jaume Marrugat,⁹ Eva Prescott,^{24,25} Jean Bernard Ruidavets,⁶ Robert Scragg,²⁶ Leon A Simons,¹⁸ Abdonas Tamosiunas,²⁷ Grethe S Tell,¹¹ Peter Vollenweider,²⁸ Pedro Marques-Vidal¹

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/heartjnl-2014-305623>).

For numbered affiliations see end of article.

Correspondence to

Dr Pedro Marques-Vidal, Institut Universitaire de Médecine Sociale et Préventive, Bâtiment Biopôle 2, Route de la Corniche 10, 1010 Lausanne, Switzerland; Pedro-Manuel.Marques-Vidal@chuv.ch

The complete list of authors' emails is provided in online supplementary appendix 1.

Received 5 February 2014

Revised 16 April 2014

Accepted 1 May 2014

Published Online First

30 May 2014

► <http://dx.doi.org/10.1136/heartjnl-2014-306160>

To cite: Marti-Soler H, Gubelmann C, Aeschbacher S, et al. *Heart* 2014;**100**:1517–1523.

ABSTRACT

Objective To assess the seasonality of cardiovascular risk factors (CVRF) in a large set of population-based studies.

Methods Cross-sectional data from 24 population-based studies from 15 countries, with a total sample size of 237 979 subjects. CVRFs included Body Mass Index (BMI) and waist circumference; systolic (SBP) and diastolic (DBP) blood pressure; total, high (HDL) and low (LDL) density lipoprotein cholesterol; triglycerides and glucose levels. Within each study, all data were adjusted for age, gender and current smoking. For blood pressure, lipids and glucose levels, further adjustments on BMI and drug treatment were performed.

Results In the Northern and Southern Hemispheres, CVRFs levels tended to be higher in winter and lower in summer months. These patterns were observed for most studies. In the Northern Hemisphere, the estimated seasonal variations were 0.26 kg/m² for BMI, 0.6 cm for waist circumference, 2.9 mm Hg for SBP, 1.4 mm Hg for DBP, 0.02 mmol/L for triglycerides, 0.10 mmol/L for total cholesterol, 0.01 mmol/L for HDL cholesterol, 0.11 mmol/L for LDL cholesterol, and 0.07 mmol/L for glycaemia. Similar results were obtained when the analysis was restricted to studies collecting fasting blood samples. Similar seasonal variations were found for most CVRFs in the Southern Hemisphere, with the exception of waist circumference, HDL, and LDL cholesterol.

Conclusions CVRFs show a seasonal pattern characterised by higher levels in winter, and lower levels in summer. This pattern could contribute to the seasonality of CV mortality.

INTRODUCTION

Cardiovascular diseases (CVD) are the main cause of death worldwide, accounting for almost 32% of all deaths in women and 27% in men.¹ Cardiovascular events have been shown to present a seasonal pattern, their rates being higher in winter than in other seasons.² Several explanations

have been put forward, from misclassification of CVD deaths³ to changes in air temperature.⁴ A study conducted in 15 European cities showed that a decrease in air temperature of 1°C could lead to a 1.72% increase in the number of CVD deaths.⁵ Seasonal patterns for cardiovascular risk factors (CVRF) have also been suggested for Body Mass Index (BMI),⁶ waist circumference,⁶ blood pressure,⁷ lipids,⁸ and glycaemia.⁹ However, most studies assessing the seasonality of CVRFs have been conducted in relatively small studies or focused on a single CVRF.

Thus, the aim of this study was to assess whether CVRFs present a seasonal pattern, and if so, if this pattern is reproducible between countries. We used data from over 230 000 subjects from 24 studies (Northern Hemisphere: 192 129, Southern Hemisphere: 45 850) to achieve this aim.

METHODS

Studies included and data collection

Data from 24 population-based studies from 15 countries were included (Australia, Belgium, Czech Republic, Denmark, France, Italy, Lithuania, New Zealand, Northern Ireland, Norway, Portugal, Principality of Liechtenstein, Russia, Spain and Switzerland). The studies were initially gathered by the Prevention, Epidemiology and Population Science section of the European Association for Cardiovascular Prevention and Rehabilitation (EACPR), and other studies considered of interest were then invited. Twenty-one studies were from the Northern Hemisphere and three from the Southern Hemisphere. Information for the following CVRFs was collected: BMI; waist circumference; systolic (SBP) and diastolic blood pressure (DBP); total, high (HDL) and low (LDL) density lipoprotein cholesterol; triglycerides (log-transformed values) and blood glucose. Only data with the following criteria were considered: adult (≥18 years) participants,

objectively measured (non-self-reported) data, and data collection at least during an entire year. Missing data for a single month was not considered as an exclusion criterion for a study.

Each study provided monthly adjusted means and their associated SEs for the CVRFs available. Within each study, BMI and waist were adjusted for age (continuous), gender and current smoking (yes/no). For blood pressure, lipids and glucose levels, further adjustments on BMI and drug treatment (yes/no) were performed. A description of the studies is provided in online supplementary table S1.

Statistical analyses

Statistical analyses were performed using R Development Core Team (2011).¹⁰ To assess seasonality of CVRFs, weighted fixed-effects models including cosinor were used for testing a sinusoidal pattern, that is, a function assuming a cyclical and symmetrical behaviour of CVRFs over time and over their estimated mean.¹¹ Due to the opposite seasons in both Hemispheres, specific models according to Northern or Southern Hemisphere were applied. For each hemisphere, a joint model including the study indicator variable besides sine and cosine functions was adjusted

$$\text{CVRF} = \beta_0 + \beta_1 \sin\left(\frac{2\pi t}{T}\right) + \beta_2 \cos\left(\frac{2\pi t}{T}\right) + \sum_{i=1}^{n-1} \lambda_i \text{study}$$

where CVRF is the vector of the reported means, β_0 is the estimated mean, β_1 and β_2 are, respectively, the associated coefficients of sine and cosine functions, $t = 1, \dots, T$ is the time period, corresponding to the monthly data, $T = 12$ is the number of time periods described by the sinusoidal curve to assess yearly seasonality with one minimum value (nadir) and one maximum value (peak) per year, and λ_i is the associated coefficient of the indicator variable for the study i , assuming that there are n studies. The amplitude, that is, the distance from the mean to the peak or nadir, was estimated as $\sqrt{\beta_1^2 + \beta_2^2}$, and therefore, the seasonal variation, that is, the maximal difference between peak and nadir, was $2 \times \text{amplitude}$. The month of the two extreme values, nadir and peak in $(0, T)$ were estimated by solving $t = \tan^{-1}((\beta_1/\beta_2)/(T/2\pi))$. If $\beta_1/\beta_2 > 0$: t indicates the month of the first extreme (the peak if $\beta_1 > 0$, the nadir otherwise) and $t+T/2$ the month of the second one (the nadir if $\beta_1 > 0$, the peak otherwise). If $\beta_1/\beta_2 \leq 0$, the month of the first extreme is found at $t+T/2$, and the month of the second one at $t+T$. The percentage of the between-month variance explained by the sinusoidal curve was estimated using the adjusted R^2 measure.

A simulated effect attributable to seasonal variations in CVRFs was modelled by calculating the 10-year risk of dying from CV for the age range 40 years to 65 years using Stata V.12.1 (Stata Corp, College Station, Texas, USA). Simulations were conducted using the SCORE model for low-risk countries¹² for countries in the Northern Hemisphere, and using the recalibrated Australian SCORE equation¹³ for countries in the Southern Hemisphere (see online supplementary appendix 2). Briefly, the absolute difference on 10-year risk of dying from CV disease between the peak and the nadir months for participants aged between 40 years and 65 years was modelled by introducing in the SCORE equation the seasonal variation of each CVRF. Within each hemisphere, the simulations were conducted taking into account gender, smoking status and age group, and using seasonal variation of systolic blood pressure and total cholesterol.

RESULTS

Characteristics of participants

The characteristics of the participants in the different studies are summarised in online supplementary table S1. There were 192 129 participants in the Northern Hemisphere and 45 850 in the Southern Hemisphere. Age ranged from 18 years to 99 years, most studies presenting a mean age between 50 years and 60 years. Women represented half or more of the participants, with the exception of BELSTRESS and the Workforce Diabetes Survey where this percentage was considerably lower; the PRIME Belfast study included only men. Finally, the percentage of current smokers varied between 15% and 70% in men and between 8% and 58% in women.

Seasonality analyses

Seasonal analyses were carried out using data adjusted for age, gender and current smoking (yes/no). For blood pressure, lipids and glucose levels, further adjustment on BMI and drug treatment (yes/no) was performed. Table 1 summarises the estimated seasonal variation, nadir and peak when a seasonal pattern was observed for a given CVRF, and adjusted R^2 . The graphical representation of seasonal variations for anthropometric data, lipid parameters and glycaemia levels is shown in online supplementary figures S1 and S2.

For anthropometric variables, BMI displayed a seasonal pattern over time. The estimated seasonal variations were 0.26 kg/m^2 and 0.52 kg/m^2 for the Northern and the Southern Hemispheres, respectively, with corresponding nadirs occurring in July and December. For waist circumference, a seasonal variation was observed in the Northern Hemisphere only, with an estimated variation of 0.57 cm , and the nadir occurring in July. In the Northern Hemisphere, the sinusoidal curve explained over 95% of the total variance for BMI and waist, while in the Southern Hemisphere, this percentage was slightly lower, 78% for BMI and 82% for waist.

SBP as well as DBP showed a seasonal pattern (figure 1). For SBP, the estimated seasonal variations were approximately 2.9 mm Hg and 3.4 mm Hg in the Northern and the Southern Hemispheres, respectively, with nadirs occurring in June and January, respectively. For DBP, the estimated seasonal variations were 1.4 mm Hg and 0.9 mm Hg in the Northern and the Southern Hemispheres, respectively, the nadirs occurring in June and November, respectively. In the Northern Hemisphere, the sinusoidal curve explained 95% and 96% of the total variance for SBP and DBP, while in the Southern Hemisphere this percentage was 98% for SBP and 59% for DBP.

Most lipid parameters showed a seasonal pattern. For triglycerides, the estimated seasonal variations were 0.02 mmol/L and 0.07 mmol/L in the Northern and the Southern Hemispheres, respectively, the nadirs occurring in April and July, respectively. In the Northern Hemisphere, for total HDL and LDL cholesterol levels, the estimated seasonal variations were 0.10 mmol/L , 0.01 mmol/L and 0.11 mmol/L , respectively, with nadirs occurring in summer months. In the Southern Hemisphere, only total cholesterol levels showed a seasonal pattern, with estimated seasonal variation of 0.21 mmol/L and the nadir occurring in October. In the Northern Hemisphere, the sinusoidal curve explained between 91% and 93% of the total variance of lipids, while in the Southern Hemisphere the percentages varied from 64% to 89%.

Glycaemia levels also showed a seasonal pattern. The estimated seasonal variations were 0.07 mmol/L and 0.17 mmol/L in the Northern and the Southern Hemispheres, respectively,

Table 1 Seasonal variation estimates of cardiovascular risk factors

	Northern Hemisphere				Southern Hemisphere			
	Seasonal variation	Nadir	Peak	Adjusted R ²	Seasonal variation	Nadir	Peak	Adjusted R ²
Body Mass Index (kg/m ²)*	0.26†	July	January	0.97	0.52†	December	June	0.78
Waist circumference (cm) *	0.57†	July	January	0.95	0.77			0.82‡
Systolic blood pressure (mm Hg)§	2.93†	June	December	0.95	3.44†	January	July	0.98
Diastolic blood pressure (mm Hg)§	1.42†	June	December	0.96	0.86†	November	May	0.59
Triglycerides (mmol/L)§	0.02†	April	October	0.94	0.07†	July	January	0.87
Total cholesterol (mmol/L)§	0.10†	June	December	0.91	0.21†	October	April	0.85
HDL cholesterol (mmol/L)§	0.01†	August	February	0.93	0.04			0.64
LDL cholesterol (mmol/L)§	0.11†	July	January	0.91	0.10			0.89‡
Glycaemia (mmol/L)§	0.07†	August	February	0.98	0.17†	October	April	0.67

Data from 21 studies from the Northern Hemisphere and 3 from the Southern Hemisphere. Results are presented as seasonal variation, nadir, peak and adjusted R² (percentage of variance explained).

*Adjusted for gender, age and smoking.

†Seasonal variation statistically significant at 5% level, that is, at least one of the associated p values of sine or cosine functions <0.05.

‡Data were available for two studies only.

§Adjusted for gender, age, smoking and Body Mass Index.

the nadirs occurring in August and October, respectively. In the Northern Hemisphere, the sinusoidal curve explained 98% of the total variance of glycaemia, while in the Southern Hemisphere this percentage was lower at 67%.

Sensitivity analyses

As lipids and plasma glucose vary according to fasting status, a sensitivity analysis was performed on the 16 studies (14 in the Northern Hemisphere and 2 in the Southern Hemisphere) that collected blood samples in the fasting state. The results are summarised in table 2. Lipid as well as glucose levels showed a seasonal pattern, with estimated seasonal variation of 0.03 mmol/L for triglycerides, 0.13 mmol/L for total cholesterol, 0.02 mmol/L for HDL cholesterol, 0.12 mmol/L for LDL cholesterol and 0.08 mmol/L for glucose. Most nadirs occurred between June and July, with the exception of triglycerides, where the nadir occurred in March. Finally, the proportion of the total variance explained by the sinusoidal curve was higher than 80% for all markers studied. Graphical representation of seasonal variations

for lipid parameters and glycaemia levels is shown in online supplementary figures S3 and S4.

Estimated impact on cardiovascular mortality

The results of the simulation of the absolute difference in 10-year risk of dying from CV disease in the peak relative to the nadir month are summarised in figure 2. Briefly, the excess risk increased with age, smoking status and male sex. For a 65-year-old man living in the Northern Hemisphere and who smoked, the excess 10-year risk of dying from CV disease in winter relative to summer was as high as 6.9%, while for a non-smoking woman aged 40 years, the excess risk was less than 0.1% (figure 2).

DISCUSSION

To our knowledge, this is the largest and most comprehensive study ever conducted to assess the seasonality of CVRFs. Our results strongly suggest that CVRFs present a seasonal pattern with lower levels in summer and higher levels in winter, suggesting that

Figure 1 Seasonal variation in blood pressure estimated using weighted fixed-effects models including cosinor function, with one peak and one nadir for a time period of 12 months. Each study is represented as a line: dot lines and diamonds for studies in the Northern Hemisphere, and solid lines and circles for studies in the Southern Hemisphere.

Table 2 Seasonal variation estimates of lipids and glycaemia, restricted to studies that collected fasting blood samples

	Northern Hemisphere				Southern Hemisphere			
	Seasonal variation	Nadir	Peak	Adjusted R ²	Seasonal variation	Nadir	Peak	Adjusted R ²
Triglycerides (mmol/L)	0.03*	April	October	0.92	0.09*	July	January	0.54
Total cholesterol (mmol/L)	0.13*	June	December	0.89	0.07			0.64
HDL cholesterol (mmol/L)	0.02*	June	December	0.85	0.08*	June	December	0.46
LDL cholesterol (mmol/L)	0.12*	July	January	0.92	0.22			0.08†
Glycaemia (mmol/L)	0.08*	June	December	0.97	0.02			0.50

Data from 14 studies from the Northern Hemisphere and 2 from the Southern Hemisphere. Data adjusted for gender, age, smoking, body mass index and hypolipidemic (for lipids) or antidiabetic (for glycaemia) drug treatment. Results are presented as seasonal variation, nadir, peak and adjusted R² (percentage of variance explained).

*Seasonal variation statistically significant at 5% level, that is, at least one of the associated p values of sine or cosine functions <0.05.

†Data were available for one study only.

at least part of the patterning might be due to changes in air/outdoor temperature. Another possible explanation would be differences in exposure to sunlight, leading to differences in vitamin D levels and, thus, in CVRFs¹⁴; still, the effect of sunlight exposure on CVRFs has been questioned.¹⁵ Other factors such as increased levels of air pollution can also play a role. Higher concentrations of some pollutants have been associated with colder periods, and the relationship between an increased level of air pollution and cardiovascular disease has previously been established.¹⁶ Changes associated with winter/cold seasons, such as social withdrawal, the consumption of a diet with higher fat content and decreased levels of physical activity¹⁷ can also partially contribute to the seasonal pattern. This patterning cannot be attributed to the seasonal variation of BMI, as all analyses were conducted using BMI-adjusted data. Indeed, CVRFs can be mediated by BMI, as evidenced by the complete remission or considerable improvement of high blood pressure, cholesterol or glucose levels following weight loss after caloric reduction or bariatric surgery.^{18, 19} Hence, adjustment on BMI was deemed necessary prior to analysing the seasonality of the other CVRFs.

This is also the first study which includes observations from both hemispheres. As winter in the Northern Hemisphere corresponds to summer in the Southern Hemisphere, this should lead to a mirror image of seasonal variations, which was actually observed in the present study.

Anthropometry

BMI and waist circumference showed a small seasonal variation, a finding already reported in the literature.⁶ The most likely explanation is changes in physical activity²⁰ or dietary intake²¹ during the warmer months. For instance, whereas total energy intake does not change between winter and summer, diet composition does, with a higher intake of fat-rich foods in winter.²²

Blood pressure

Blood pressure values were higher in winter than in summer; this finding has been repeatedly reported.^{23, 24} The seasonal variations for SBP were approximately 2.9 mm Hg and 3.4 mm Hg in the Northern and the Southern Hemispheres, respectively. The most likely explanation is changes in lifestyle or in temperature. Temperature shifts lead to arteriolar vasoconstriction, decreased sweating and increased salt load.^{24, 25} Seasonal variation in the concentration of air pollutants might also influence blood pressure levels through the modulation of the autonomic nervous system, the activation of the renin-angiotensin system and the increase in endothelin.²⁶ The higher consumption of vegetables and fruits in summer²¹ could also intervene via an increase in potassium intake.²⁷ Finally, the reduced synthesis of vitamin D in winter could increase blood pressure levels, as vitamin D has been shown to inhibit the renin-angiotensin system²⁵; however, this hypothesis has been challenged.¹⁵

Figure 2 Effect of seasonal variation in cardiovascular risk factors on 10-year risk of dying from cardiovascular disease. The chart shows the absolute risk difference on 10-year risk of dying from CV disease between the peak and the nadir months for each hemisphere and for different gender, age and smoking combinations. Simulations were performed using the SCORE equation for low-risk countries for countries in the Northern Hemisphere, and using the recalibrated Australian SCORE equation for countries in the Southern Hemisphere.

Interestingly, the seasonal variations observed in our study are comparable and even slightly higher than those reported for genetic scores for blood pressure,²⁸ suggesting that season might exert a similar or even greater effect on average blood pressure levels than currently known genetic markers. Our results also suggest that studies assessing small effects on blood pressure (such as genome-wide association studies) should take into account the effect of season on their analysis.

Lipids

Seasonality was observed for triglycerides, total HDL and LDL cholesterol levels in the Northern Hemisphere. These findings are in agreement with several studies^{8, 27} and partly with others²⁹ where seasonality was found in men only. Possible explanations include a higher intake of fat in winter²² and an increased consumption of vegetables and fruits in summer²⁷ although no significant changes in dietary intake have been reported by others.⁸ Seasonal variations in plasma volume (haemodilution in summer and haemoconcentration in winter) have also been suggested.⁸

Seasonal variation of lipid levels had already been acknowledged in the European guidelines for management of dyslipidaemias³⁰ and could be of clinical importance in clinical trials or epidemiological studies. For instance, in the PROVE-IT trial, achievement of the LDL-goal was significantly higher in summer than in winter.³¹ Similarly, epidemiological studies conducted during the cold months might report higher levels of CVRFs than in warm months. Thus, it would be of interest that epidemiological studies be conducted throughout a whole year or, alternatively, that the exact timing of the study be reported.

Glycaemia

Seasonality of glucose levels has been reported among diabetic patients, with a peak in winter and a nadir in summer.³² A similar pattern has been observed for glycosylated haemoglobin among non-diabetic subjects.³³ In this study, a slight seasonal variation in glycaemia was observed, and the estimated variation was not modified when the analysis was restricted to studies that collected blood samples in the fasting state (see online supplementary figure S2). Several explanations have been put forward, namely an increase in hyperglycaemic hormones,³³ partly compensated by a parallel increase in insulin levels.⁹ Other possible explanations include changes in insulin sensitivity due to decreased temperatures in winter or increased physical activity in summer.³³

Estimated impact on cardiovascular mortality

The seasonal variations in CVRFs led to significant differences in estimated 10-year risk of dying from CV disease according to season. In elderly subjects, this difference ranged between 1.8% and 9.9%. Although the increase in CVRFs in the colder months might not translate into a higher risk of CVD in the future at the individual level, it does have clinical and population consequences. For instance, at the clinical level, a subject screened in the colder months will have a higher calculated 10-year risk of CVD and, thus, a higher likelihood of receiving preventive advice or treatment than in the warmer months. At the population level, the higher risk of future CVD event in colder months could also translate into a higher immediate risk: higher BP levels in the colder months could promote the triggering of CVD, such as stroke or myocardial infarction. Although the calculations were performed mainly for illustrative purposes, they suggest that the seasonal variation of CVRFs could partly explain the seasonality of CVD deaths in most countries.^{34, 35} Nevertheless, further studies are needed to better assess this point.

Strengths and weaknesses

To our knowledge, this is the largest and most comprehensive study ever conducted to assess the seasonality of CVRFs. Our results replicate between studies, and also between the Northern and the Southern Hemispheres.

This study has also some weaknesses worth acknowledging. First, only data from two countries of the Southern Hemisphere (Australia and New Zealand) were available. This led to smaller sample sizes, which might have prevented the assessment of seasonal patterns for some CVRFs in the Southern Hemisphere. Indeed, the percentage of the between-month variability explained by the sinusoidal curve was generally lower in the Southern Hemisphere than in the Northern Hemisphere. Nevertheless, only for DBP, HDL cholesterol and glycaemia, the percentage of variance explained by cosinor curve was lower than 70%. Second, the use of the cosinor model might be questionable, and a mixed-effects model might have been appropriate. However, preliminary results showed a similar seasonal pattern in most within-study analyses. Thus, no clear heterogeneity was found between studies, that is, studies with high values of CVRFs did not present a stronger seasonal pattern than studies with lower values. Further, the cosinor model can be described as a linear regression model using sine and cosine functions,³⁶ which has several advantages: it allows us to correctly estimate seasonal variations, and it is more flexible than polynomial regression, such as quadratic or cubic polynomials. Still, in the context of linear regression, the cosinor model provides a straightforward estimation of the seasonal variation, nadir and peak, as well as a measure of the goodness-of-fit. Third, the SCORE equation for low-risk countries was used, and it is likely that using the high-risk equation would have led to bigger seasonal risk differences. Hence, our estimations can be considered as conservative. Finally, seasonality was assessed using one single measurement from each study participant, instead of multiple measurements performed over time in the

Key messages

What is known on this subject?

Cardiovascular events have been shown to present a seasonal pattern, their rates being higher in winter than in other seasons. Seasonal patterns for cardiovascular risk factors (CVRFs) have also been suggested for Body Mass Index (BMI), waist circumference, blood pressure, lipids and glycaemia. However, most studies assessing the seasonality of CVRFs have been conducted in relatively small studies or focused on a single CVRF.

What might this study add?

A seasonal pattern characterised by higher levels in winter and lower levels in summer of CVRF levels was demonstrated in 24 populations of both hemispheres using a common methodology. The magnitude of the seasonal effect is comparable to the pooled effect of genetic markers. The seasonal pattern of CVRFs could explain the seasonality of CV mortality.

How might this impact on clinical practice?

Interpretation of individual risk factors levels might be different depending on season: a low value in winter is definitely a low 'yearly' value, but a low value in summer does not mean a low value in winter.

same persons. However, the results were adjusted for the main confounders and it is unlikely that the same recruiting bias (ie, subjects with low levels of CVRFs being recruited only in summer) would appear in most studies.

CONCLUSION

Cardiovascular risk factors show a seasonal pattern characterised by higher levels in winter and lower levels in summer. This pattern could contribute to the seasonality of CV mortality.

Author affiliations

- ¹Institute of Social and Preventive Medicine, University of Lausanne, Lausanne, Switzerland
- ²Department of Medicine, University Hospital Basel, Basel, Switzerland
- ³Department of Clinical Epidemiology, Predictive Medicine and Public Health, University of Porto Medical School, Porto, Portugal
- ⁴Institute of Public Health of the University of Porto, Porto, Portugal
- ⁵Department of Epidemiology and Public Health, University College London, London, UK
- ⁶Department of Epidemiology, UMR 1027- INSERM, Toulouse University—CHU Toulouse, Toulouse, France
- ⁷Department of Public Health, Ghent University, Ghent, Belgium
- ⁸Department of Epidemiology and Prevention, IRCCS Istituto Neurologico Mediterraneo NEUROMED, Pozzilli (IS), Italy
- ⁹Cardiovascular and Genetic Epidemiology Research Group (ULEC-EGEC), Inflammatory and Cardiovascular Disease Programme (RICAD), IMIM, Barcelona, Spain
- ¹⁰Department of Community Medicine, Preventive care and Emergency Medicine, Geneva University Hospitals, Geneva, Switzerland
- ¹¹Department of Global Public Health and Primary Care, University of Bergen, Bergen, Norway
- ¹²Research Centre for Prevention and Health, Capital Region of Denmark, Glostrup, Denmark
- ¹³Faculty of Health Science, University of Copenhagen, Denmark
- ¹⁴Laboratory of Internal Medicine, Institute of Internal Medicine, Siberian Branch RAMS, Novosibirsk, Russia
- ¹⁵UKCRC Centre of Excellence for Public Health, Queens University of Belfast, Belfast, UK
- ¹⁶Unit of Epidemiology of Cerebro and Cardiovascular Diseases, National Centre of Epidemiology, Surveillance and Health Promotion, Istituto Superiore di Sanità, Rome, Italy
- ¹⁷Research Unit and Docent Unit of Family Medicine Girona, Primary Care Research Institute Jordi Gol, Girona, Spain
- ¹⁸Departament de Medicina, Universitat de Girona, Girona, Spain
- ¹⁹Lipid Research Department, University of New South Wales, St Vincent's Hospital, Sydney, Australia
- ²⁰Centro di Prevenzione Cardiovascolare, ASS 4 'Medio Friuli', Udine, Italy
- ²¹Cancer Epidemiology Centre, Cancer Council Victoria, Victoria, Australia
- ²²Centre for Health Monitoring, National Institute of Public Health, Prague, Czech Republic
- ²³Novosibirsk State Medical University, Novosibirsk, Russia
- ²⁴Department of Cardiology, Bispebjerg University Hospital, Copenhagen, Denmark
- ²⁵Copenhagen City Heart Study, Frederiksberg Hospital, Copenhagen, Denmark
- ²⁶School of Population Health, University of Auckland, Auckland, New Zealand
- ²⁷Institute of Cardiology, Lithuanian University of Health Sciences, Kaunas, Lithuania
- ²⁸Department of Internal Medicine, University Hospital and University of Lausanne, Lausanne, Switzerland

Contributors HM-S, CG and PM-V wrote the manuscript. HM-S was responsible for analysis and interpretation data. HM-S, SA, LA, MB, VB, EC, GdG, ADC, RE, JF, IG, JJ, TJ, YN, MGO, LP, RR, JS, GS, DV, JV, HB, AB, DC, DDB, CD, J-MG, SG, GGG, LI, FK, RK, SM, JM, EP, JBR, RS, LAS, AT, GST, PV and PM-V were involved in acquisition of data. All authors critically revised the manuscript for important intellectual content. PMV supervised the study and is the guarantor.

Funding BELSTRESS was supported by grants from the Federal Office for Scientific, Technical, and Cultural Affairs (ST/02/007), the FWO-Vlaanderen, and the National Fund for Scientific Research (FNRS). The Bus Santé study is funded by the Geneva University Hospitals through the General Directorate of Health (Canton of Geneva). Idris Guessous is supported by a grant from the Swiss National Science Foundation (grant #33CM30-124087). The CoLaus/PsyCoLaus study was and is supported by research grants from GlaxoSmithKline, the Faculty of Biology and Medicine of Lausanne, Switzerland and three grants of the Swiss National Science Foundation (grants #3200B0-105993, #3200B0-118308, #33CSO-122661 and FN 33CSO-139468). The Copenhagen City Heart Study was mainly supported by the Danish Heart Foundation. The Dubbo Study was supported by the National Health &

Medical Research Council of Australia. The Epiporto study was financed by the Fundação para a Ciência e a Tecnologia, Portugal [PTDC/SAU-ESA/108315/2008]. The GAPP study was supported by the Liechtenstein Government, the Swiss Heart Foundation, the Swiss Society of Hypertension, the University of Basel, the University Hospital Basel, the Hanel Foundation, Schiller AG and Novartis, Switzerland. David Conen was supported by a grant of the Swiss National Science Foundation (PP00P3_133681). The HAPIEE Project is supported by research grants from Wellcome Trust (grants #064947/Z/01/Z, # 081081/Z/06/Z), MacArthur Foundation 'Health and Social Upheaval (a research network)' and National Institute on Aging (grant #1R01 AG23522). The Health2006 study was financially supported by grants from the Velux Foundation; the Danish Medical Research Council, Danish Agency for Science, Technology and Innovation; the Aase and Ejner Danielsens Foundation; ALK-Abelló A/S (Hørsholm, Denmark), Timber Merchant Vilhelm Bangs Foundation, MEKOS Laboratories (Denmark) and Research Centre for Prevention and Health, the Capital Region of Denmark. The Melbourne Collaborative Cohort Study (MCCS) recruitment and ongoing maintenance is funded by Cancer Council Victoria. It is also supported by the Australian National Health and Medical Research Council (grants #209057, #251533 and #396414). The enrolment phase of the Moli-sani Project was supported by research grants from Pfizer Foundation (Rome, Italy) and the Italian Ministry of University and Research (MIUR, Rome, Italy)—Programma Triennale di Ricerca, Decreto no.1588. MONA LISA was funded by Institut de Veille Sanitaire (InVS), Institut National de la Santé et de la Recherche Médicale (INSERM), Agence Nationale de la Recherche (ANR) and Pfizer. The Osservatorio Epidemiologico Cardiovascolare (OEC) and the Osservatorio Epidemiologico Cardiovascolare/Health Examination Survey (OEC/HES) are part of the CUORE Project of the Istituto Superiore di Sanità, Rome; both surveys were supported partly from the Italian Ministry of Health, grants of the Centre of Disease Control, and partly from the Italian Association of Hospital Cardiologists (ANMCO). The PRIME study was supported by grants from the Institut National de la Santé et de la Recherche Médicale (INSERM) and the Merck, Sharpe and Dohme-Chibret Laboratory. The Belfast PRIME Study was funded by the 'HSC Research and Development Office for Northern Ireland'. REGICOR Funding: Spanish Health Research Fund [FIS 94/0539, FIS PI020471]; Agency for Management of University and Research Grants [2005SGR00577], and Spain's Ministry of Science and Innovation through the Carlos III Health Institute & European Regional Development Funds (ERDF) [HERACLES Program RD12/0042]. The Workforce Diabetes Survey was supported by research grants from the Health Research Council of New Zealand.

Competing interests None.

Patient consent Obtained.

Ethics approval All studies were approved by the local or national ethics committees.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement The dataset is available from the corresponding author at Pedro-Manuel.Marques-Vidal@chuv.ch.

REFERENCES

- 1 World Health Organization. *The global burden of disease. 2004 update*. Geneva, Switzerland: World Health Organization, 2008.
- 2 Abrignani MG, Corrao S, Biondo GB, *et al*. Influence of climatic variables on acute myocardial infarction hospital admissions. *Int J Cardiol* 2009;137:123–9.
- 3 Wexelman BA, Eden E, Rose KM. Survey of New York City resident physicians on cause-of-death reporting, 2010. *Prev Chronic Dis* 2013;10:E76.
- 4 Ballester F, Corella D, Perez-Hoyos S, *et al*. Mortality as a function of temperature. A study in Valencia, Spain, 1991–1993. *Int J Epidemiol* 1997;26:551–61.
- 5 Analitis A, Katsouyanni K, Biggeri A, *et al*. Effects of cold weather on mortality: results from 15 European cities within the PHEWE project. *Am J Epidemiol* 2008;168:1397–408.
- 6 Visscher TL, Seidell JC. Time trends (1993–1997) and seasonal variation in body mass index and waist circumference in the Netherlands. *Int J Obes Relat Metab Disord* 2004;28:1309–16.
- 7 Lewington S, Li L, Sherliker P, *et al*. Seasonal variation in blood pressure and its relationship with outdoor temperature in 10 diverse regions of China: the China Kadoorie Biobank. *J Hypertens* 2012;30:1383–91.
- 8 Maclaughlan J, Claudia W. *Death certification reform: a case study on the potential impact on mortality statistics, England and Wales*. London, UK: Office for National Statistics, 2012.
- 9 Behall KM, Scholfield DJ, Hallfrisch JG, *et al*. Seasonal variation in plasma glucose and hormone levels in adult men and women. *Am J Clin Nutr* 1984;40:1352–6.
- 10 R Foundation for Statistical Computing. R: A language and environment for statistical computing. 2008. <http://www.R-project.org>
- 11 Stolwijk AM, Straatman H, Zielhuis GA. Studying seasonality by using sine and cosine functions in regression analysis. *J Epidemiol Community Health* 1999;53:235–8.

- 12 Conroy RM, Pyorala K, Fitzgerald AP, *et al.* Estimation of ten-year risk of fatal cardiovascular disease in Europe: the SCORE project. *Eur Heart J* 2003;24:987–1003.
- 13 Chen L, Tonkin AM, Moon L, *et al.* Recalibration and validation of the SCORE risk chart in the Australian population: the AusSCORE chart. *Eur J Cardiovasc Prev Rehabil* 2009;16:562–70.
- 14 Martins D, Wolf M, Pan D, *et al.* Prevalence of cardiovascular risk factors and the serum levels of 25-hydroxyvitamin D in the United States: data from the Third National Health and Nutrition Examination Survey. *Arch Intern Med* 2007;167:1159–65.
- 15 Scragg R, Wishart J, Stewart A, *et al.* No effect of ultraviolet radiation on blood pressure and other cardiovascular risk factors. *J Hypertens* 2011;29:1749–56.
- 16 Shah AS, Langrish JP, Nair H, *et al.* Global association of air pollution and heart failure: a systematic review and meta-analysis. *Lancet* 2013;382:1039–48.
- 17 Kasper S, Wehr TA, Bartko JJ, *et al.* Epidemiological findings of seasonal changes in mood and behavior. A telephone survey of Montgomery County, Maryland. *Arch Gen Psychiatry* 1989;46:823–33.
- 18 Heneghan HM, Meron-Eldar S, Brethauer SA, *et al.* Effect of bariatric surgery on cardiovascular risk profile. *Am J Cardiol* 2011;108:1499–507.
- 19 Sarkhosh K, Birch DW, Shi X, *et al.* The impact of sleeve gastrectomy on hypertension: a systematic review. *Obes Surg* 2012;22:832–7.
- 20 Matthews CE, Freedson PS, Hebert JR, *et al.* Seasonal variation in household, occupational, and leisure time physical activity: longitudinal analyses from the seasonal variation of blood cholesterol study. *Am J Epidemiol* 2001;153:172–83.
- 21 Ma Y, Olendzki BC, Li W, *et al.* Seasonal variation in food intake, physical activity, and body weight in a predominantly overweight population. *Eur J Clin Nutr* 2006;60:519–28.
- 22 Shahar DR, Yerushalmi N, Lubin F, *et al.* Seasonal variations in dietary intake affect the consistency of dietary assessment. *Eur J Epidemiol* 2001;17:129–33.
- 23 Brennan PJ, Greenberg G, Miall WE, *et al.* Seasonal variation in arterial blood pressure. *Br Med J (Clin Res Ed)* 1982;285:919–23.
- 24 Alperovitch A, Lacombe JM, Hanon O, *et al.* Relationship between blood pressure and outdoor temperature in a large sample of elderly individuals: the Three-City study. *Arch Intern Med* 2009;169:75–80.
- 25 Cuspidi C, Ochoa JE, Parati G. Seasonal variations in blood pressure: a complex phenomenon. *J Hypertens* 2012;30:1315–20.
- 26 Peng RD, Dominici F, Pastor-Barriuso R, *et al.* Seasonal analyses of air pollution and mortality in 100 US cities. *Am J Epidemiol* 2005;161:585–94.
- 27 Nazir DJ, Roberts RS, Hill SA, *et al.* Monthly intra-individual variation in lipids over a 1-year period in 22 normal subjects. *Clin Biochem* 1999;32:381–9.
- 28 Ehret GB, Munroe PB, Rice KM, *et al.*, Internati onal Consortium for Blood Pressure Genome-Wide Association S. Genetic variants in novel pathways influence blood pressure and cardiovascular disease risk. *Nature* 2011;478:103–9.
- 29 Hadaegh F, Harati H, Zabetian A, *et al.* Seasonal variability of serum lipids in adults: Tehran Lipid and Glucose Study. *Med J Malaysia* 2006;61:332–8.
- 30 Catapano AL, Reiner Z, De Backer G, *et al.* ESC/EAS Guidelines for the management of dyslipidaemias The Task Force for the management of dyslipidaemias of the European Society of Cardiology (ESC) and the European Atherosclerosis Society (EAS). *Atherosclerosis* 2011;217:3–46.
- 31 Tung P, Wiviott SD, Cannon CP, *et al.* Seasonal variation in lipids in patients following acute coronary syndrome on fixed doses of Pravastatin (40 mg) or Atorvastatin (80 mg) (from the Pravastatin or Atorvastatin Evaluation and Infection Therapy-Thrombolysis In Myocardial Infarction 22 [PROVE IT-TIMI 22] Study). *Am J Cardiol* 2009;103:1056–60.
- 32 Kershenbaum A, Kershenbaum A, Tarabeia J, *et al.* Unraveling seasonality in population averages: an examination of seasonal variation in glucose levels in diabetes patients using a large population-based data set. *Chronobiol Int* 2011;28:352–60.
- 33 MacDonald MJ, Liston L, Carlson I. Seasonality in glycosylated hemoglobin in normal subjects. Does seasonal incidence in insulin-dependent diabetes suggest specific etiology? *Diabetes* 1987;36:265–8.
- 34 Reavey M, Saner H, Paccaud F, *et al.* Exploring the periodicity of cardiovascular events in Switzerland: Variation in deaths and hospitalizations across seasons, day of the week and hour of the day. *Int J Cardiol* 2013;168:2195–200.
- 35 Gemmell I, McLoone P, Boddy FA, *et al.* Seasonal variation in mortality in Scotland. *Int J Epidemiol* 2000;29:274–9.
- 36 Mikulich SK, Zerbe GO, Jones RH, *et al.* Comparing linear and nonlinear mixed model approaches to cosinor analysis. *Stat Med* 2003;22:3195–211.