Survival by stroke volume index in low-gradient normal EF severe aortic stenosis: insights into myocardial systolic dysfunction

To the Editor,

We read with interest the recent article on survival by stroke volume index (SVI) in patients with low-gradient normal EF severe aortic stenosis, which demonstrated lower SVI is incrementally associated with mortality.¹

The authors discuss a putative mechanism of low stroke volume (SV) secondary to concentric remodelling which results in reduced LV cavity size. This impedes LV diastolic filling culminating in diminished systolic function despite normal EF. The authors quote evidence of systolic impairment, for example reduced longitudinal strain, in similar cohorts with preserved EF.² In the current study, the subgroup with lowest SVI, and therefore presumed most severe systolic impairment despite EF >50%, demonstrated the thickest relative wall measurements. We believe this observation helps to explain the apparent paradox between significant myocardial dysfunction and preservation of EF in this cohort and in the wider 'heart failure with preserved ejection fraction' context. Recent mathematical modelling of LV contraction has shown that both myocardial shortening and end-diastolic wall thickness are determinants of EF.³ Essentially, absolute LV wall thickening, as defined by the absolute difference between wall thickness at end-systole and end-diastole, may be nearly normal in patients with concentric LV hypertrophy (LVH) because absolute systolic thickening will be augmented in response to increased end-diastolic LV wall thickness. As a result, the endocardial displacement and EF will also be normal, as the external LV volume remains fairly constant throughout the cardiac cycle⁴ and the absolute wall thickening may appear to compensate for any contractile strain abnormality. The development of concentric LVH may be viewed as a compensatory response that normalises contractile stress and total contractile force. However, if contractile stress remains reduced, the contractile force will be inadequate and result in a fall in stroke volume despite the preserved EF. In order to understand the apparent discrepancy in SV and EF, one must distinguish between contractile strain and stress and the relationship between end-diastolic wall thickness and EF.

The authors elected to investigate SV indexed to body surface area. However, it would be interesting to know whether correcting EF for the presence of concentric LVH (EF_c), as described in mathematical modelling studies of the LV³ would be a useful prognostic marker in this cohort of patients. After all, EF_c is potentially an even more relevant allometric indexed value given the importance of end-diastolic wall thickness in patients with concentric LVH and systolic impairment but preserved EF.

Jonathan C L Rodrigues,^{1,2}
Amardeep Ghosh Dastidar,¹ Stephen Rohan,³
David H MacIver^{4,5}

¹CMR Department, NIHR Bristol Cardiovascular Biomedical Research Unit, Bristol Heart Institute, Bristol, UK

²School of Physiology and Pharmacology, The University of Bristol, Bristol, UK

³Medical School, The University of Bristol, Bristol, UK

⁴Department of Cardiology, Taunton and Somerset Hospital, Taunton, UK

⁵Department of Biological Physics, The University of Manchester, Manchester, UK

Correspondence to Dr Jonathan C L Rodrigues, CMR Unit, NIHR Bristol Cardiovascular Biomedical Research Unit, Bristol Heart Institute, Bristol Royal Infirmary, Upper Maudlin Street, Bristol BS2 8HW, UK; jonathan.rodrigues@uhbristol.nhs.uk

Acknowledgements NIHR Bristol Cardiovascular Biomedical Research Unit. The views expressed are those of the authors and not necessarily those of the National Health Service, National Institute for Health Research or Department of Health.

Contributors JCLR: contributed to conception of the work, drafted the manuscript, approved the final version, agreed to be accountable for all aspects of the work.

AGD and DHM: contributed to conception of the work, reviewed the manuscript, approved the final version, agreed to be accountable for all aspects of the work. SR: reviewed the manuscript, approved the final version, agreed to be accountable for all aspects of the work.

Competing interests None.

Provenance and peer review Not commissioned; internally peer reviewed.

CrossMark

To cite Rodrigues JCL, Ghosh Dastidar A, Rohan S, et al. *Heart* 2015;**101**:83–84.

Published Online First 28 October 2014

► <http://dx.doi.org/10.1136/heartjnl-2014-306151>

Heart 2015;**101**:83–84.
doi:10.1136/heartjnl-2014-306933

REFERENCES

- 1 Eleid MF, Sorajla P, Michelena HI, et al. Survival by stroke volume index in patients with low-gradient normal EF severe aortic stenosis. *Heart* 2015;**101**: 23–9.
- 2 Adda J, Mielot C, Giorgi R, et al. Low-flow, low-gradient severe aortic stenosis despite normal ejection fraction is associated with severe left ventricular dysfunction as assessed by speckle-tracking echocardiography: a multicenter study. *Circ Cardiovasc Imaging* 2012;**5**:27–35.
- 3 Maciver DH. A new method for quantification of left ventricular systolic function using a corrected ejection fraction. *Eur J Echocardiogr* 2011;**12**:228–34.
- 4 Emilsson K, Brudin L, Wandt B. The mode of left ventricular pumping: is there an outer contour change in addition to atrioventricular plane displacement? *Clin Physiol* 2001;**21**:437–46.