Effect of a new beta-adrenergic blocking agent, ICI 66082, on exercise haemodynamics and airway resistance in angina pectoris

Hans Åström and Hans Vallin

From the Department of Clinical Physiology, Thoracic Clinics, Karolinska Sjukhuset, Stockholm, Sweden

A new cardioselective beta-adrenergic blocking agent, ICI 66082, was studied in 10 patients with severe effort-limiting angina pectoris, of whom all but one later went to bypass operation. The effect on maximal working capacity and airways resistance was compared before and after a single intravenous dose of 5 mg ICI 66082 and propranolol. The haemodynamic effects of ICI 66082 were also studied during cardiac catheterization in the same patients during supine exercise provoking chest pain.

Propranolol and ICI 66082 were equipotential in their blocking effect on exercise heart rate. In contrast to propranolol, ICI 66082 did not block the beta-receptors of the lung. Maximal working capacity was higher after ICI 66082 than after propranolol (+9%) and compared to the control period (+19%). This could be explained by a decrease found, after ICI 66082 administration, in both cardiac output (stroke volume unchanged) and aortic pressure during exercise with no change in the peripheral vascular resistance.

The beneficial effect of beta-adrenergic blockade in the treatment of angina pectoris is well documented (Epstein and Braunwald, 1966). Since propranolol was introduced, several other beta-adrenergic blocking agents with slightly different pharmacological properties have been introduced, in an effort to avoid the side effects of generalized beta-blockade, particularly with respect to the beta-receptors in the lung. Propranolol has been shown to have a significant therapeutic effect in hypertension, in spite of its beta-blocking effect on peripheral vessels (Prichard and Gillam, 1964). Theoretically this peripheral effect should hinder the achievement of an immediate pressure lowering effect.

Preliminary animal studies have shown that ICI 66082, 4-(2-hydroxy-3-isopropylaminopropoxy)-phenyl acetamide, blocks the beta-receptors specifically in the heart (Barrett *et al.*, 1973). It does not block peripheral vascular receptors and possesses no intrinsic sympathicomimetic or membrane stabilizing property. Furthermore it does not cross the blood-brain barrier. These qualities should be of advantage in the treatment of angina pectoris and hypertension. The findings in the present study suggest that the cardioselective ICI 66082 has such advantages as compared to propranolol in the treatment of angina pectoris.

Received 22 April 1974.

Subjects, methods, and procedure

Ten male subjects, aged 36 to 62 years, were investigated. All had effort induced angina pectoris with significant coronary artery stenosis verified with selective coronary angiography. Some clinical and anthropometric data are listed in Table 1. All but one later went to bypass operation because of intractable angina pectoris. All medication was withdrawn at least 2 days before the present investigations. None was on digitalis or diuretics and none had hypertension or diabetes.

Work tests were performed on a bicycle ergometer with increments of 10 Ws every minute from an initial work load of 10 Ws. All subjects exercised to their tolerance limits for chest pain and this was called W_{max} . A six-lead chest electrocardiogram was recorded during the whole exercise procedure. The heart rate was calculated from the electrocardiogram during the last 10 seconds at every load. This work test was repeated on two separate days after administration of 5 mg propranolol and ICI 66082 i.v. in a random order. Ten minutes after this administration and immediately before the exercise test the airway resistance was measured with a body plethysmograph (Brunes and Holmgren, 1966). All exercise tests were performed in the morning at least 1 hour after a light breakfast on the same electrically braked bicycle ergometer. The reproducibility of a single W_{max} test calculated from duplicate determinations performed within 4 days in another group of patients with similar severity of coronary artery disease has proven to be 6.6 per cent in our laboratory. Thus

TABLE 1 Anthropometric, clinical, and functional data in 10 patients with angina pectoris

Case No.	Age (yr)	Weight (kg)	BSA (m ²)	Time from infarction to investigation	Angina pectoris duration	Heart volume (ml/m ²)	$W_{\max}$ sitting (Watts)	Work supine at catheterization (Watts)
1	56	77	1.94	—	7 yr	400	—	20
2	45	63	1.74	7 mth	1 yr	420	80	20
3	43	83	2.04	2 yr	2 yr	420	70	10
4	36	65	1.77	—	3 mth	420	90	10
5	56	75	1.81	1 yr	10 yr	460	100	20
6	62	85	2.03	—	3 yr	630	—	20
7	45	80	1.95	—	2 yr	370	100	20
8	59	91	2.14	—	1 yr	470	90	10
9	49	77	1.86	—	1 yr	370	150	20
10	48	82	1.99	1 yr	4 yr	330	70	10

differences in $W_{\max}$ of more than 10 Ws (i.e. one step) is expected to be significant at work rates around 100 Ws.

The last step of the investigation was a right heart catheterization, performed with a double-lumen catheter introduced percutaneously in a right cubital vein. The left heart catheterization was performed via a 'teflon' catheter introduced percutaneously in the right brachial artery. The cardiac output was determined with the direct Fick method. The rate-pressure index was calculated as the product of systolic pressure in aorta and heart rate (Robinson, 1967). pH, carbon dioxide, and base excess were determined with the micro-equilibration technique. The techniques and procedures were the same as described previously in more detail from our laboratory (Åström, 1968).

Heart catheterization was performed in the morning after oral premedication with 0.8 g meprobamate. The subjects were studied at rest and during work in supine position. The work load had been determined previously to induce moderate chest pain but still work could be continued for 6 to 8 minutes. After a resting period of 30 to 40 minutes, 5 mg ICI 66082 was given in the pulmonary artery and 10 minutes later the procedure was repeated.

The nature and purpose of the investigation were explained to the patients and their consent obtained.

Results

Working capacity ($W_{\max}$)

Eight patients performed a maximal exercise test before and after both ICI 66082 and propranolol. Fig. 1 shows the close linear correlation between work load in 1-minute steps and heart rate. The $W_{\max}$ in the control period was on an average 93.8 ± 9.1 (SEM) Ws and increased after administration of ICI 66082 to 111.3 ± 7.7 Ws ($P < 0.05$) and after propranolol to 102.5 ± 8.8 Ws (n.s.). The difference between the two drugs in the effect on $W_{\max}$ was significant, while the maximal heart rate

FIG. 1 Relation between mean values for heart rate (HR) and work load at 1-minute step exercise before and after 5 mg of ICI 66082 and propranolol. Symbols with circle represent mean maximal point. Individual control values in Table 1.

was equal (Fig. 2). After both drugs the subjects stopped the work because of chest pain but at a lower heart rate (Fig. 2).

Airway resistance (Fig. 3)

Airway resistance was within the normal range in most patients. After ICI 66082 the resistance decreased by 7 per cent ($P < 0.05$) and the airway conductance per litre thoracic gas volume increased by 15 per cent ($P < 0.01$). It is noteworthy that the patients with signs of slight airway obstruction (i.e. control airway resistance above 3.0 cm H₂O/l. per sec. or specific airway conductance below 0.12 l./sec per cm H₂O) all had an increase in airway conductance

FIG. 2 Maximal power (W_{MAX}) and maximal heart rate (HR_{MAX}) performed before and after ICI 66082 and propranolol. Open circles represent mean value.

FIG. 3 Airway resistance (R_a) and airway conductance per litre thoracic gas volume (C_A/TGV) before and after ICI 66082 and propranolol.

TABLE 2 Mean values for haemodynamic variables before and after 5 mg ICI 66082 i.v. for individual differences and significance of differences

	<i>O₂</i> <i>uptake</i>	<i>Heart rate</i>	<i>Arterio- venous diff</i>	<i>Cardiac output</i>	<i>Stroke volume</i>	<i>Pressure (mmHg) (mean)</i>			<i>Rate-pressure</i>
	(ml/min)	(beats/min)	(ml/l.)	(l./min)	(ml)	<i>Aorta</i>	<i>PCV</i>	<i>PA</i>	<i>index</i> × 10 ⁻²
<i>Rest</i>									
Before	257	83	46	5.7	70	97	10	16	97.6
After	243	69	53	4.6	68	95	11	17	79.9
Difference	14	14	-7	1.1	2	2	-1	-1	17.7
P	<0.02	<0.001	<0.01	<0.001	NS	NS	NS	NS	<0.001
<i>Exercise</i>									
Before	601	101	77	8.0	79	115	22	32	146.9
After	592	85	91	6.7	79	103	19	28	110.5
Difference	9	16	-14	1.3	0	12	3	4	36.4
P	NS	<0.001	<0.01	<0.001	NS	<0.001	NS	NS	<0.001

P=probability, paired t-test.

FIG. 4 Stroke volume (SV), cardiac output ($\dot{Q}$), and heart rate (HR) before and after ICI 66082 in 10 patients with angina pectoris at rest (○) and during exercise (●). Square with circle represents mean value.

after ICI 66082 but a further decrease after the same amount of propranolol was administered.

Haemodynamics (Table 2)

Oxygen consumption was slightly reduced after ICI 66082 at rest ($P < 0.05$) but was unchanged

during exercise. All subjects could work without chest pain after ICI 66082 at that work load which caused pain before treatment. This fact did not affect the total oxygen uptake and thus not the mechanical efficiency. Cardiac output at rest was diminished by 19 per cent and during work by 15 per cent. This change was caused by a decrease in

FIG. 5 Aortic mean pressure ($\bar{P}_{AO}$), rate-pressure index, and pulmonary artery wedge pressure ($\bar{P}_{PCV}$) before and after ICI 66082. Symbols as in Fig. 4.

heart rate alone as stroke volume was unchanged both at rest and during work (Fig. 4). At rest the pressures in aorta, pulmonary artery wedge, pulmonary artery, and right atrium were all within normal range, and no significant change was found after ICI 66082. During exercise before beta-blockade provoking angina pectoris in all subjects the wedge pressure increased abnormally. The magnitude of this increase was not significantly changed after blockade. However, in 3 patients with wedge pressure above 25 mmHg this pressure decreased after blockade by an average of 12 mmHg (Fig. 5). The pulmonary artery mean pressure was also unchanged after blockade. The aortic pressure increased during exercise but the increment after ICI 66082 was significantly less (12.6 mmHg, $P < 0.001$) than during control period. The systemic vascular resistance during work was unchanged by drug administration. The rate-pressure index decreased by 18 per cent at rest and 25 per cent at work ($P < 0.001$). There was no difference found in pH, base excess, or carbon dioxide either at rest or during work.

Discussion

All patients in this study had effort induced angina pectoris and all but one later went to bypass opera-

tion. A standardized multistage exercise test was used to evaluate the tolerance limit for angina pectoris. From this test it is also seen that ICI 66082 and propranolol are equipotential concerning the blocking effect on heart rate increase during exercise (Fig. 3).

In the present study the maximal work rate was higher after ICI 66082 than after propranolol. The relative bradycardia was the same for both agents. The difference in W_{max} might be explained by their different action on the arterial pressure, as this, together with heart rate, is one of the major determinants for the myocardial oxygen consumption (Sarnoff *et al.*, 1958). The rate-pressure index decreased by 25 per cent after ICI 66082. From an earlier study on the haemodynamic effects of propranolol (Åström, 1968), a recalculation of the rate-pressure index showed a decrease of 15 per cent (Fig. 6). It was recently shown that the rate-pressure product was a better index of the myocardial oxygen consumption than the tension-time index despite a change in contractility induced by propranolol (Jorgensen *et al.*, 1973).

The aortic pressure was unchanged at rest, with an increase in the peripheral vascular resistance after blockade with ICI 66082. During exercise, however, the resistance was unchanged with a

lower aortic pressure related to the decrease in cardiac output. Propranolol increases the peripheral vascular resistance both at rest and during exercise when given acutely as it blocks the beta-adrenergic vasodilator receptors (Hamer and Sowton, 1965). The cardioselective beta-blocking agent, practolol, has been reported not to change systemic arterial pressure nor peripheral resistance during erect exercise in normals (Gibson and Sowton, 1968). In patients with coronary heart disease practolol has been reported to decrease the arterial pressure during exercise, inducing angina pectoris (Rousseau, Brasseur, and Detry, 1973).

exercise the stroke volume was unchanged after this new beta-blocking agent as also found after propranolol (Åström, 1968).

ICI 66082 did not increase airway resistance, particularly not in patients with high resistance during the control period. This is in contrast to the non-selective beta-blocking agent propranolol (McNeill, 1964). The actual slight decrease in airway resistance after ICI 66082 might be explained by a reflex beta-stimulation of the lungs elicited by the haemodynamic changes. This implies less risk if the drug be given to asthmatic subjects.

In conclusion ICI 66082 has proved to be a cardioselective beta-adrenergic blocking agent with apparently no adverse effect on respiratory mechanics. Its haemodynamic action seems similar to that of practolol, with no effect on the peripheral vascular resistance during exercise. ICI 66082 is supposed not to enter the brain and would thus give less side-effects such as troublesome dreams sometimes produced by propranolol and practolol (Wiseman, 1971).

We are grateful to P. E. Hörnkvist, ICI-Pharma, Sweden for supplies of ICI 66082.

References

- Åström, H. (1968). Haemodynamic effects of beta-adrenergic blockade. *British Heart Journal*, **30**, 44.
- Barrett, A. M., Carter, J., Fitzgerald, J. D., Hull, R., and Le Count, D. (1973). A new type of cardioselective adrenoceptive blocking drug. *British Journal of Pharmacology*, **48** (2), 340 P.
- Brunes, L., and Holmgren, A. (1966). Total airway resistance and its relationship to body size and lung volumes in healthy young women. *Scandinavian Journal of Clinical and Laboratory Investigation*, **18**, 316.
- Epstein, S. E., and Braunwald, E. (1966). Beta-adrenergic receptor blocking drugs. Mechanisms of action and clinical applications. *New England Journal of Medicine*, **275**, 1106.
- Gibson, D., and Sowton, E. (1968). Effects of ICI 50172 in man during erect exercise. *British Medical Journal*, **1**, 213.
- Gibson, D. G., and Coltart, D. J. (1972). Haemodynamic effects of practolol. *British Heart Journal*, **34**, 95.
- Hamer, J., and Sowton, E. (1965). Cardiac output after beta-adrenergic blockade in ischaemic heart disease. *British Heart Journal*, **27**, 892.
- Jorgensen, C. R., Wang, K., Wang, Y., Gobel, F. L., Nelson, R. R., and Taylor, H. (1973). Effect of propranolol on myocardial oxygen consumption and its hemodynamic correlates during upright exercise. *Circulation*, **48**, 1173.
- McNeill, R. S. (1964). Effect of beta-adrenergic-blocking agent on asthmatics. *Lancet*, **2**, 1101.
- Prichard, B. N. C., and Gillam, P. M. S. (1964). The use of propranolol in the treatment of hypertension. *British Medical Journal*, **2**, 725.
- Robinson, B. F. (1967). Relation of heart rate and systolic blood pressure to the onset of pain in angina pectoris. *Circulation*, **35**, 1073.
- Rousseau, M. F., Brasseur, L. A., and Detry, J.-M. R. (1973). Haemodynamic and electrocardiographic effects of practolol during upright exercise in coronary heart disease. *Cardiovascular Research*, **7**, 306.
- Sarnoff, S. J., Braunwald, E., Welch, G. H., Jr., Case, R. B., Stainsby, W. N., and Macruz, R. (1958). Hemodynamic determinants of oxygen consumption of the heart with special reference to the tension-time index. *American Journal of Physiology*, **192**, 148.
- Wiseman, R. A. (1971). Practolol - accumulated data on unwanted effects. *Postgraduate Medical Journal*, **47**, January, Suppl. 68.

Requests for reprints to Dr. Hans Åström, Department of Clinical Physiology, Thoracic Clinics, Karolinska Sjukhuset, S-104 01 Stockholm 60, Sweden.