

British Heart Journal

ogy
h

<i>David Short</i> Maximal exercise testing in patients with spontaneous angina pectoris associated with transient ST segment elevation. Risks and electrocardiographic findings <i>J.-M. R. Detry, P. Mengeot, M. F. Rousseau, J. Cosyns, R. Ponlot, and L. A. Brasseur</i>	on 893	Variations in clinical presentation of Fallot's tetralogy in infancy. Angiographic and pathogenetic implications <i>E. A. Shinebourne, R. H. Anderson, and Jean J. Bowyer</i>	946
Continuous assessment of left ventricular shape in man <i>D. G. Gibson and D. J. Brown</i>	897	Comparative study of sinoatrial conduction time and sinus node recovery time <i>G. Steinbeck and B. Lüderitz</i>	956
Non-invasive left ventricular volume determination by two-dimensional echocardiography <i>J. Gehrke, S. Leeman, M. Raphael, and R. B. Pridie</i>	904	Localization of left ventricular ischaemia in angina pectoris by cineangiography during exercise <i>B. Sharma and S. H. Taylor</i>	963
Hypertrophic obstructive cardiomyopathy. Assessment by echocardiographic and Doppler ultrasound techniques <i>D. R. Boughner, R. L. Schuld, and J. A. Persaud</i>	911	Echocardiographic left ventricular dimensions in pressure and volume overload. Their use in assessing aortic stenosis <i>D. H. Bennett, D. W. Evans, and M. V. J. Raj</i>	971
Long-term survival of patients with incomplete bundle-branch block complicating acute myocardial infarction <i>E. Lichstein, P. K. Gupta, and K. D. Chadda</i>	917	Case reports Effect of coexistent coarctation of pulmonary trunk in natural history of complete absence of pulmonary valve with ventricular septal defect <i>T. Peter, R. Harper, J. Vohra, and D. Hunt</i>	978
Surgical implications of right aortic arch with isolation of left subclavian artery <i>L. Rodriguez, T. Izukawa, C. A. F. Moës, G. A. Trusler, and W. G. Williams</i>	924	Complete heart block and systemic lupus erythematosus <i>P. Wray and M. Iveson</i>	982
Coarctation of aorta with right aortic arch. Report of surgical correction in 2 cases: one with associated anomalous origin of left circumflex coronary artery from the right pulmonary artery <i>M. Honey, J. C. R. Lincoln, M. P. Osborne, and D. P. de Bono</i>	931	Significance of exercise-induced second degree atrioventricular block <i>A. Bakst, B. Goldberg, and L. Schamroth</i>	984
		Correspondence	987
		Proceedings of the British Cardiac Society. Private business	989
		Notices	992
	937	British Medical Association	