Two-stage replacement of tricuspid valve in active endocarditis

PER ARNEBORN, VIKING OLOV BJÖRK, LUIS RODRIGUEZ,
AND MAJ SVANBOM

From the Thoracic Surgical Clinic, Karolinska sjukhuset, and the Hospital for Infectious Diseases (Roslagstulls sjukhus), Stockholm, Sweden

One month after a 21-year-old female drug addict with tricuspid staphylococcal endocarditis seemed to have been cured by antibiotic therapy she relapsed. Further antibiotic treatment for one month failed to control the infection. Tricuspidectomy was then performed and the infection was eliminated. A valve prosthesis was inserted 3 months later. She recovered and is now fully restored to health. We recommend this two-stage surgical technique in similar cases.

Right-sided endocarditis, mainly involving the tricuspid valve, is no longer a rare disease, its frequency increasing from less than 5 per cent (Thayer, 1931; Bain *et al.*, 1958) to 10 to 20 per cent in the past decade (Roberts and Buchbinder, 1972; Arbulu *et al.*, 1973); this can be ascribed largely to the increased incidence of narcotic addiction (Conway, 1969; Banks *et al.*, 1973; Graham *et al.*, 1973; Svanbom *et al.*, 1973).

Valve replacement during the active stage of endocarditis is now an accepted form of treatment in many places (Strandell *et al.*, 1969; Manhas *et al.*, 1972; Graham *et al.*, 1973; Black *et al.*, 1974), but not in all. Good results have been obtained in many cases of tricuspid valve endocarditis by means of valve resection alone (Arbulu *et al.*, 1973; Graham *et al.*, 1973; Michl *et al.*, 1973) or with the insertion of an artificial valve at a second operation (Arbulu *et al.*, 1973; Simberkoff *et al.*, 1974).

Case report

The patient was a 21-year-old woman without known heart disease, who had taken narcotics intravenously for about 4 years. Some weeks before the onset of the present illness, she had an infection in one foot after treading on a piece of glass.

- On 14 June 1974 during a holiday abroad, she fell ill with malaise and fever above 40°C and was treated with antibiotics. On returning to Sweden on 24 June, she was admitted to Roslagstull Hospital for persisting high fever.

On admission she was febrile, but had no signs of cardiac failure or peripheral emboli. A systolic

murmur grade 2/6 was heard over the apex. The liver and spleen were enlarged. X-ray film showed infiltrations in the right middle and left lower lung lobes. Electrocardiogram was normal apart from sinus tachycardia. Phonocardiogram showed a faint systolic murmur of medium frequency, thought, however, to be consistent with a normal flow in the pulmonary artery.

On the third day in hospital a growth of penicillinase-producing *Staphylococcus aureus* from blood was reported and cloxacillin plus benzyl penicillin were administered. On the next day, she became dyspnoeic and x-ray examination showed a pleural effusion. Despite changes in antibiotic therapy, the lung shadows increased with cavity formation in the right lower lobe.

On 15 July, transient murmurs and electrocardiographic abnormalities suggestive of pericarditis were noted and pulmonary scintigraphy indicated changes consistent with multiple small emboli. Right-sided cardiac catheterisation showed normal pressures. Angiocardiography defined a walnut-sized mass on the tricuspid valve, with extreme regurgitation.

After further antibiotic treatment, the patient was discharged on 1 October, afebrile and asymptomatic, but was readmitted on 28 October with fever and cough. A chest x-ray film showed further infiltration and scintigraphy revealed fresh infarction of the left lower lobe. The heart size had increased from 300 to 400 ml per m² body surface area. The electrocardiogram was initially normal but signs of right ventricular strain appeared later. Though repeated blood cultures were negative, carbenicillin

and gentamicin were given and the temperature subsided.

On 25 November, the tricuspid valve was resected under total cardiopulmonary bypass at the Thoracic Surgical Clinic, Karolinska sjukhuset, Stockholm. Large vegetations were seen on the posterior leaflet with acute inflammatory reaction. Bacteria were not seen on microscopy or cultured. Increased pulsation of the jugular veins and the liver was noted after operation. Right heart catheterisation showed a low stroke volume, normal pressures in the pulmonary capillary venous and pulmonary arterial positions, and a high right ventricular end-diastolic pressure. Heart size was now 510 ml per m² body surface area. She was dyspnoeic on walking but otherwise felt well.

On account of the persistent right-sided failure, a 31 mm Björk-Shiley mitral valve prosthesis was inserted on 12 February 1975, and 15 days later she was discharged. One year after operation, she was in good health, working full-time, and had resumed water-skiing.

Discussion

The combination of narcotic addiction, staphylococcal septicaemia with right-sided endocarditis, and pulmonary embolism was first described by Hussey *et al.* (1944), and has often been reported since then (Hussey and Katz, 1950; Olsson and Romansky, 1962; Conway, 1969; Roberts and Buchbinder, 1972; Menda and Gorbach, 1973; Svanbom *et al.*, 1973). These authors drew attention to the possibility of tricuspid valve involvement. In this form of endocarditis cardiac symptoms are initially often obscure (Bain *et al.*, 1958; Conway, 1969; Arbulu *et al.*, 1973; Graham *et al.*, 1973), and the diagnosis and treatment may be delayed. Later in the course of the disease, typical signs of tricuspid regurgitation (Banks *et al.*, 1973) and increasing heart size are frequent. These symptoms were observed in our patient and the diagnosis was also verified by right-sided ventriculography.

Staphylococci are still the most common agents in this connection (Roberts and Buchbinder, 1972; Banks *et al.*, 1973; Graham *et al.*, 1973; Menda and Gorbach, 1973), but in recent years other organisms, particularly Gram-negative bacilli, have also been implicated (Carruthers and Kanokvechayant, 1973; Graham *et al.*, 1973; Rosenblatt *et al.*, 1973). Mixed infections have also been reported (Menda and Gorbach, 1973; Simberkoff *et al.*, 1974).

Staphylococcal tricuspid endocarditis may respond to antibiotic therapy alone (Arbulu *et al.*, 1973; Menda and Gorbach, 1973), and initially this seemed to apply in our case as the patient was dis-

charged apparently without active infection, after 10 weeks of antibiotic treatment. Her second illness with fever and pulmonary emboli suggested reactivation of her endocarditis. The signs of acute inflammatory reaction in the excised valve supported this suggestion but the aetiology of the relapse remained obscure since repeated blood cultures were negative and examination of the excised valve gave no indication of the causative agent.

Infectious foci in heart valves are difficult to treat. The bacteria may acquire resistance to the antibiotics, or L-forms of the bacteria may develop. In our case, a superinfection may have occurred when the patient may have been giving herself injections. A 'mixed' infection with Gram-negative bacteria in addition to staphylococci (Simberkoff *et al.*, 1974) may have developed. The favourable effect of gentamicin and carbenicillin supports this.

The primary treatment of bacterial endocarditis is conservative. Surgical intervention was used only for patients with haemodynamic failure after healed endocarditis (Yeh *et al.*, 1964), until Wallace *et al.* (1965) for the first time operated during the active stage of endocarditis when conservative treatment had failed. Since then several similar reports have been published (Strandell *et al.*, 1969; Manhas *et al.*, 1972; Graham *et al.*, 1973; Black *et al.*, 1974). The favourable results tally with the well-known fact that the elimination of infected foci is important for the successful treatment of septicaemia.

Valve replacement during active endocarditis is not always successful, and there are reports of paraprosthetic endocarditis and failure of sutures (Manhas *et al.*, 1972). Consequently, a decision on surgical intervention must not be premature, and should only be undertaken when antibiotic treatment has been shown to be ineffective. On the other hand, the probabilities of left heart involvement and toxic myocarditis increase with time. Arbulu *et al.* (1973) recommend surgical intervention when antibiotic treatment has not healed the infection within 6 weeks.

The small pressure differences within the right heart make damage to its valvular system less detrimental than such damage to the left heart. Traumatic tricuspid damage, which persisted for several decades without any symptoms, has been described previously (Croxxson *et al.*, 1971; Morgan and Forker, 1971; Marvin *et al.*, 1973). In the series of Arbulu (Arbulu *et al.*, 1973) during a follow-up time of 3 to 27 months, only 1 of 10 survivors required insertion of a prosthetic valve. However, experience from animal experiments (Spellman and Balkissoon, 1956; Michl *et al.*, 1973), from traumatic tricuspid regurgitation, and from patients with valve resections (Michl *et al.*, 1973; Robin *et*

al., 1974), indicates that right heart failure may develop sooner or later.

Adaptation to tricuspid regurgitation in young and otherwise healthy persons with normal pulmonary resistance may be considerable, but the insertion of a prosthesis could result in completely normal haemodynamic function. Our patient is now working full time and water skis, which is indicative of good haemodynamic function.

We consider that tricuspid valve replacement is safe when a low profile prosthesis with optimal haemodynamic properties, such as the Björk-Shiley valve, is used.

References

- Arbulu, A., Kafi, A., Thoms, N. W., and Wilson, R. F. (1973). Right-sided bacterial endocarditis. *Annals of Thoracic Surgery*, **16**, 136-140.
- Bain, R. C., Edwards, J. E., Scheifley, C. H., and Geraci, J. E. (1958). Right-sided bacterial endocarditis and endarteritis. A clinical and pathologic study. *American Journal of Medicine*, **24**, 98-110.
- Banks, T., Fletcher, R., and Ali, N. (1973). Infective endocarditis in heroin addicts. *American Journal of Medicine*, **55**, 444-451.
- Black, S., O'Rourke, R. A., and Karliner, J. S. (1974). Role of surgery in the treatment of primary infective endocarditis. *American Journal of Medicine*, **56**, 357-369.
- Carruthers, M. M., and Kanokvechayant, R. (1973). *Pseudomonas aeruginosa* endocarditis. Report of a case, with review of the literature. *American Journal of Medicine*, **55**, 811-818.
- Conway, N. (1969). Endocarditis in heroin addicts. *British Heart Journal*, **31**, 543-545.
- Croxson, M. S., O'Brien, K. P., and Lowe, J. B. (1971). Traumatic tricuspid regurgitation. Long-term survival. *British Heart Journal*, **33**, 750-755.
- Graham, D. Y., Reul, G. J., Martin, R., Morton, J., and Kennedy, J. H. (1973). Infective endocarditis in drug addicts. Experiences with medical and surgical treatment. *Circulation*, **47** and **48**, Suppl. III, 37-41.
- Hussey, H. H., and Katz, S. (1950). Infections resulting from narcotic addiction. *American Journal of Medicine*, **9**, 186-193.
- Hussey, H., Keliher, T. F., Schaefer, B. F., and Walsh, B. J. (1944). Septicemia and bacterial endocarditis resulting from heroin addiction. *Journal of the American Medical Association*, **126**, 535-538.
- Manhas, D. R., Mohri, H., Hessel, E. A., and Merendino, K. A. (1972). Experience with surgical management of primary infective endocarditis. A collected review of 139 patients. *American Heart Journal*, **84**, 738-747.
- Marvin, R. F., Schrank, J. P., and Nolan, S. P. (1973). Traumatic tricuspid insufficiency. *American Journal of Cardiology*, **32**, 723-726.
- Menda, K. B., and Gorbach, S. L. (1973). Favourable experience with bacterial endocarditis in heroin addicts. *Annals of Internal Medicine*, **78**, 25-32.
- Michl, L., Shumacker, H. B., and Hobbs, D. (1973). Tricuspid valvectomy. *Surgery, Gynecology and Obstetrics*, **137**, 590-592.
- Morgan, J. R., and Forker, A. D. (1971). Isolated tricuspid insufficiency. *Circulation*, **43**, 559-564.
- Olsson, R. A., and Romansky, M. J. (1962). Staphylococcal tricuspid endocarditis in heroin addicts. *Annals of Internal Medicine*, **57**, 755-762.
- Roberts, W. C., and Buchbinder, N. A. (1972). Right-sided valvular infective endocarditis. A clinicopathologic study of twelve necropsy patients. *American Journal of Medicine*, **53**, 7-19.
- Robin, E., Belamaric, J., Thoms, N. W., Arbulu, A., and Ganguly, S. N. (1974). Consequences of total tricuspid valvectomy without prosthetic replacement in treatment of *Pseudomonas* endocarditis. *Journal of Thoracic and Cardiovascular Surgery*, **68**, 461-465.
- Rosenblatt, J. E., Dahlgren, J. G., Fishbach, R. S., and Tally, F. P. (1973). Gram-negative bacterial endocarditis in narcotic addicts. *California Medicine*, **118** (2), 1-4.
- Simberloff, M. S., Isom, W., Smithivas, T., Noriega, E. R., and Rahal, J. J. (1974). Two-stage tricuspid valve replacement for mixed bacterial endocarditis. *Archives of Internal Medicine*, **133**, 212-216.
- Spellman, M. W., and Balkissoon, B. (1956). The production and evaluation of ascites secondary to isolated tricuspid insufficiency in dogs. *Surgery*, **39**, 37-48.
- Strandell, T., Bengtsson, E., Björk, V. O., Michaeli, E., Svanbom, M., and Tunevall, G. (1969). Fulminant bakteriell endokardit med ruptur av aortaklaffarna, behandlad med aortaprotet under akut stadium. *Nordisk Medicin*, **82**, 1083-1084.
- Svanbom, M., Bengtsson, E., Strandell, T., and Tunevall, G. (1973). Septicemia in narcotic addicts. *Scandinavian Journal of Infectious Diseases*, **5**, 97-110.
- Thayer, W. S. (1931). Bacterial or infective endocarditis. *Edinburgh Medical Journal*, **38**, 237-265; 307-334.
- Wallace, A. C., Young, W. G., and Osterhout, S. (1965). Treatment of acute bacterial endocarditis by valve excision and replacement. *Circulation*, **31**, 450-453.
- Yeh, T. J., Hall, D. P., and Ellison, R. G. (1964). Surgical treatment of aortic valve perforation due to bacterial endocarditis. *American Surgeon*, **30**, 766-769.

Requests for reprints to Dr. Maj Svanbom, Roslagstulls sjukhus, Box 5901, S-114 89 Stockholm, Sweden.