Electrophysiological effects of lorcinide, a new antiarrhythmic drug

Observations in patients with and without pre-excitation

FRITS W BÄR, JERÓNIMO FARRÉ, DAVID ROSS, EDGAR J VANAGT, ANTON P GORGELS, HEIN J J WELLENS

From the Department of Cardiology, University of Limburg, Annadal Hospital, Maastricht, The Netherlands

SUMMARY The electrophysiological effects of the intravenous administration of a new antiarrhythmic drug, lorcinide, were evaluated by programmed electrical stimulation of the heart in 23 patients with atrioventricular conduction disturbances (four patients), ventricular tachycardia (five patients), and accessory atrioventricular pathway (14 patients). Lorcinide did not affect the refractory period of the atrium, ventricle, atrioventricular node, or the AH interval. It lengthened the duration of the HV interval, the refractory period of the accessory pathway, and the width of the QRS complex. The drug terminated ventricular tachycardia in four of five patients. It is concluded that the drug may be of potential benefit in patients with ventricular tachycardia or accessory atrioventricular pathways (especially those with a short refractory period).

Lorcinide is contraindicated in patients with bundle-branch block and prolonged HV interval.

Lorcinide (R 15 889) is a benzene-acetamide hydrochloride derivative. It is a new antiarrhythmic agent with local anaesthetic properties. The action of the drug can be partly explained by its selective reducing effect of sodium conductance.¹

Our study was performed to investigate the specific electrophysiological effects of lorcinide in man.

Subjects and methods

Twenty-three patients were studied.

Four patients (two men and two women) had atrioventricular conduction disturbances. The ages of these patients ranged from 33 to 79, with a mean of 57 years. The conduction disturbance was located at the level of the atrioventricular node in one patient, the His bundle in one patient, and the bundle-branch system in two patients.

Five patients (two men and three women) had ventricular tachycardia. Their ages were 22 to 69, mean 44 years. In two of these patients an old myocardial infarction was present. The other three were idiopathic. Twelve patients (10 male and two female) suffered from the Wolff-Parkinson-White syndrome.

Received for publication 12 May 1980

The age of these patients was 8 to 58, mean 36 years. In nine the accessory pathway was located on the left side of the heart and in three on the right side. Ten of these patients had circus movement tachycardia.

A further two patients (one man of 43 years and one woman of 33 years) had a left sided concealed bypass tract.

Our methods of recording and stimulation during programmed electrical stimulation of the heart have been described previously.²

Informed consent was obtained, and four to five wires were passed through the femoral vein using the Seldinger technique. Two bipolar (or one quadripolar) wire(s) were positioned high on the lateral wall of the right atrium. One was used for stimulation, the other for recording an intra-atrial electrogram. A bipolar wire was placed in the region of the bundle of His to record the His bundle electrogram. A fourth (bipolar) wire was positioned in the apex of the right ventricle and used for ventricular stimulation.

In 13 patients a left atrial electrogram was recorded by way of a fifth wire (bi- or quadripolar) in the coronary sinus. This wire was also used for left atrial stimulation. In two patients a left atrial

electrogram was also recorded from a wire in the main stem of the pulmonary artery.

In three patients with ventricular tachycardia a second bipolar wire was introduced in the ventricle: in two patients in the outflow tract of the right ventricle and in one patient in the left ventricle. Stimulation was performed at twice diastolic threshold both before and after lorcaïnide. All data obtained during the stimulation studies were recorded on a 16 channel Elema recorder. Leads I, II, III, V₁, and V₆, a right atrial lead, a left atrial lead, a His bundle recording, and sometimes a ventricular lead were recorded simultaneously.

Using the single test stimulus method during atrial stimulation the effective refractory period of the right atrium and the atrioventricular node, the functional refractory period of the atrioventricular node, the A₁H₁, the maximal A₂H₂, the H₁V₁, and the maximal H₂V₂ interval were determined. The width of the His bundle electrogram and QRS complex were also measured.

During single test stimulation of the right ventricle the mode of VA conduction, the effective refractory period of the right ventricle, QRS width, and V₁H₁ and V₁A₁ intervals were determined. In patients with ventricular tachycardia, the rate and tachycardia zone were measured. In patients with an accessory pathway the effective refractory period of the extra connection in the anterograde and retrograde directions, tachycardia zone, rate, and AH, HV, and VA intervals during tachycardia were measured. The modes of initiation and termination of tachycardia were determined in all patients with tachycardia.

Previously described criteria³ were applied to determine the pathway used during ventriculoatrial conduction and the pathway used during re-entrant tachycardia. Measurements were repeated immediately after and continuously during the first 15

minutes, and then at 30, 45, and 60 minute intervals after the intravenous injection of lorcaïnide in a dose of 2 mg/kg bodyweight given over a five-minute period. Under results only the maximal changes in electrophysiological variables after lorcaïnide administration will be given. In patients where the exact value of the effective refractory period of the accessory pathway could not be determined because conduction over that structure occurred up to the effective refractory period of the atrium or ventricle, the increase after lorcaïnide can only be an approximation. Statistical analyses were performed using the paired t test.

Results

ANTEROGRADE CONDUCTION OVER ATRIOVENTRICULAR NODE—HIS BUNDLE AXIS (Table 1)

During atrial pacing lorcaïnide had no significant effect on the effective refractory period of the right atrium, the effective refractory period of the atrioventricular node, the functional refractory period of the atrioventricular node, and the A₁H₁ and A₂H₂ intervals. The width of the His bundle electrogram increased by a mean of 2.7 ± 2.6 ms. In addition, the H₁V₁ interval increased by a mean of 22.7 ± 40.0 ms. The maximal H₂V₂ interval did not change significantly, but the QRS width during regular atrial pacing lengthened by a mean of 20.4 ± 14.4 ms. The 12 patients with anterograde conduction over the accessory pathway are not included in Table 1.

After drug administration right bundle-branch block developed in two patients.

VA (HIS-ATRIOVENTRICULAR NODE) CONDUCTION

These observations were made in the nine patients

Table 1 Effect of lorcaïnide on electrophysiological data during atrial pacing*

	No. of patients	Before		After		P
		Range (ms)	Mean $\pm$ SD (ms)	Range (ms)	Mean $\pm$ SD (ms)	
ERP _{RA}	6	190–280	242 $\pm$ 31	200–270	245 $\pm$ 29	NS
ERP _{AVN}	6	190–490	287 $\pm$ 104	200–430	272 $\pm$ 82	NS
FRP _{AVN}	6	300–520	403 $\pm$ 73	310–440	378 $\pm$ 52	NS
A ₁ H ₁	11	70–220	105 $\pm$ 44	70–160	103 $\pm$ 24	NS
A ₂ H ₂ maximal	6	150–330	237 $\pm$ 75	140–410	207 $\pm$ 103	NS
His	11	10–30	20 $\pm$ 6	10–35	22 $\pm$ 7	< 0.01
H ₁ V ₁	11	35–130	55 $\pm$ 27	40–270**	78 $\pm$ 65	NS
H ₂ V ₂ maximal	6	35–70	52 $\pm$ 14	40–75	71 $\pm$ 32	NS
QRS width	11	80–170	109 $\pm$ 34	85–210	129 $\pm$ 46	< 0.001

*Patients with Wolff-Parkinson-White syndrome are not included. Data on all parameters before and after drug administration were not available in all 11 patients.

**One patient developed complete HV block after lorcaïnide.

ERP, effective refractory period; RA, right atrium; AVN, atrioventricular node; FRP, functional refractory period.

without accessory pathways. No significant effect of lorainide was seen on the effective refractory period of the right ventricle. Before the drug the range was 230 to 290, mean 256 ± 25 ms, and after lorainide the range was 220 to 320, mean 260 ± 37 ms (NS). The QRS width during regular ventricular pacing increased. Before the drug the range was 120 to 190, mean 158 ± 23 ms, and after the drug it was 135 to 215 ms, mean 178 ± 25 ms ($p < 0.001$). VA delay developed after the drug in four out of the five patients showing retrograde conduction over the His-atrioventricular node pathway. Two of these four patients developed complete retrograde block below the His bundle.

ATRIOVENTRICULAR CONDUCTION DISTURBANCES

In the patients with atrioventricular nodal and intraHisian block lorainide had no effect on anterograde conduction.

This was in contrast to the two patients with conduction disturbances distal to the His bundle. Before the drug both patients had complete right bundle-branch block, left axis deviation, and an HV interval of 130 ms.

After drug administration complete block distal to the His bundle developed in one patient and a 2:1 distal block in the other. The latter is shown in Fig. 1.

VENTRICULAR TACHYCARDIA

Before drug administration ventricular tachycardia could be initiated by a single test stimulus in four out of five patients. In the fifth patient spontaneous reinitiation of tachycardia always occurred im-

mediately after pacing-induced termination of tachycardia. The RR interval during tachycardia before the drug varied from 280 to 470, mean 405 ± 87 ms. After lorainide administration the RR interval during ventricular tachycardia increased in all patients, with a mean increase of 71 ± 30 ms. The slowing in tachycardia rate was followed by spontaneous termination of tachycardia in four out of five patients (Fig. 2). In only one patient were we not able to reinitiate tachycardia during programmed stimulation after lorainide. There were no significant differences in mode of initiation and width of tachycardia zone in the other three patients where ventricular tachycardia could be induced before lorainide.

EFFECT OF LORCAINIDE IN PATIENTS WITH ACCESSORY PATHWAY

Eleven patients with Wolff-Parkinson-White syndrome had sinus rhythm and both anterograde and retrograde conduction over the accessory pathway. One patient had atrial fibrillation. Two patients had a left sided concealed bypass and sinus rhythm.

ANTEROGRADE CONDUCTION OVER ACCESSORY PATHWAY (Table 2)

In the 11 patients with sinus rhythm the effective refractory period of the accessory pathway was determined before and after drug administration. When possible we compared the effect on the effective refractory period of the accessory pathway from data obtained during stimulation of the atrial lead which was located closest to the atrial end of the accessory pathway.

If this was not possible (cases 14, 17, and 20) data

Fig. 1 Development of 2:1 block distal to His during lorainide injection. Leads III, V1, a high right atrial lead (HRA), and a His bundle lead were recorded simultaneously.

from stimulation by way of the right atrial wire were used. Before the drug the effective refractory period of the accessory pathway in the anterograde direction varied from <220 to 410 ms.

In eight out of the 11 patients drug administration resulted in a complete anterograde block of the accessory pathway (Fig. 3). In one patient the effective refractory period of the accessory pathway lengthened by 130 ms. In the remaining two patients no effect was seen. Only one of the five patients with an effective refractory period of the accessory pathway ≤ 270 ms had conduction over the extra connection after the drug.

Atrial fibrillation was induced in six out of the 11

Table 2 *Effect of lorcaïnide on anterograde ERP_{AP} and tachycardia zone in 13 patients with WPW syndrome or concealed accessory pathway*

Case no.	Site of stimulation	Basic cycle length (ms)	ERP _{AP}		Tachycardia zone	
			Before (ms)	After (ms)	Before (ms)	After (ms)
1	CS	500	270	>500	50	120
2	HRA	500	410	>500	60	20
3	HRA	600	320	>600	30	30
4	CS	600	290	420	50	100
5	HRA	470	260	>470	60	60
6	HRA	600	280	280	0	0
7	CS	600	260	250	0	10
8	HRA	600	<220	>600	0	80
9	HRA	500	310	>500	15	0
10	CS	500	400	>500	140	NM
11	CS	500	230	>660	10	NM
12*	CS	600	—	—	90	NM
13*	HRA	500	—	—	40	20

ERP, effective refractory period; AP, accessory pathway; CS, coronary sinus; HRA, high right atrium; NM, not measured.
*Patients with a concealed left sided accessory pathway.

Fig. 2 Upper panel: ventricular tachycardia before drug administration. Lower panel: lorcaïnide administration results in a slowing in the rate of tachycardia followed by its termination. Leads I, V₁, a high right atrial lead (RA), and a His bundle lead were recorded simultaneously.

patients with sinus rhythm by pacing the atrium at high rates. In three out of these six patients atrial fibrillation could again be initiated after drug administration. In one patient lorcaïnide was given during atrial fibrillation; this patient had permanent atrial fibrillation. In the four patients showing atrial fibrillation before and after lorcaïnide the shortest RR interval between QRS complexes with pre-excitation was 220 to 250 ms before the drug. After lorcaïnide administration two patients had exclusive conduction over the atrioventricular nodal His connection. In the other two patients conduction over the accessory pathway continued. In one patient the shortest RR interval during pre-excitation lengthened from 220 to 410 ms; in the other patient no difference was observed.

RETROGRADE CONDUCTION OVER ACCESSORY PATHWAY (Table 3)

Ventricular pacing was always performed at the apex of the right ventricle. The retrograde effective refractory period of the accessory pathway could be determined before and after the drug in 11 out of the 14 patients (the two patients with a concealed bypass are included). One patient had atrial fibrillation. In two patients the His-AV nodal pathway had a shorter refractory period than the accessory pathway.

In two patients complete block in retrograde conduction over the accessory pathway developed after the drug. The remaining nine patients showed only a minor increase in the effective refractory period of the accessory pathway.

Fig. 3 Development of anterograde block in the accessory pathway during lorainide injection in a patient with the Wolff-Parkinson-White syndrome. The patient received a total amount of 120 mg over a five minute period. Already after 30 seconds, when 12 mg was given, block occurred in the accessory pathway. Leads II, V1, V6, a high right atrial (HRA) lead, coronary sinus (CS) lead, and a His bundle lead were recorded simultaneously.

EFFECT ON CIRCUS MOVEMENT TACHYCARDIAS USING ACCESSORY PATHWAY

Atrial stimulation before and/or after drug administration was done in 13 patients (Table 2). In 10 of these patients we were able to initiate tachycardia before, and in eight patients after, drug administration. In these patients we carefully determined the width of the tachycardia zone. No significant change was seen after drug administration.

Ventricular stimulation before and/or after

lorainide was done in 13 patients (Table 3). In six out of these 13 patients we were able to initiate tachycardia before, and in four patients after, lorainide. No significant change in the width of the tachycardia zone occurred after drug administration.

EFFECT OF LORCAINIDE ON HEART RATE DURING TACHYCARDIA AND TERMINATION OF TACHYCARDIA

Before lorainide all patients showing circus movement tachycardias required at least one extra

Fig. 4 Termination of circus movement tachycardia during lorainide injection. On the left (panel A) before drug administration. On the right (panel B) a small amount of lorainide (10 mg) resulted in termination of tachycardia by creating retrograde block in the accessory pathway. Leads II, V1, V6, a high right atrial (HRA) lead, a His bundle lead, and a coronary sinus (CS) lead were recorded simultaneously.

stimulus from the atrium or ventricle to terminate the arrhythmia.

When lorcaïnide was given during circus movement tachycardias spontaneous termination of tachycardia by block in the accessory pathway in the VA direction occurred in four out of nine patients (Fig. 4). As shown in Table 4, lorcaïnide resulted in a significant slowing in rate of tachycardia, which was based upon an increase in the HV and VA interval.

Maximal effect on all electrophysiological indices was seen five minutes after the injection of the drug was stopped. The effect of lorcaïnide lasted for 45 to 60 minutes. No intra-arterial blood pressures were measured. Based upon the frequency of the sinus node and absence of complaints of the patients, we conclude that lorcaïnide does not have an important effect on blood pressure in the supine position.

No complications resulted from the catheterisation or the drug.

Discussion

Our data show that lorcaïnide has no significant effect on the electrophysiological properties of the atrioventricular node or the refractory period of the atrium or ventricle. A decrease was seen in conduction velocity in the His bundle, bundle-branches, and ventricular muscle. The width of the His bundle electrogram, the duration of the HV interval, and the width of the QRS complex during atrial and ventricular stimulation increased significantly.

This effect, as shown in our two patients with subnodal conduction disturbances, makes the drug

Table 4 Effect of lorcaïnide on RR, AH, HV, and VA interval during circus movement tachycardia in nine patients with WPW syndrome or concealed accessory pathway

Interval	Before (ms)		After (ms)		P
	Range	Mean $\pm$ SD	Range	Mean $\pm$ SD	
RR	235-390	289 $\pm$ 48	260-400	333 $\pm$ 48	< 0.01
AH	70-180	118 $\pm$ 36	65-170	110 $\pm$ 41	NS
HV	25- 55	41 $\pm$ 11	35- 60	47 $\pm$ 9	< 0.01
VA	80-190	131 $\pm$ 32	85-240	175 $\pm$ 50	< 0.01

dangerous in patients with impaired conduction in the His-Purkinje system.⁴⁻⁶

Lorcaïnide had a beneficial effect in patients with ventricular arrhythmias. Administration of the drug resulted in slowing followed by termination of tachycardia in four out of five patients. In three of these four patients, however, the tachycardia could still be reinitiated by premature ventricular stimuli.

As we have shown previously,⁷ antiarrhythmic drugs such as procainamide may not prevent reinitiation of tachycardia by timed stimuli and may even increase the width of the tachycardia zone in patients with ventricular tachycardia. The slowing of the tachycardia rate and the tendency to termination, however, suggests that lorcaïnide could be a clinically effective agent in these patients. Clinically this has been observed both after intravenous and chronic oral administration of the drug.⁸⁻⁹

The effect of lorcaïnide in patients with accessory atrioventricular connection who suffer from circus movement tachycardia or atrial fibrillation is interesting. In most patients there was complete block or distinct prolongation of the refractory period of the accessory pathway in the anterograde direction. In contrast to our previous observations with procainamide, amiodarone, quinidine, and ajmaline,¹⁰ lorcaïnide blocked or considerably lengthened the effective refractory period of the accessory pathway in patients with an effective refractory period of their accessory pathway of 270 ms or less. This has also been observed by others.⁵ This suggests that the drug might be useful in protecting patients with a short refractory period of their accessory pathway, who are at risk for ventricular fibrillation if atrial fibrillation supervenes. As observed with other drugs in patients with the Wolff-Parkinson-White syndrome, the effect of lorcaïnide on retrograde conduction over the accessory pathway was less obvious.¹¹⁻¹³

Nevertheless in two out of 10 patients there was complete retrograde block over the accessory pathway. In five patients there was slight pro-

Table 3 Effect of lorcaïnide on retrograde ERP_{AP} and tachycardia zone in 13 patients with WPW syndrome and concealed accessory pathway

Case no.	Basic cycle length (ms)	Retrograde ERP _{AP}		Tachycardia zone	
		Before (ms)	After (ms)	Before (ms)	After (ms)
1	500	280	300	0	0
2	500	< 240	< 240	0	0
3	600	330	350	10	0
4	600	310	> 600	5	0
5	500	< 260	< 250	60	80
6	500	280	360	0	0
7	500	< 220	< 240	0	10
8	500	260	290	30	20
9	500	< 240	> 470	0	0
10	420	< 210	< 210	5	30
11	475	< 270	NM	0	NM
12*	500	< 210	NM	70	NM
13*	430	220	230	0	0

ERP, effective refractory period; AP, accessory pathway; NM, not measured.

*Patients with a concealed accessory pathway.

longation of the effective refractory period of the accessory pathway in retrograde direction. The occurrence of spontaneous termination of circus movement tachycardia in four patients caused by retrograde block of the accessory pathway is important. Lorcinide also slowed the rate of tachycardia by increasing the HV and VA interval.

As reported after amiodarone administration,¹³ reinitiation of circus movement tachycardia by timed atrial or ventricular stimuli was still possible in most patients after lorcinide.

In view of our observations we conclude that the effect of chronic oral lorcinide administration should be assessed in patients with the Wolff-Parkinson-White syndrome and patients with ventricular tachycardia. The drug seems to be contraindicated in patients with sub-atrioventricular nodal conduction disturbances.

References

- 1 Carmeliet E, Janssen P, Marsboom R, Van Nueten J, Xhonneux R. Antiarrhythmic, electrophysiologic and hemodynamic effects of lorcinide. *Arch Int Pharmacodyn Ther* 1978; **231**: 104-30.
- 2 Wellens HJJ. Value and limitations of programmed electrical stimulation of the heart in the study and treatment of tachycardias. *Circulation* 1978; **57**: 845-53.
- 3 Wellens HJ, Durrer D. Patterns of ventriculo-atrial conduction in the Wolff-Parkinson-White syndrome. *Circulation* 1974; **49**: 22-31.
- 4 Manz M, Steinbeck G, Lüderitz B. Action of lorcinide (RT 15889) on sino-atrial node function and intracardial excitation conduction. *Herz Kreislauf* 1979; **11**: 192-7.
- 5 Kasper W, Meinertz T, Kersting F, Löllgen H, Lang K, Just H. Electrophysiological actions of lorcinide in patients with cardiac disease. *J Cardiovasc Pharmacol* 1979; **1**: 343-52.
- 6 Ng C, Gstöttner M, Gmeiner R. Intracardiac electrophysiological effects of lorcinide in man. *Eur J Clin Pharmacol* 1979; **15**: 241-7.
- 7 Wellens HJJ, Bär FWHM, Lie KI, Düren DR, Dohmen HJ. Effect of procainamide, propranolol, and verapamil on mechanism of recurrent ventricular tachycardia. *Am J Cardiol* 1977; **40**: 579-85.
- 8 Cocco G, Strozzi C. Initial clinical experience of lorcinide (Ro 13-1042), a new antiarrhythmic agent. *Eur J Clin Pharmacol* 1978; **14**: 105-9.
- 9 Kesteloot H, Stroobandt R. Clinical experience with lorcinide (R 15 889), a new anti-arrhythmic drug. *Arch Int Pharmacodyn Ther* 1977; **230**: 225-34.
- 10 Wellens HJ, Bär FW, Gorgels AP. Effect of drugs in WPW syndrome. Importance of initial length of effective refractory period of accessory pathway (abstract). *Am J Cardiol* 1978; **41**: 372.
- 11 Wellens HJJ, Durrer D. Effect of procaine amide, quinidine, and ajmaline in the Wolff-Parkinson-White syndrome. *Circulation* 1974; **50**: 114-20.
- 12 Wellens HJ. Effect of drugs in the Wolff-Parkinson-White syndrome. *Adv Cardiol* 1975; **14**: 233-40.
- 13 Wellens HJJ, Lie KI, Bär FW, et al. Effect of amiodarone in the Wolff-Parkinson-White syndrome. *Am J Cardiol* 1976; **38**: 189-94.

Requests for reprints to Dr Frits W Bär, Department of Cardiology, University of Limburg, Annadal Hospital, 6201 Bx Maastricht, The Netherlands.