Table 2 Details of patients with confirmed diaphragmatic palsy

Patient No	Age	Diagnosis	Operation	Result	Outcome
1	15 months	Fallot	Repair	L Palsy	—
2	3 years	Fallot, LMBT, WS	Repair	L Palsy	Plication
3	8 months	DS, AVSD	PAB	L Palsy	Plication
4	16 months	Fallot	Repair	L Palsy	Plication
5	22 months	Fallot, DA	Repair	L Palsy	—

L, left hemidiaphragm; LMBT, previous left modified Blalock-Taussig shunt; WS, previous Waterston shunt; DS, Down syndrome; AVSD, atrioventricular septal defect; PAB, pulmonary artery banding; DA, ductus arteriosus.

usually suspected until there is difficulty weaning a patient off artificial ventilation. We found that ultrasound examination at the bedside was as effective as fluoroscopy in the diagnosis of a paralysed hemidiaphragm. Bedside examination facilitates early diagnosis and is of particular benefit in infants, when plication is often necessary to achieve extubation. There were no cases of paralysed right hemidiaphragm in our group, but we would not expect the diagnosis to be difficult with ultrasound because it is generally easier to image the right hemidiaphragm than the left. Bilateral palsy can be missed on fluoroscopic examination unless careful attention is given to the phases of respiration, because the two sides move in concert—albeit paradoxically. This error is less likely with ultrasound assessment because each hemidiaphragm is imaged independently. Ultrasound imaging of the left hemidiaphragm can be difficult in bigger chil-

dren and adults, but diaphragmatic palsy is hardly ever the cause of major respiratory morbidity in this group and they rarely require plication.

We thank Dr Ian James, Dr Edward Sumner, and Dr Duncan Macrae of the Cardiac Intensive Care Unit for allowing us to study patients under their care.

1 Watanabe T, Trusler GA, Williams WG, Edmonds JF, Coles JG, Hosokawa Y. Phrenic nerve paralysis after paediatric cardiac surgery. Retrospective study of 125 cases. *J Thorac Cardiovasc Surg* 1987;94:383–8.
2 Affatato A, Villagra F, De Leon JP, et al. Phrenic nerve paralysis following paediatric cardiac surgery. Role of diaphragmatic plication. *J Cardiovasc Surg* 1988;29: 606–9.
3 Haber K, Asher WM, Freimanis AK. Echographic evaluation of diaphragmatic motion in intra-abdominal diseases. *Radiology* 1975;114:141–4.
4 Kangaroo H, Sukov R, Sample WF, Lipson M, Smith L. Ultrasonographic evaluation of juxtadiaphragmatic masses in children. *Radiology* 1977;125:785–7.
5 Haller JO, Schneider M, Kassner EG, Friedman AP, Waldroup LD. Sonographic evaluation of the chest in infants and children. *Am J Radiol* 1980;134:1019–27.
6 Aldrich TK, Herman JH, Rochester DF. Bilateral diaphragmatic paralysis in the new born infant. *J Pediatr* 1980; 97:988–90.
7 Harris RS, Giovannetti M, Kim BK. Normal ventilatory movement of the right hemidiaphragm studied by ultrasonography and pneumotachography. *Radiology* 1983; 146:141–4.
8 Laing IA, Teele RL, Stark AR. Diaphragmatic movement in newborn infants. *J Pediatr* 1988;112:638–43.
9 Diamant MJ, Boechat MI, Kangaroo H. Real-time sector ultrasound in the evaluation of suspected abnormalities of diaphragmatic motion. *J Clin Ultrasound* 1985;13:539–43.
10 Mok Q, Ross-Russell R, Shinebourne EA. Prospective study of the frequency of phrenic nerve paralysis after cardiac surgery in children [Abstract]. *Br Heart J* 1989;61:99.

VIEWS FROM THE PAST Ernest Henry Starling

Starling had already published his law of the heart (see page 4) before his military service during the first world war. Using cardiac catheterisation, John McMichael, Peter Sharpey-Schafer, and Sheila Howarth questioned the general applicability of this law in studies started during the second world war (Lewis collection).

DENNIS M KRIKLER