endocarditis was reported,⁴ more and more cases have been described where it was possible to preserve the tricuspid valve by careful removal of the vegetations.⁵⁻⁷ The use of autologous pericardium for tricuspid valve repair has been described before but was used only if two minor leaflets or one major leaflet were involved.⁸

Our case was complicated by the associated endocarditis of the mitral valve where it was not possible to preserve the valve because of coexisting degenerative changes. Apart from this, the tricuspid valve also had vegetations in all three cusps and removal of these with part of the valve cusps resulted in three significant defects. Repair of these defects with autologous pericardium enabled us to maintain good apposition of the valve cusps. This was further reinforced with the insertion of a Duran ring.

In a large series of cases of tricuspid valve surgery, there was no difference in mortality between valve repair and valve replacement.³ It is, however, our view that conservative surgery of the atrioventricular valve should be attempted when practical and possible. This case illustrates the feasibility of conserving the

tricuspid valve despite the presence of vegetations on all the leaflets.

The excellent postoperative valve function after this repair is very encouraging but more cases are needed to establish that this should be attempted every time this clinical situation occurs.

- 1 Chan P, Ogilby JD, Segal B. Tricuspid valve endocarditis. *Am Heart J* 1989;117:1140-6.
- 2 Buda AJ, Zatz RJ, Le Mire MS, Bach DS. Prognostic significance of vegetations detected by two dimensional echocardiography in infective endocarditis. *Am Heart J* 1986;112:1291-6.
- 3 McGrath LB, Gonzales-Lavin L, Bailey BM, Grunkemeier GL, Fernandez JF, Laub GW. Tricuspid valve operations in 530 patients. *J Thorac Cardiovasc Surg* 1990;99:124-33.
- 4 Arbulu A, Thoms MW, Wilson RF. Valvectomy without prosthetic replacement, a lifesaving operation for tricuspid pseudomonas endocarditis. *J Thorac Cardiovasc Surg* 1972;64:103-7.
- 5 Jagger JD, McGaughan BC, Pawsey CGK. Tricuspid valve endocarditis cured by excision of a single vegetation. *Am Heart J* 1986;112:626-7.
- 6 Chandraratna PA, Reagan RB, Imaizumi T, Langevin E, Elkins RC. Infective endocarditis cured by resection of a tricuspid valve vegetation. *Ann Intern Med* 1978;89:517-8.
- 7 Tanaka M, Abe T, Hosokawa S, Suengaga Y, Hikosaka H. Tricuspid valve candida endocarditis cured by valve-sparing debridement. *Ann Thorac Surg* 1989;48:857-8.
- 8 Yee ES, Ulyott DJ. Reparative approach for right sided endocarditis. *J Thorac Cardiovasc Surg* 1988;96:133-40.

Comment

There are good reasons for Anderson *et al* to be proud of their technical achievement in repairing all three cusps of a tricuspid valve damaged by endocarditis. None the less, the question remains as to whether for the next patient they encounter this provides a better prospect than replacing the valve. Their view "that conservative surgery of the atrioventricular valve(s) should be attempted when practical and possible" reflects the current tide of opinion. However, in this instance, the attempt at repairing the mitral valve was abandoned because the surgeons regarded the attempted repair as unsatisfactory. The success they achieved vindicates their intraoperative decisions. The case emphasises that the excellent results reported for mitral valve repair include a selection process at this stage, and if bad intraoperative results had been accepted by the pioneers of mitral valve conservation it is likely that conservative operations on the mitral valve would have fallen into disrepute. In the case of the mitral valve there is a growing body of evidence to suggest that repair gives superior short and long term results for the patient. The ideal policy is much harder to establish in tricuspid valve operations because they are so relatively infrequent.¹ McGrath *et al* in a series of tricuspid valve

operations in 530 cases were unable to demonstrate benefit for a conservative policy over the results achieved with valve replacement.² A recent report, also from Leeds, reports excellent long term results with a tissue valve tricuspid valve replacement in a series of 73 patients followed for, on average, nearly 10 years.³ The evidence suggests that, although Anderson, Scott, and Nair achieved an excellent result by conserving the valve, a strategy aimed at replacing the valve with a reliable prosthesis might have done just as well. On the evidence available, an elaborate attempt to conserve a tricuspid valve in a patient with endocarditis who has to have mitral valve replacement to save his life seems hard to justify.

TOM TREASURE

St George's Hospital,
Blackshaw Road,
London SW17 0QT

- 1 Surgery for tricuspid regurgitation [editorial]. *Lancet* 1988;ii:1061-2.
- 2 McGrath LB, Gonzales-Lavin L, Bailey BM, Grunkemeier GL, Fernandez JF, Laub GW. Tricuspid valve operations in 530 patients. *J Thorac Cardiovasc Surg* 1990;99:124-33.
- 3 Eng J, Ravichandran PS, Kay PH, Murday AJ. Long-term results of Ionescu-Shiley valve in tricuspid position. *Ann Thorac Surg* 1990;51:200-3.