LETTERS TO THE EDITOR

- The British Heart Journal welcomes letters commenting on papers that it has published within the past six months.
- All letters must be typed with double spacing and signed by all authors.
- No letter should be more than 600 words.
- In general, no letter should contain more than six references (also typed with double spacing).

Heart rate variability in left ventricular hypertrophy

SIR,—Mandawat *et al* in their paper (*Br Heart J* 1995;73:139-44) showed that heart rate variability was significantly reduced in patients with left ventricular hypertrophy secondary to hypertension or aortic valve disease. In addition they showed that heart rate variability was not affected by β blocker treatment. However, they made no comment on the effect of other antihypertensive agents such as thiazide diuretics and certain calcium antagonists such as nifedipine which can cause reflex sympathetic activation and hence reduce heart rate variability. Nearly 60% of their patients were taking diuretics. We have recently found in a study of Chinese patients with hypertension that thiazide treatment was associated with reduced heart rate variability assessed by non-spectral and spectral methods, compared with a 4 week control during which potassium replacement treatment alone was given (unpublished). Like Mandawat *et al* we found a significant reduction in the SD of all NN intervals over 24 (SDNN) and in addition root mean square of difference of successive RRs (rMSSD), proportion of adjacent RRs more than 50 ms different (pNN50), with reduced low frequency and high frequency spectral power.

Though left ventricular hypertrophy may well be independently associated with reduced heart rate variability, we were surprised that no comment is made about the possible effect of treatments, other than β blockers, on heart rate variability in hypertensive subjects. In view of the possibility of an increase in sudden death in hypertensive patients taking thiazide diuretics¹ we would be interested to know whether Mandawat *et al* have any comparable data on the relation of thiazide treatment to heart rate variability in hypertensive subjects with left ventricular hypertrophy.

JOHN E SANDERSON
BRIAN TOMLINSON

Department of Medicine and Clinical Pharmacology,
Chinese University of Hong Kong,
Prince of Wales Hospital,
Shatin, NT Hong Kong

1 Siscovick DS, Raghunathan TE, Psaty BM, *et al*. Diuretic therapy for hypertension and the risk of primary cardiac arrest. *N Engl J Med* 1994;330:1852-7.

This letter was shown to the authors, one of whom replies on behalf of his co-authors as follows:

SIR,—The observation by Sanderson and Tomlinson of reduced heart rate variability during treatment with a thiazide diuretic raises the interesting possibility of altered cardiac autonomic tone as a potential mechanism for the increased risk of sudden death associated with this treatment. Unfortunately this suggestion is neither supported nor refuted by our study. Most of our subjects were receiving a combination of antihypertensive drugs (mean 2.5). The inclusion of a diuretic with the regimen did not influence indices of heart rate variability (SD of all NN intervals over 24 h; triangular index; and SD of the mean NN interval for all 5 min segments of a 24 h recording), whether analysed as any diuretic drug ($n = 54$) or, more specifically, as a thiazide preparation ($n = 32$ of 82) (heart rate variability corrected for RR interval). Similarly, neither calcium antagonists ($n = 39$) nor angiotensin converting enzyme inhibitors ($n = 21$) affected time heart rate variability in these subjects with left ventricular hypertrophy. During prospective follow up of this cohort over 5.3 years, all-cause mortality was reduced by β blocker treatment but was unaffected by diuretic treatment (unpublished).

DAVID R WALLBRIDGE
Department of Cardiology,
Universitätsklinikum Essen,
Hufelandstrasse 55,
45122 Essen, Germany

Risks and results of surgery

SIR,—We congratulate Professor Tom Treasure on his interesting and forward looking article.¹ However, we suggest that the objectives in this instance should not be limited merely to "redressing apparent differences in results by correcting for differential casemix". Much more important potential benefits await the widespread introduction of an appropriate means of risk adjustment from observational databases.^{2,3} These include an ability in coronary artery surgery to produce predictive data comparable to randomised controlled trials.^{4,5}

Risk stratification on the basis of preoperative, perioperative, and postoperative variables is the cornerstone of any realistic comparisons but the system chosen must be robust and not open to manipulation or subjective elements.^{6,7} The best system available for the task must be used. A system described in 1990 has been endorsed by the Association of Cardiothoracic Anaesthetists of Great Britain and Ireland (ACTA).⁸ This ACTA system has been demonstrated to perform better (with improved receiver operating characteristic curves) than the Parsonnet system.⁹ In this comparison the explanatory components of the Parsonnet and ACTA scoring systems were subjected to multiple logistic regression (MLR) analysis. The MLR package always selected the ACTA items rather than the Parsonnet items. Moreover the subjective elements of the Parsonnet data set ("catastrophic state" and "PTCA crash") were not selected as explanators of increased mortality or prolonged stay in the intensive therapy unit (ITU) in this analysis.

There are other systems available, one of which (from New York State) has recently gained considerable publicity for the claims made for it and the criticism it has received.^{7,10} This system and the Parsonnet system both contain subjective elements, which in the Parsonnet system are not good explanators of outcome but can be used subjectively to increase the score a patient achieves.

The Parsonnet and New York State data sets confine their outcome analysis to death in hospital, but there are more valuable measures of the process undertaken in different centres. Length of ITU stay (included in the ACTA system), length of hospital stay, and estimated in hospital cost are also important outcomes that can be estimated before cardiac surgery and are valuable both to the patient and to the centre undertaking the surgery.

The data becoming available from this type of observational study are increasingly valuable in addressing the planning needs of healthcare purchasers worldwide. Conclusions drawn from such observational studies with casemix adjustment have been recognised to be as valuable as randomised controlled trials. Acceptance of a proven inferior system would considerably retard the laudable objectives of Professor Treasure and others.

S N C BOLSLIN
A M S BLACK
A J BRYAN
C J E DAY
Directorate of Anaesthesia,
Bristol Royal Infirmary,
Bristol BS2 8HW

- 1 Treasure T. Risks and results of surgery. *Br Heart J* 1995;74:11-2.
- 2 Bolsin SNC, Bryan AJ. Last word . . . Focus on Outcome Analysis 1995;1:19-20.
- 3 Hannan EL, Kilburn HJ, Racz M, Shields E, Chassin MR. Improving the outcomes of coronary artery bypass surgery in New York State. *J Am Med Ass* 1994;271:761-6.
- 4 Hlatky MA, Califf RM, Harrell FE, Lee KL, Mark DB, Pryor DB. Comparison of predictions based on observational data with the results of randomised controlled trials of coronary artery bypass surgery. *J Am Coll Cardiol* 1988;11:237-45.
- 5 Sowden AJ, Deeks JJ, Sheldon TA. Volume and outcome in coronary artery bypass graft surgery: true association or artifact? *BMJ* 1995;311:151-5.
- 6 Black AMS, Harris RI. Comparing two head injury treatments by linear logistic model. *Acta Neurochir Suppl (Wien)* 1982;62:31-46.
- 7 Green J, Wintfeld N. Report cards on cardiac surgeons. Assessing New York State's approach. *N Engl J Med* 1995;332:1229-32.
- 8 Bolsin SN, Morgan C, Ray R. A simple, pre-operative scoring system for predicting mortality and intensive care unit stay in adult patients undergoing heart surgery with cardiopulmonary bypass. *J Cardiothorac Vasc Anesth* 1990;4(suppl 1):139.
- 9 Day C, Duncan F, Lopatzidis A, Feneck R, Bolsin S, Black A. Pre-weighted sum scores and stepwise logistic regression in explaining mortality from cardiac surgery [abstr]. *Br J Anaesth* 1995;75:242P.
- 10 Hannan EL, Kumar D, Racz M, Sui AL, Chassin MR. New York State's Cardiac Surgery Reporting System: four years later. *Ann Thorac Surg* 1995;58:1852-7.

Percutaneous balloon dilatation of the mitral valve in critically ill young patients with intractable heart failure

SIR,—Patel and colleagues demonstrated the important role of emergency percutaneous

balloon dilatation of the mitral valve using the Inoue technique in young patients who are critically ill because of severe mitral valve stenosis.¹ We note that the procedure was performed while three of their patients were on assisted ventilation. Three patients had atrial fibrillation, but it is not clear whether this group overlapped with the patients who required ventilatory support.

In an earlier study percutaneous balloon dilatation of the mitral valve was associated with a 2–4% incidence of arterial emboli.² We believe that patients at high risk for embolic events should have transoesophageal echocardiographic (TOE) screening before valve dilatation because it has been shown that TOE is significantly superior to transthoracic imaging in detecting of important left atrial thrombus.^{2,3}

We have successfully carried out Inoue balloon mitral valve dilatation in two patients (aged 61 and 59 years). Both had atrial fibrillation with severe left ventricular failure secondary to critical mitral stenosis and could not be weaned from intravenous inotropes or ventilatory support. In both patients transoesophageal assessment before the procedure showed spontaneous contrast in the left atrium but no left atrial thrombus. The mitral valve echocardiographic scores (Wilkins *et al*⁴) were 8 and 10. A significant reduction was obtained in the transmitral end diastolic gradient (50% and 80% respectively) and there was a 2–3 fold increase in the mitral valve area on echocardiography. Both patients were successfully weaned off ventilatory support within 24 hours of valve dilatation.

We suggest that percutaneous dilatation of the mitral valve using the Inoue balloon technique is as valuable in older patients who cannot be weaned from assisted ventilation as a result of critical mitral stenosis as it is in younger patients. Older patients are more likely to be in atrial fibrillation and should have transoesophageal echocardiography.

G K DAVIS
D H ROBERTS
J S WRIGHT
Regional Cardiac Centre,
Blackpool Victoria Hospital,
Blackpool

- 1 Patel JJ, Munclinger MJ, Mitha AS, Patel N. Percutaneous balloon dilatation of the mitral valve in critically ill young patients with intractable heart failure. *Br Heart J* 1995; 73:555–8.
- 2 Thomas MR, Monaghan MJ, Smyth DW, Metcalfe JM, Jewitt DE. Comparative value of transthoracic and transoesophageal echocardiography before balloon dilatation of the mitral valve. *Br Heart J* 1992;68: 493–7.
- 3 Manning WJ, Reis GJ, Douglas PS. Use of transoesophageal echocardiography to detect left atrial thrombi before percutaneous balloon dilatation of the mitral valve: a prospective study. *Br Heart J* 1992;67: 170–3.
- 4 Wilkins GT, Weyman HF, Abascal VM, Block PC, Palicou IF. Percutaneous balloon dilatation of the mitral valve: an analysis of the echocardiographic variables related to outcome and the mechanism of dilatation. *Br Heart J* 1988;60:299–308.

This letter was shown to the authors, one of whom replies as follows:

SIR,—I appreciate the concern voiced by Davis and his colleagues regarding the role

of transoesophageal echocardiography to exclude left atrial thrombus before percutaneous balloon dilatation of the mitral valve. At our hospital we have stopped performing routine transoesophageal assessment because in our experience the yield, following on a carefully performed transthoracic cross sectional echocardiographic study, was low.¹ This may be because most of our patients are young (mean age 30 years) and in sinus rhythm. Only 16% of all our patients undergoing balloon dilatation of the mitral valve have transoesophageal echocardiography. I do not, however, underestimate its important role in elderly patients who are in atrial fibrillation.

J PATEL
Cardiac Unit,
Wentworth Hospital/University of Natal,
Durban, South Africa

- 1 Patel JJ, Dyer RB, Pillay R, Munclinger MJ, Mitha AS. Do all patients undergoing Inoue balloon mitral valvotomy need transoesophageal echo (TEE) to exclude left atrial thrombus? [Abstr]. *J Am Coll Cardiol* 1995; 89A.

Strut fracture of the convexo-concave Björk-Shiley mitral valve prosthesis

SIR,—Goodfield *et al* (*Br Heart J* 1995;73: 391) recommend an overpenetrated chest radiograph in posteroanterior and lateral or oblique views if strut fracture is suspected. This takes time when there is no time. Also, in a breathless patient with pulmonary oedema it would be difficult to acquire the high quality penetrated views required to show absence of the occluder.

Sudden total mitral regurgitation transforms a person from normal health to desperate straits virtually instantaneously. In a patient with this prosthetic valve, for practical purposes, there is no differential diagnosis.

Valve thrombosis has a more gradual onset and is less disastrous, with auscultatory signs of mixed stenosis and regurgitation. Loss of a mitral occluder produces pulmonary oedema without murmurs (no pressure gradient between ventricle and atrium). Prosthetic clicks are absent in both. Echocardiography is fast and fully informative but should only be done en route to the operating room.

Goodfield *et al* successfully treated their patient. The x rays which they show were taken with the chest open on the operating table.

CELIA M OAKLEY
Royal Postgraduate Medical School,
Hammersmith Hospital,
Du Cane Road,
London W12 0NN

This letter was shown to the authors, who reply as follows:

SIR,—Professor Oakley is of course correct in her comments about our dramatic image

in cardiology showing an intraoperative chest x ray with the disc occluder of a mitral Björk-Shiley convexo-concave valve lying within the aorta. The other x ray showed the fractured strut lying within the femoral artery. If a patient who is known to have a convexo-concave (CC) valve presents with sudden cardiovascular collapse and pulmonary oedema to a centre with cardiothoracic surgery, then of course the valve must immediately be replaced.

Many patients with implanted Björk-Shiley CC valves have not been identified and therefore do not carry information stating that they have this valve in place.¹ Most patients presenting to hospital with acute pulmonary oedema do not present to one that has cardiac surgery immediately available and with an experienced cardiologist to see that individual in that casualty department and make the diagnosis. Patients with prosthetic valves in situ who develop sudden pulmonary oedema should always of course be suspected of having valve dysfunction but this may also have other causes such as dehiscence secondary to endocarditis, or may be related to ventricular dysfunction, for example associated with arrhythmia. The absence of prosthetic clicks in a patient in extremis in a noisy accident and emergency department may be apparent only to very experienced observers. Cardiac surgeons would be unnecessarily overwhelmed if all such patients were transferred on presentation for immediate valve replacement.

There is a need for immediate and accurate diagnosis of outlet strut fracture and Hiratzka *et al* in a review of the subject have recommended “the single most expeditious study that should be readily obtainable in virtually any hospital or emergency treatment centre and that is highly effective in making the diagnosis of outlet strut fracture is an overpenetrated chest radiography, preferably in two views (posterior anterior and lateral or oblique).” We believe that this remains sound advice. Patients presenting with cardiovascular collapse caused by strut fracture of the valve may need immediate ventilation and a penetrated chest x ray may then permit the diagnosis. An intraoperative x ray can help identify the displaced components of the prosthesis and facilitate their removal, as was the case in the patient presented in our illustration.

Almost 100 000 patients are seen annually in the casualty department in our hospital and there are many cases of acute pulmonary oedema. The diagnosis may be obvious to an experienced cardiologist looking at a single illustration after the diagnosis has been made, but what is needed is straightforward advice for those dealing with a common medical emergency with an uncommon but catastrophic cause, and we believe that advice to obtain a penetrated radiograph in such cases remains sound.

P BLOOMFIELD
N E R GOODFIELD
Cardiology Department,
The Royal Infirmary of Edinburgh,
Lauriston Place,
Edinburgh EH3 9YW

- 1 Hiratzka LF, Kouchoukos NT, Grunkemeier GL, Miller C, Skully HE, Wechsler AS. Outlet strut fracture of Björk-Shiley convexo-concave valve: current information and recommendations for patient care. *J Am Coll Cardiol* 1988;11:1130–7.