- analysis of the courses of 43 patients with univentricular heart palliated by pulmonary artery banding. *Circulation* 1986;73:758-64.
- 15 Franklin RC, Sullivan ID, Anderson RH, Shinebourne EA, Deanfield JE. Is banding of the pulmonary trunk obsolete for infants with tricuspid atresia and double inlet ventricle with a discordant ventriculo-arterial connection? Role

- of aortic arch obstruction and subaortic stenosis. *J Am Coll Cardiol* 1990;16:1455-64.
- 16 Wessel DL, Adatia I, Giglia TM, Thompson JE, Kulik TJ. Use of inhaled nitric oxide and acetylcholine in the evaluation of pulmonary hypertension and endothelial function after cardiopulmonary bypass. *Circulation* 1993;88:2128-38.

SHORT CASES IN CARDIOLOGY

Large left ventricular thrombus in a patient with Friedreich's ataxia

K Reynen, D Claus, A Bornemann

Department of
Internal Medicine II,
University of
Erlangen-Nürnberg,
Erlangen, Germany
K Reynen

Department of
Neurology, University
of Erlangen-Nürnberg,
Erlangen, Germany
D Claus

Department of
Pathology, University
of Erlangen-Nürnberg,
Erlangen, Germany
A Bornemann

Correspondence to:
Dr K Reynen, Department
of Internal Medicine II,
University of Erlangen-
Nürnberg, Östliche
Stadtmauerstrasse 29,
D-91054 Erlangen,
Germany.

Accepted for publication
17 July 1995

Large apical thrombus
(arrow) in the left ventricle,
which was of normal size,
visualised by cross sectional
echocardiography (trans-
thoracic approach, apical
view). LV, left ventricle;
RV, right ventricle; LA,
left atrium; RA, right
atrium.

A 25 year old woman was admitted to the hospital because of fatigue, palpitation, and dyspnoea on minor exertion. Nineteen years before, Friedreich's ataxia had been diagnosed. On physical examination, the patient presented characteristic findings of Friedreich's ataxia, including marked ataxia, dysarthria, generalised weakness, severe scoliosis, and cavus deformity of the foot. No heart murmur was audible. The electrocardiogram showed atrial flutter with 2:1 atrioventricular conduction; the QRS rate was 170/min. Echocardiography revealed biventricular hypertrophy with left ventricular wall thickness of 13 mm, severe global hypokinesis of a left ventricle of normal size, and a large apical thrombus (3.5 × 2.5 cm) (figure).

About 20 hours after the echocardiographic examination, the patient suddenly lost consciousness; she was intubated because of respiratory insufficiency. Although pulses of the carotid arteries were strongly palpable, pulses at all four extremities were not. Thrombus embolisation was therefore suspected, and confirmed by echocardiography. The neuro-

logical disorder and the apparent cerebrovascular involvement ruled out vascular surgical intervention.

At necropsy residual thrombotic material was present in the apex of the left ventricle. Occlusive thrombotic material was found in the left internal carotid artery, the left subclavian and right brachial artery, and the aortic bifurcation.

Although Friedreich's ataxia is primarily an inherited progressive neurological disease, heart failure is the most common cause of death in these patients.¹ Concentric or, less frequently, asymmetrical left ventricular wall thickening may readily be detected by echocardiography: rarely, a picture of dilated cardiomyopathy is found.^{2,3} Echocardiography also allows assessment of systolic and diastolic ventricular function.³ Increasing physical immobility can mask myocardial insufficiency for a long time. Atrial arrhythmias raise suspicions of left ventricular dysfunction and indicate a poor prognosis.^{1,3} Occasional peripheral emboli resulting from mural thrombi or, more usually, from atrial thrombi in cases of atrial fibrillation may occur in Friedreich's ataxia. The thrombus was exceptionally large in this case and there was massive lethal embolisation. Anticoagulation should be considered in patients with Friedreich's ataxia and impaired ventricular wall motion or atrial fibrillation.

- 1 Hewer RL. Study of fatal cases of Friedreich's ataxia. *BMJ* 1968;iii:649-52.
- 2 Child JS, Perloff JK, Bach PM, Wolfe AD, Perlman S, Kark RAP. Cardiac involvement in Friedreich's ataxia: a clinical study of 75 patients. *J Am Coll Cardiol* 1986; 7:1370-8.
- 3 Alboliras ET, Shub C, Gomez MR, Edwards WD, Hagler DJ, Reeder GS, et al. Spectrum of cardiac involvement in Friedreich's ataxia: clinical, electrocardiographic and echocardiographic observations. *Am J Cardiol* 1986; 58:518-24.