CARDIOVASCULAR MEDICINE

Meta-analysis of randomised controlled trials of the effectiveness of antiarrhythmic agents at promoting sinus rhythm in patients with atrial fibrillation

G Nichol, F McAlister, B Pham, A Laupacis, B Shea, M Green, A Tang, G Wells

Heart 2002;87:535-543

Objective: To conduct a meta-analysis of randomised controlled trials to estimate the effectiveness of antiarrhythmic drugs at promoting sinus rhythm in patients with atrial fibrillation.

Design: Articles were identified by using a comprehensive search of English language papers indexed in Medline from 1966 to August 2001. For the outcomes of sinus rhythm and death, a random effects model was used to model repeated assessments within a study at different time points.

Setting: Emergency departments and ambulatory clinics.

Patients: Patients with atrial fibrillation.

Interventions: Antiarrhythmic agents grouped according to their Vaughan-Williams class.

Main outcome measures: Sinus rhythm and mortality.

Results: 91 articles met a priori criteria for inclusion in the analysis. Median duration of follow up was one day (range 0.04-1096, mean (SD) 46 (136) days). The median proportion of patients in sinus rhythm at follow up was 55% (range 0-100%) and 32% (range 0-90%) receiving active treatment and placebo, respectively. Median survival was 99% (range 55-100%) and 99% (range 55-100%). Compared with placebo, the following drug classes were associated with increased sinus rhythm at follow up: IA (treatment difference 21.5%, 95% confidence interval (CI) 16.3% to 26.8%); IC (treatment difference 33.1%, 95% CI 23.3% to 42.9%); and III (treatment difference 17.4%, 95% CI 11.5% to 23.3%). Class IC drugs were associated with increased sinus rhythm at follow up compared with class IV drugs (treatment difference 43.2%, 95% CI 11.5% to 75.0%). There was no significant difference in mortality between any drug classes.

Conclusions: Class IA, IC, and III drugs are associated with increased sinus rhythm at follow up compared with placebo. It is unclear whether any antiarrhythmic drug class is associated with increased or decreased mortality.

See end of article for authors' affiliations

Correspondence to:
Dr Graham Nichol, F6,
Ottawa Hospital, Civic
Site, 1053 Carling
Avenue, Ottawa, Ontario,
Canada K1Y 4E9;
grahamnichol@earthlink.net

Accepted
27 February 2002

Atrial fibrillation is the most common cardiac arrhythmia¹ and can be associated with disabling complications.^{1,2} At least 5% of admissions for cardiovascular disease are for management of atrial fibrillation.¹

Patients in atrial fibrillation can receive treatment to decrease embolic complications, to promote sinus rhythm, or to control ventricular rate. Each of these treatments is associated with particular risks and benefits. Although the effectiveness of anticoagulant treatment in atrial fibrillation is well established, use of electrical cardioversion or antiarrhythmic drugs may offer additional benefits. Compared with patients with atrial fibrillation, patients in whom sinus rhythm is restored have fewer strokes,² greater cardiac output,³ and reversal of cardiomyopathy.³⁻⁵ However, attempts to maintain sinus rhythm may be associated with increased mortality.⁶ Furthermore, as many as 75% of patients relapse into chronic atrial fibrillation within 12 months of restoration of sinus rhythm.⁷ Any reduction in the mortality or morbidity caused by stroke that is attained by use of antiarrhythmic drugs may be offset by adverse drug effects or relapse to atrial fibrillation.

Antiarrhythmic treatment to restore and maintain sinus rhythm after atrial fibrillation has been evaluated in many small trials. These trials evaluated several drugs, each of which has unique mechanisms of action and side effects. A large trial underway is evaluating the effectiveness of antiarrhythmic drugs in patients with atrial fibrillation. In the AFFIRM study (atrial fibrillation follow-up investigation of rhythm management), the choice of antiarrhythmic drug was at the discretion of the physician.⁸ Therefore, the results of this study are

unlikely to determine whether a specific antiarrhythmic drug has long term effectiveness in atrial fibrillation. Despite ongoing and previous trials, there is major interphysician variability in prescribing behaviours related to antiarrhythmic treatment in patients with atrial fibrillation.⁹

Meta-analysis is a method of pooling the results of individual trials statistically to provide an overall estimate of the effectiveness of a treatment. The combination of several studies can yield additional power to detect significant differences between treatments. Therefore, the purpose of this meta-analysis was to estimate the effectiveness of various classes of antiarrhythmic drugs in patients with atrial fibrillation. Using a protocol that was developed a priori, we conducted a meta-analysis based on conventional techniques.^{10,11} The protocol consisted of selection criteria for the primary studies, definitions of the primary end points, and an analysis plan.

METHODS

Identification of trials

Our aim was to identify all randomised controlled trials for which results were available by August 2001. A randomised controlled trial was defined as a research study in which patients were assigned prospectively to a treatment group by random allocation. Studies were identified by using a comprehensive search of English language articles indexed in Medline for the keywords "atrial fibrillation" and "antiarrhythmic agents". Potentially relevant randomised trials were identified by combining these keywords with the

Table 1 Effect of antiarrhythmic agents on sinus rhythm in placebo controlled trials

Comparators	Number of trials	Follow up* (days), median (range)	Proportion in sinus rhythm at follow up in treatment arm (%), mean (95% CI)	Proportion in sinus rhythm at follow up in control arm (%), mean (95% CI)	Treatment difference† (%), mean (95% CI)	p Value
IA v Placebo	7	93 (0.04–372)	54.7 (50.7 to 58.6)	33.1 (29.2 to 37.0)	21.5 (16.3 to 26.8)	<0.0001
IC v placebo	17	0.29 (0.04–186)	60.2 (52.6 to 67.8)	27.1 (19.6 to 34.6)	33.1 (23.3 to 42.9)	<0.0001
III v placebo	19	0.33 (0.04–1096)	46.5 (41.4 to 51.6)	29.1 (23.9 to 34.4)	17.4 (11.5 to 23.3)	0.001
Digoxin v placebo	4	0.58 (0.04–0.75)	48.4 (31.0 to 65.7)	35.3 (19.3 to 51.3)	13.1 (–7.2 to 33.3)	0.15
IA v III	6	1.0 (0.04–186)	44.3 (33.0 to 55.6)	40.1 (29.1 to 51.2)	4.1 (–8.0 to 16.2)	0.42
IC v III	10	0.91 (0.04–372)	56.7 (45.2 to 68.2)	60.1 (49.6 to 70.5)	–3.4 (17.2 to 10.5)	0.6
IC v IV	4	0.04 (0.04–2.0)	77.8 (42.7 to 100)	34.5 (0 to 77.2)	43.2 (11.5 to 75.0)	0.03
III v digoxin	2	1 (1–2)	82.9 (46.8 to 100)	67.7 (17.6 to 100)	15.2 (–43.7 to 74.2)	0.19

*Estimates from the control groups; †Percentage difference between treatment and control groups. CI, confidence interval.

Cochrane Collaboration's search strategy.¹² Only references published in the English language were considered. The bibliographies of published studies were reviewed to identify additional references. The authors of the primary studies were not contacted to identify additional studies. All published primary studies, but not unpublished studies or abstracts, were considered for inclusion in the analysis.

Classification of antiarrhythmic agents

All antiarrhythmic agents were grouped according to their mechanism of action by using the Vaughan-Williams classification system. Accordingly, drugs that were studied were classified as class I, which block sodium channels; class II, which block β receptors (for example, practolol); class III, which prolong repolarisation (for example, amiodarone, dofetilide, ibutilide, sotalol, and lidoflazine); class IV, which block calcium channels (for example, clonidine, diltiazem, magnesium, and verapamil); and other agents (for example, digoxin). Class I agents were subclassified as IA, which depress action potential upstroke, slow conduction, and prolong repolarisation (for example, cibenzoline, disopyramide, procainamide, and quinidine); and class IC, which depress upstroke depolarisation, slow conduction, and have a slight effect on repolarisation (for example, flecainide, pilsicainide, and propafenone.)

Inclusion and exclusion criteria

A priori exclusion criteria were studies that did not use a randomised study design; enrolled patients with induced atrial fibrillation; evaluated treatments other than antiarrhythmic agents; evaluated use of antiarrhythmic agents for prophylaxis of atrial fibrillation alone; followed up patients for < 60 minutes after drug administration; or lacked data on the proportion of patients in sinus rhythm at study follow up.

Study quality

The quality of each primary study was evaluated by using a validated assessment tool.¹³ This is a three item scale that measures the methodological quality of clinical studies. Individual items are randomisation, blinding, and withdrawal and dropout. Allocation concealment was also included (with two additional bonus points).¹⁴ Therefore, the maximum number of points per study was seven. Numbers were abstracted twice by two independent reviewers, and checked for accuracy after data entry.

Outcomes and data abstraction

The following variables were extracted from each study, if available: type and dose of intervention and control drugs; type and dose of concomitant drugs; number, age, and sex of patients randomly allocated; duration of follow up; number of patients who dropped out; number of patients who withdrew; number of patients in sinus rhythm at each follow up; and number of patients deceased. For the purpose of this analysis, patients were considered to have dropped out of the study if

they were lost to follow up. Patients were considered to have withdrawn if they stopped taking the study drug.

All studies were independently reviewed for eligibility, data abstraction, and study quality (by MG, FM, GN, BS, and AT). Differences were resolved by discussion until consensus was achieved. Data were abstracted twice and checked for accuracy after data entry.

Primary analyses

Data were analysed by using the SAS mixed procedure.¹⁵ Primary analyses compared the effect of each class of antiarrhythmic agent on the two outcome measures: the proportion of patients in sinus rhythm at the time of study follow up, and mortality.

For each outcome measure, a repeated measures random effects model was used to model the repeated assessments within a study at different time points.¹⁶ In the model, the outcome measure at each time point was weighted by the inverse variance of the outcome measure estimate. The model included treatment, assessment time, and a trial factor to force treatment comparison within the trial. Standard residual diagnostics were used to check for model goodness of fit. Least square means and their 95% confidence intervals (CI) for treatment difference were derived from the fitted models at the median follow up duration. Treatment difference was defined as the absolute difference in the proportion of patients who had experienced the outcome of interest (that is, sinus rhythm or mortality.) To avoid overfitting, effects were considered for classes of drugs rather than for individual drugs. Indicator variables were used for each class of drug, regardless of whether they were used singly or in combination. Two tailed $p = 0.05$ was considered significant.

Sensitivity analyses

For normal sinus rhythm at follow up, we performed the following sensitivity analyses. Firstly, we conducted subgroup analyses to determine the effectiveness of antiarrhythmic agents in trials that reported follow up of less than one week or at least one week. These analyses were conducted to identify whether any observed effectiveness was transient or sustained. They were also conducted to differentiate use of antiarrhythmic agents for conversion to sinus rhythm from use for maintenance of sinus rhythm. These analyses were done only for comparators where at least three trials were available since the model would not run with one or two trials.

Secondly, we included cohorts of patients who were taking one of the medications under comparison in the main model.¹⁶ These cohorts were extracted from other controlled trials that were included in the meta-analysis but that did not directly compare the medications under comparison. For example, if the main model compared class IA drugs with placebo, we checked the results of this model by including all the cohorts of patients who were taking either class IA drugs or placebo even though these additional cohorts were not part of a trial that directly compared class IA drugs with placebo.

Figure 1 Absolute treatment difference in proportion of patients with normal sinus rhythm (%) at end of study between class IA drugs and placebo.

Thirdly, we accounted for variation in trial quality by weighting an outcome on both its precision and its quality.¹⁷

For mortality data, we also performed three sensitivity analyses. Firstly, we included cohorts of patients as described above. Secondly, we assumed that no death was observed for trials that did not report mortality. Thirdly, we accounted for variation in trial quality.

Since amiodarone has several mechanisms of action, we considered the effect of amiodarone versus any other antiarrhythmic agent on sinus rhythm or mortality.

Lastly, we considered a large randomised controlled trial that included patients who had recently had atrial fibrillation but did not necessarily have it at the time of enrolment.¹⁸ Although this study considered a slightly different population from the other studies included in this analysis, it was considered because of its large sample size and relatively long duration of follow up.

RESULTS

Literature review

Four hundred twenty eight potentially relevant articles were identified. Reasons for exclusion were non-randomised study

design ($n = 128$); induced atrial fibrillation ($n = 3$); treatments other than antiarrhythmic agents ($n = 34$); use of antiarrhythmic agents for prophylaxis alone ($n = 56$); follow up for < 60 minutes after drug administration ($n = 8$); lack of data on the proportion of patients in sinus rhythm at study follow up ($n = 92$); and duplicate publication ($n = 16$). Ninety one trials met the criteria for inclusion in the analysis.¹⁹⁻¹⁰⁹

The appendix (on Heart website only) shows the data abstracted from these articles. There were 43, 29, 47, 1, 51, 21, and 12 trials for placebo, class IA, IC, II, III or IV drugs, and digoxin, respectively. Some trials evaluated more than two drugs or evaluated combinations of drugs. These articles described trials that were conducted in 24 countries and published between 1970 and 2001. The total number of enrolled patients was 8563 (range 18–506). The median duration of follow up was one day (range 0.04–1096, mean (SD) 46 (136) days.) The median proportion of patients in sinus rhythm at follow up was 55% (range 0–100%) and 32% (range 0–90%) for patients receiving active treatment and placebo, respectively. The median survival was 99% (range 55–100%) and 99%

Figure 2 Absolute treatment difference in proportion of patients with normal sinus rhythm (%) at end of study between class IC agents and placebo.

Figure 3 Absolute treatment difference in proportion of patients with normal sinus rhythm (%) at end of study between class III agents and placebo.

(range 55–100%) for patients receiving active treatment and placebo, respectively. The median quality score was 4 (range 0–7).

Effect of antiarrhythmic agents on promoting sinus rhythm at follow up

Compared with placebo, use of the following classes of drugs was associated with significant increases in the proportion of patients who were in sinus rhythm, adjusted for variation in follow up duration (table 1): IA (treatment difference 21.5%, 95% CI 16.3% to 26.8%, $p < 0.0001$; fig 1; IC (treatment difference 33.1%, 95% CI 23.3% to 42.9%, $p < 0.0001$; fig 2; III (treatment difference 17.4%, 95% CI 11.5% to 23.3%, $p = 0.001$; fig 3). Class IC was associated with significant increases in the proportion of patients who were in sinus rhythm at the end of study follow up compared with class IV (treatment difference 43.2%, 95% CI 11.5% to 75.0%, $p = 0.03$). For each figure, the combined treatment effect was estimated at the median follow up time.

Effect of antiarrhythmic agents on mortality

There was no significant difference in mortality between active treatment and placebo for any of the classes of antiarrhythmic drugs considered (table 2).

Adverse effects of antiarrhythmic agents

Although we attempted to abstract data describing any adverse effects that were observed in each trial, these were not reported frequently or consistently. Therefore, we were unable to analyse the association between individual classes of drugs and adverse effects.

Sensitivity analyses

Among trials that followed up patients for less than seven days, class IC and class III antiarrhythmic drugs were associated with increased sinus rhythm compared with placebo (table 3). Among the same group of trials, no class antiarrhythmic drugs were associated with a significant decrease in mortality.

Among trials that followed up patients for at least seven days, class IA and class III antiarrhythmic drugs were associated with increased sinus rhythm compared with placebo (table 3). Among the same group of trials, there were no significant differences in mortality between the classes of drugs compared.

Among trials that compared amiodarone with other antiarrhythmic drugs, there was no significant difference in sinus rhythm at follow up or in mortality (table 4). When a recent large randomised trial was included in the secondary analysis,¹⁸ the p value was closer to the critical value but was

Table 2 Effect of antiarrhythmic agents on mortality in placebo controlled trials

Comparators	Number of trials	Proportion dead at follow up in treatment arm (%), mean (95% CI)	Proportion dead at follow up in control arm (%), mean (95% CI)	Treatment difference (%), mean (95% CI)	p Value
IA v placebo	5	2.53 (1.41 to 3.64)	1.57 (0.58 to 2.56)	0.96 (-0.45 to 2.36)	0.14
IC v placebo	14	1.20 (1.10 to 1.30)	1.28 (1.17 to 1.39)	-0.08 (-0.20 to 0.04)	0.16
III v placebo	17	4.28 (4.06 to 4.50)	4.39 (4.16 to 4.62)	-0.11 (-0.32 to 0.09)	0.25
Digoxin v placebo	4	1.76 (1.71 to 1.80)	1.73 (1.69 to 1.78)	0.02 (-0.01 to 0.06)	0.12
IA v III	5	7.29 (6.12 to 8.45)	7.12 (6.01 to 8.23)	0.17 (-0.58 to 0.92)	0.53
IC v III	8	1.76 (1.07 to 2.45)	2.01 (1.28 to 2.74)	-0.26 (-1.10 to 0.59)	0.50
IC v IV	3	1.85 (0.53 to 3.17)	1.91 (0.23 to 3.59)	-0.06 (-2.26 to 2.13)	0.91
III v digoxin	2	3.29 (0 to 14.37)	2.82 (0 to 13.26)	0.48 (-13.71 to 14.67)	0.74

Table 3 Short term and long term effect of antiarrhythmic agents on sinus rhythm in placebo controlled trials

Comparators	Follow up duration (days)	Number of trials	Treatment difference (%), mean (95% CI),	p Value
IA v placebo	≥7 (all studies)	6	21.1 (15.0 to 27.2)	0.0003
IC v placebo	<7	17	31.9 (22.6 to 41.1)	<0.0001
III v placebo	<7	12	17.3 (9.6 to 25.0)	0.0002
III v placebo	≥7	3	17.6 (3.3 to 31.9)	0.03
Digoxin v placebo	<7 (all studies)	4	13.1 (-7.2, 33.3)	0.15
IA v III	<7	4	12.5 (-29.7 to 33.3)	0.11
IA v III	≥7	2	-6.6 (-36.7 to 23.5)	0.22
IC v III	<7	9	2.5 (-12.2 to 17.2)	0.71

Table 4 Effect of amiodarone versus other antiarrhythmic agents in placebo controlled trials

Outcome	Number of trials	Amiodarone estimate (%), mean (95% CI)	Other active drugs* estimate (%), mean (95% CI)	Treatment difference (%), mean (95% CI)	p Value
Sinus rhythm	20	50.0 (6.7 to 93.3)	47.2 (1.84 to 92.6)	2.7 (-51.2 to 56.7)	0.64
Mortality	20	2.12 (0.68 to 3.56)	2.29 (0.88 to 3.69)	-0.17 (-1.66 to 1.32)	0.38

*Other active drugs are digoxin, verapamil, propafenone, quinidine, procainamide, and flecainide.

still not significant (details available from authors). Similar results to those of the primary analysis were obtained when the analysis adjusted for differences in study quality or included study cohorts (details available from authors).

DISCUSSION

Compared with placebo, use of several of the classes of drugs considered in this analysis was associated with a significant increase in the proportion of patients who were in sinus rhythm at the time of follow up. However, there were limited data to show whether one class of drug was better than another. Compared with placebo, use of any of the classes of drugs considered in this analysis was not associated with a significant increase or decrease in mortality at the time of follow up. Compared with any other active antiarrhythmic agent, amiodarone was not associated with a significant difference in sinus rhythm at follow up or mortality.

The primary reason to use antiarrhythmic agents in patients with atrial fibrillation is to improve survival, quality of life, or both. Improvement in quality of life may be caused by decreased symptomatic arrhythmia or to decreased disabling complications such as stroke. No randomised trials have had sufficient power to address these issues. Whether sinus rhythm is present several months after antiarrhythmic treatment is initiated is at best a surrogate marker for either survival or quality of life. Meta-analysis of previous randomised controlled trials provides the best available information to address this issue but does not provide sufficient information on which to base clinical recommendations confidently. However, it appears that there is little difference between classes of antiarrhythmic agents in terms of achievement and maintenance of sinus rhythm. Furthermore, there is no evidence that any class of antiarrhythmic agents is associated with increased or decreased mortality. This lack of effect is important since previous trials of antiarrhythmic agents for other arrhythmias have shown that these agents are associated with increased mortality.

Based on this analysis, if a patient or physician chooses to use an antiarrhythmic agent as a treatment for atrial fibrillation, use of class III drugs as first line agents for treatment of atrial fibrillation is reasonable since they are both effective and safe. The present analysis did not find a significant difference in effectiveness between amiodarone and other class III drugs. However, previous trials showed increased mortality after myocardial infarction in patients who received D-sotalol¹¹⁰ or

proarrhythmia in patients with ventricular dysfunction who received dofetilide.¹¹¹ Therefore, caution is required when using class III agents other than amiodarone to promote sinus rhythm in patients with pre-existing heart disease.

The use of class IA or IC drugs may be associated with increased mortality.^{7 112 113} Therefore, these drugs are now not usually used in patients known to have coronary artery disease. The use of class IV drugs is restricted to situations where the primary goal is control of ventricular rate rather than restoration of sinus rhythm.

A previous meta-analysis of placebo controlled trials in patients with atrial fibrillation showed that quinidine treatment was more effective than no antiarrhythmic treatment but was associated with increased mortality.⁷ These findings are consistent with our finding that use of class IA antiarrhythmics is associated with increased sinus rhythm. It is also consistent with our finding of no difference in mortality, since the confidence intervals for the effect estimate from the quinidine analysis and that of the present analysis both overlapped zero. However, the previous analysis considered only the effect of quinidine rather than all class IA drugs.

Controlled trials of class III agents such as amiodarone for ventricular arrhythmias showed small but significant reductions in all cause mortality or arrhythmic death in patients who received treatment rather than the control treatment,¹¹⁴⁻¹¹⁷ in contrast to controlled trials of class IC agents.¹¹²⁻¹¹³ However, it would be premature to generalise the results of these trials to all patients with atrial fibrillation.

Others have considered the cost effectiveness of treatments for patients with atrial fibrillation.^{118 119} A decision analytical model was used to show that cardioversion followed by aspirin with or without amiodarone was effective and inexpensive.¹¹⁸ Cardioversion followed by treatment with warfarin and amiodarone was effective but more expensive. The present analysis supports the findings that amiodarone is effective and appropriate in patients with atrial fibrillation.

Another decision analysis had slightly different findings.¹¹⁹ Of the treatment strategies that were considered, cardioversion followed by repeat cardioversion plus amiodarone on relapse was most cost effective. Our findings support the use of initial treatment with amiodarone rather than delayed treatment, since class III agents are associated with increased sinus rhythm and may be associated with decreased mortality. However, we have not incorporated our results into a decision model since we were unable to show a significant difference in mortality between any class of antiarrhythmic drugs.

A large randomised controlled trial underway is evaluating the use of antiarrhythmic agents by patients with atrial fibrillation.⁶ The results of the AFFIRM trial will be reported in 2002. It evaluated the effectiveness of either ventricular rate control or rhythm control with antiarrhythmic agents for total mortality and stroke. All patients received concurrent anticoagulation treatment. Since selection of the type or class of drug was not randomised in this trial, any observed difference between types of antiarrhythmic agents may be confounded by differences in other factors.

Since the present analysis is complex, we will now review the rationale for the inclusion criteria we used and discuss their implications. Firstly, we excluded studies not published in English. Compared with meta-analyses that considered only English articles, those that considered English and non-English articles did not differ with respect to the estimate of benefit of the effectiveness of an intervention.¹²⁰ Also, use of such articles would require translation, which was outside of the scope of this study because of the anticipated difficulty in abstracting information about duration of follow up, and rates of dropout and withdrawal. Thus, we relied on citations that were identified by review of English language papers indexed in Medline and of the bibliographies of these articles.

Studies that considered patients with induced atrial fibrillation were excluded since the efficacy of antiarrhythmic agents in patients in an electrophysiology laboratory may not be generalisable to their effectiveness in a less selected population. Treatments other than antiarrhythmic agents, such as ablation, are not widely available. Therefore, the results of studies of these interventions are not generalisable to the population as a whole.

Studies that considered the use of antiarrhythmic agents for prophylaxis alone were also excluded. Since the frequency of the arrhythmia in these patients is unclear, it is difficult to interpret the presence of sinus rhythm at the time of follow up.

Finally, studies that followed up patients for < 60 minutes after drug administration were excluded, since transient conversion may be unsustainable. Such transient benefits are unlikely to affect patient morbidity or quality of life.

The effectiveness of each class of drug was expressed as the percentage difference in this analysis. A common alternative approach is to express the effectiveness of each class as an odds ratio. However, odds ratios may be misleading when events are common or when they are observed in a randomised trial rather than a case-control study.¹²¹⁻¹²³ Both conditions apply to this analysis. Also, odds ratios do not convey the absolute magnitude of the benefit of treatment.

Limitations of analysis

This analysis has several limitations because of the constraints of the existing literature. Firstly, there are several ways of classifying the underlying pattern of atrial fibrillation.¹²⁴ In our analysis, it was sometimes difficult to determine whether patients who were enrolled in a particular clinical trial had isolated, paroxysmal, persistent, or chronic atrial fibrillation. Despite this, we found consistent results when the data were pooled and analysed in different ways. Thus, we believe that it is unlikely that considering alternative ways of classifying atrial fibrillation would yield different results.

Secondly, the analysis lacked statistical power to detect differences between individual antiarrhythmic agents or to detect small differences between classes of antiarrhythmic agents. Nevertheless, this is the first meta-analysis to discriminate between each class of antiarrhythmic agent and is the largest analysis performed to date of the use of these agents in patients with atrial fibrillation.

Thirdly, the analysis adjusted for variation in follow up duration but not for other parameters of potential interest

including use of electrical cardioversion before or after initiation of antiarrhythmic agents, use of transoesophageal echocardiography to identify clot before cardioversion, or clinical characteristics of the patients. Although we attempted to extract such data, the vast majority of the studies analysed did not report them. Therefore, we were unable to evaluate the influence of these variables on sinus rhythm, survival, or other clinical outcomes.

Despite the limitations of the data described above, we believe that this analysis is the most comprehensive assessment to date of the effectiveness of different antiarrhythmic agents in patients with atrial fibrillation. It offers estimates of the effectiveness of different classes of drugs. No other studies have pooled the results of different studies using statistical analysis and adjusted for differences in follow up time. Given the large number of antiarrhythmic agents and large uncertainty about which agent is effective, this meta-analysis is an important advance. Future studies should directly compare specific class III agents and be of sufficient size to determine effectiveness and safety.

In summary, evaluation of the effectiveness of different classes of antiarrhythmic agents continues to be difficult because of inadequate data. The effectiveness estimates derived in this analysis must be interpreted with caution because of the wide variety of classes of drugs considered and of differences in the duration of follow up.

Conclusion

On the basis of a meta-analysis of data from 91 randomised trials of antiarrhythmic agents promoting sinus rhythm in patients with atrial fibrillation, we confirm that class IA, IC, and III drugs are associated with increased sinus rhythm compared with placebo. Class IC drugs are associated with increased sinus rhythm compared with class IV drugs. We are unable to confirm whether any antiarrhythmic class of drug is associated with increased or decreased mortality in this population. Patients, physicians, and policy makers should consider the impact of each class of drug on sinus rhythm, mortality, quality of life, and patient compliance.

ACKNOWLEDGEMENTS

Funding for this project was provided in part by the Bickell Foundation of Canada. Dr Nichol is supported by an Ontario Ministry of Health Career Scientist Award. Dr McAlister is supported by a Population Health Investigatorship from the Alberta Heritage Foundation for Medical Research. Dr Laupacis is a Senior Scientist of the Canadian Institute of Health Research.

Summary of studies examined in the meta-analysis appear on the Heart website—www.heartjnl.com

Authors' affiliations

G Nichol, Clinical Epidemiology Program, Ottawa Health Research Institute, Ottawa Hospital, Ottawa, Ontario, Canada
F McAlister, Division of General Internal Medicine, Department of Medicine, University of Alberta, Edmonton, Alberta, Canada
B Pham, Glaxo Wellcome Inc, Mississauga, Ontario, Canada
A Laupacis, Institute of Clinical Evaluative Sciences in Ontario, Toronto, Ontario, Canada
B Shea, Institute of Population Health, Ottawa, Ontario, Canada
M Green, A Tang, University of Ottawa Heart Institute, Ottawa, Ontario, Canada
G Wells, Department of Epidemiology and Community Medicine, University of Ottawa, Ottawa, Ontario, Canada

REFERENCES

- 1 Bailey D, Lehmann MH, Schumacher DN, *et al*. Hospitalization for arrhythmias in the United States: importance of atrial fibrillation. *J Am Coll Cardiol* 1992;19:41A.
- 2 Stroke Prevention in Atrial Fibrillation Investigators. Predictors of thromboembolism in atrial fibrillation. I. Clinical features of patients at risk. *Ann Intern Med* 1992;116:1-5.
- 3 Kieny JR, Sacrez A, Facello A, *et al*. Increase in radionuclide left ventricular ejection fraction after cardioversion of chronic atrial fibrillation in idiopathic dilated cardiomyopathy. *Eur Heart J* 1992;13:1290-5.
- 4 Kieny JR, Sacrez A, Facello A, *et al*. The causes of dilated cardiomyopathy: a clinical pathological review of 673 consecutive patients. *J Am Coll Cardiol* 1994;23:586-90.
- 5 Sarembock IJ, Horak AR, Commerford PJ. Tachycardia-induced left ventricular dysfunction: a report of two cases. *S Afr Med J* 1988;73:484-5.
- 6 Flaker GC, Blackshear JL, McBride R, *et al*. Antiarrhythmic drug therapy and cardiac mortality in atrial fibrillation. The Stroke Prevention in Atrial Fibrillation Investigators. *J Am Coll Cardiol* 1992;20:527-32.
- 7 Coplen SE, Antman EM, Berlin JA, *et al*. Efficacy and safety of quinidine therapy for maintenance of sinus rhythm after cardioversion: a meta-analysis of randomized controlled trials. *Circulation* 1990;82:1106-16.
- 8 Planning and Steering Committees of the AFFIRM study for the NHLBI AFFIRM investigators. Atrial fibrillation follow-up investigation of rhythm management: the AFFIRM study design. *Am J Cardiol* 1997;79:1198-202.
- 9 McAlister FA, Ackman M, Tsuyuki R, *et al*, for the CQIN Investigators. Contemporary utilization for digoxin in patients with atrial fibrillation. *Ann Pharmacother* 1999;33:289-93.
- 10 L'Abbe KA, Detsky AS, O'Rourke K. Metaanalysis in clinical research. *Ann Intern Med* 1987;107:224-33.
- 11 Sacks HS, Berrier J, Reitman D, *et al*. Meta-analyses of randomized controlled trials. *N Engl J Med* 1987;316:450-5.
- 12 Dickersin K, Scherer R, Lefebvre C. Identifying relevant studies for systematic reviews. *BMJ* 1994;309:1286-91.
- 13 Jadad AR, Moore RA, Carroll D, *et al*. Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Control Clin Trials* 1996;17:1-12.
- 14 Schulz KF, Chalmers I, Hayes RJ, *et al*. Empirical evidence of bias: dimensions of methodological quality associated with estimates of treatment effects in controlled trials. *JAMA* 1995;273:408-12.
- 15 Normand SL. Meta-analysis: formulating, evaluating, combining, and reporting. *Stat Med* 1999;18:321-59.
- 16 Dear KBG. Iterative generalized least squares for meta-analysis of survival data at multiple times. *Biometrics* 1994;50:989-1002.
- 17 Moher D, Pham B, Jones A, *et al*. Does quality of reports of randomised trials affect estimates of intervention efficacy reported in meta-analyses? *Lancet* 1998;352:609-13.
- 18 Roy D, Talajic M, Dorian P, *et al*. Amiodarone to prevent recurrence of atrial fibrillation. Canadian Trial of Atrial Fibrillation Investigators. *N Engl J Med* 2000;342:913-20.
- 19 Byrne-Quinn E, Wing AJ. Maintenance of sinus rhythm after DC reversion of atrial fibrillation: a double-blind controlled trial of long-acting quinidine bisulphate. *Br Heart J* 1970;32:370-6.
- 20 Levi G, Proto C, Rovetta A. Double-blind evaluation of practolol and quinidine in the treatment of chronic atrial fibrillation. *Cardiology* 1973;58:364-8.
- 21 Hartel G, Louhija A, Kontinen A. Disopyramide in the prevention of recurrence of atrial fibrillation after electroconversion. *Clin Pharmacol Ther* 1974;15:551-5.
- 22 Sodermark T, Jonsson B, Olsson A, *et al*. Effect of quinidine on maintaining sinus rhythm after conversion of atrial fibrillation or flutter: a multicentre study from Stockholm. *Br Heart J* 1975;37:486-92.
- 23 Normand JP, Legendre M, Kahn JC, *et al*. Comparative efficacy of short-acting and long-acting quinidine for maintenance of sinus rhythm after electrical conversion of atrial fibrillation. *Br Heart J* 1976;38:381-7.
- 24 Kennelly BM. Comparison of lidoflazine and quinidine in prophylactic treatment of arrhythmias. *Br Heart J* 1977;39:540-6.
- 25 Boissel JP, Wolf E, Gillet J, *et al*. Controlled trial of a long-acting quinidine for maintenance of sinus rhythm after conversion of sustained atrial fibrillation. *Eur Heart J* 1981;2:49-55.
- 26 Rasmussen K, Wang H, Fausa D. Comparative efficiency of quinidine and verapamil in the maintenance of sinus rhythm after DC conversion of atrial fibrillation: a controlled clinical trial. *Acta Med Scand Suppl* 1981;645:23-8.
- 27 Lloyd EA, Gersh BJ, Forman R. The efficacy of quinidine and disopyramide in the maintenance of sinus rhythm after electroconversion from atrial fibrillation: a double-blind study comparing quinidine, disopyramide and placebo. *S Afr Med J* 1984;65:367-9.
- 28 Campbell TJ, Gavanagh TP, Morgan JJ. Intravenous sotalol for the treatment of atrial fibrillation and flutter after cardiopulmonary bypass: comparison with disopyramide and digoxin in a randomised trial. *Br Heart J* 1985;54:86-90.
- 29 Ochs HR, Anda L, Eichelbaum M, *et al*. Diltiazem, verapamil, and quinidine in patients with chronic atrial fibrillation. *J Clin Pharmacol* 1985;25:204-9.
- 30 Cowan JC, Gardiner P, Reid DS, *et al*. A comparison of amiodarone and digoxin in the treatment of atrial fibrillation complicating suspected acute myocardial infarction. *J Cardiovasc Pharmacol* 1986;8:252-6.
- 31 Falk RH, Knowlton AA, Bernard SA, *et al*. Digoxin for converting recent-onset atrial fibrillation to sinus rhythm: a randomized, double-blinded trial. *Ann Intern Med* 1987;106:503-6.
- 32 Karlsson BW, Torstensson I, Abjorn C, *et al*. Disopyramide in the maintenance of sinus rhythm after electroconversion of atrial fibrillation: a placebo-controlled one-year follow-up study. *Eur Heart J* 1988;9:284-90.
- 33 Suttrop MJ, Kingma JH, Lie AHL, *et al*. Intravenous flecainide versus verapamil for acute conversion of paroxysmal atrial fibrillation or flutter to sinus rhythm. *Am J Cardiol* 1989;63:693-6.
- 34 Van Gelder IC, Crijns HJ, Van Gilst WH, *et al*. Efficacy and safety of flecainide acetate in the maintenance of sinus rhythm after electrical cardioversion of chronic atrial fibrillation or atrial flutter. *Am J Cardiol* 1989;64:1317-21.
- 35 Bertini G, Conti A, Fradella G, *et al*. Propafenone versus amiodarone in field treatment of primary atrial tachydysrhythmias. *J Emerg Med* 1990;8:15-20.
- 36 Juul-Moller S, Edvardsson N, Rehnqvist-Ahlberg N. Sotalol versus quinidine for the maintenance of sinus rhythm after direct current conversion of atrial fibrillation. *Circulation* 1990;82:1932-9.
- 37 Kondili A, Kastrati A, Papa Y. Comparative evaluation of verapamil, flecainide and propafenone for the acute conversion of atrial fibrillation to sinus rhythm. *Wien Klin Wochenschr* 1990;102:510-3.
- 38 McAlister HF, Luke RA, Whitlock RM, *et al*. Intravenous amiodarone bolus versus oral quinidine for atrial flutter and fibrillation after cardiac operations. *J Thorac Cardiovasc Surg* 1990;99:911-8.
- 39 Noc M, Stajer D, Horvat M. Intravenous amiodarone versus verapamil for acute conversion of paroxysmal atrial fibrillation to sinus rhythm. *Am J Cardiol* 1990;65:679-80.
- 40 Suttrop MJ, Kingma JH, Jessurun ER, *et al*. The value of class IC antiarrhythmic drugs for acute conversion of paroxysmal atrial fibrillation or flutter to sinus rhythm. *J Am Coll Cardiol* 1990;16:1722-7.
- 41 Donovan KD, Dobb GJ, Coombs LJ, *et al*. Reversion of recent-onset atrial fibrillation to sinus rhythm by intravenous flecainide. *Am J Cardiol* 1991;67:137-41.
- 42 Kuhlkamp V, Schmid F, Risler T, *et al*. Randomized comparison of flecainide and cibenzoline in the conversion of atrial fibrillation. *Int J Cardiol* 1991;31:65-9.
- 43 Singh S, Saini RK, DiMarco J, *et al*. Efficacy and safety of sotalol in digitalized patients with chronic atrial fibrillation. The Sotalol Study Group. *Am J Cardiol* 1991;68:1227-30.
- 44 Capucci A, Lenzi T, Boriani G, *et al*. Effectiveness of loading oral flecainide for converting recent-onset atrial fibrillation to sinus rhythm in patients without organic heart disease or with only systemic hypertension. *Am J Cardiol* 1992;70:69-72.
- 45 Hjelms E. Procainamide conversion of acute atrial fibrillation after open-heart surgery compared with digoxin treatment. *Scand J Thorac Cardiovasc Surg* 1992;26:193-6.
- 46 Kingma JH, Suttrop MJ. Acute pharmacologic conversion of atrial fibrillation and flutter: the role of flecainide, propafenone, and verapamil. *Am J Cardiol* 1992;70:56A-60A, discussion 60A-1A.
- 47 Roth A, Kaluski E, Felner S, *et al*. Clonidine for patients with rapid atrial fibrillation. *Ann Intern Med* 1992;116:388-90.
- 48 Zehender M, Hohnloser S, Muller B, *et al*. Effects of amiodarone versus quinidine and verapamil in patients with chronic atrial fibrillation: results of a comparative study and a 2-year follow-up. *J Am Coll Cardiol* 1992;19:1054-9.
- 49 Chapman MJ, Moran JL, O'Fathartaigh MS, *et al*. Management of atrial tachyarrhythmias in the critically ill: a comparison of intravenous procainamide and amiodarone. *Intensive Care Med* 1993;19:48-52.
- 50 Gullesstad L, Birkeland K, Molstad P, *et al*. The effect of magnesium versus verapamil on supraventricular arrhythmias. *Clin Cardiol* 1993;16:429-34.
- 51 Madrid AH, Moro C, Marin-Huerta E, *et al*. Comparison of flecainide and procainamide in cardioversion of atrial fibrillation. *Eur Heart J* 1993;14:1127-31.
- 52 Reimold SC, Cantillon CO, Friedman PL, *et al*. Propafenone versus sotalol for suppression of recurrent symptomatic atrial fibrillation. *Am J Cardiol* 1993;71:558-63.
- 53 Capucci A, Boriani G, Rubino I, *et al*. A controlled study on oral propafenone versus digoxin plus quinidine in converting recent onset atrial fibrillation to sinus rhythm. *Int J Cardiol* 1994;43:305-13.
- 54 Cochrane AD, Siddins M, Rosenfeldt FL, *et al*. A comparison of amiodarone and digoxin for treatment of supraventricular arrhythmias after cardiac surgery. *Eur J Cardiothorac Surg* 1994;8:194-8.
- 55 Weiner P, Ganam R, Ganem R, *et al*. Clinical course of recent-onset atrial fibrillation treated with oral propafenone. *Chest* 1994;105:1013-6.
- 56 Bellandi F, Cantini F, Pedone T, *et al*. Effectiveness of intravenous propafenone for conversion of recent-onset atrial fibrillation: a placebo-controlled study. *Clin Cardiol* 1995;18:631-4.
- 57 Boriani G, Capucci A, Lenzi T, *et al*. Propafenone for conversion of recent-onset atrial fibrillation: a controlled comparison between oral loading dose and intravenous administration. *Chest* 1995;108:355-8.
- 58 Boudonas G, Lefkos N, Efthymiadis AP, *et al*. Intravenous administration of diltiazem in the treatment of supraventricular tachyarrhythmias. *Acta Cardiol* 1995;50:125-34.
- 59 Di Biasi P, Scrofani R, Pajce A, *et al*. Intravenous amiodarone vs propafenone for atrial fibrillation and flutter after cardiac operation. *Eur J Cardiothorac Surg* 1995;9:587-91.

- 60 **Donovan KD**, Power BM, Hockings BE, *et al*. Intravenous flecainide versus amiodarone for recent-onset atrial fibrillation. *Am J Cardiol* 1995;**75**:693-7.
- 61 **Halinen MO**, Huttunen M, Paakkinen S, *et al*. Comparison of sotalol with digoxin-quinidine for conversion of acute atrial fibrillation to sinus rhythm (the sotalol-digoxin-quinidine trial). *Am J Cardiol* 1995;**76**:495-8.
- 62 **Hohnloser SH**, van de Loo A, Baedeker F. Efficacy and proarrhythmic hazards of pharmacologic cardioversion of atrial fibrillation: prospective comparison of sotalol versus quinidine. *J Am Coll Cardiol* 1995;**26**:852-8.
- 63 **Hou ZY**, Chang MS, Chen CY, *et al*. Acute treatment of recent-onset atrial fibrillation and flutter with a tailored dosing regimen of intravenous amiodarone: a randomized, digoxin-controlled study. *Eur Heart J* 1995;**16**:521-8.
- 64 **Atarashi H**, Inoue H, Hiejima K, *et al*. Conversion of recent-onset atrial fibrillation by a single oral dose of pilsicainide (pilsicainide suppression trial on atrial fibrillation). The PSTAF Investigators. *Am J Cardiol* 1996;**78**:694-7.
- 65 **Bianconi L**, Mennuni M, Lukic V, *et al*. Effects of oral propafenone administration before electrical cardioversion of chronic atrial fibrillation: a placebo-controlled study. *J Am Coll Cardiol* 1996;**28**:700-6.
- 66 **Botto GL**, Bonini W, Broffoni T, *et al*. Conversion of recent onset atrial fibrillation with single loading oral dose of propafenone: is in-hospital admission absolutely necessary? *Pacing Clin Electrophysiol* 1996;**19**:1939-43.
- 67 **Crijns HJ**, Gosselink AT, Lie KI. Propafenone versus disopyramide for maintenance of sinus rhythm after electrical cardioversion of chronic atrial fibrillation: a randomized, double-blind study. PRODIS Study Group. *Cardiovasc Drugs Ther* 1996;**10**:145-52.
- 68 **Ellenbogen KA**, Clemo HF, Stambler BS, *et al*. Efficacy of ibutilide for termination of atrial fibrillation and flutter. *Am J Cardiol* 1996;**78**:42-5.
- 69 **Fresco C**, Proclemer A, Pavan A, *et al*. Intravenous propafenone in paroxysmal atrial fibrillation: a randomized, placebo-controlled, double-blind, multicenter clinical trial. Paroxysmal Atrial Fibrillation Italian Trial (PAFIT)-2 Investigators. *Clin Cardiol* 1996;**19**:409-12.
- 70 **Galve E**, Rius T, Ballester R, *et al*. Intravenous amiodarone in treatment of recent-onset atrial fibrillation: results of a randomized, controlled study. *J Am Coll Cardiol* 1996;**27**:1079-82.
- 71 **Larbuissou R**, Venneman I, Stiels B. The efficacy and safety of intravenous propafenone versus intravenous amiodarone in the conversion of atrial fibrillation or flutter after cardiac surgery. *J Cardiothorac Vasc Anesth* 1996;**10**:229-34.
- 72 **Stambler BS**, Wood MA, Ellenbogen KA, *et al*. Efficacy and safety of repeated intravenous doses of ibutilide for rapid conversion of atrial flutter or fibrillation. Ibutilide Repeat Dose Study Investigators. *Circulation* 1996;**94**:1613-21.
- 73 **Digitalis in Acute Atrial Fibrillation (DAAF) Trial Group**. Intravenous digoxin in acute atrial fibrillation: results of a randomized, placebo-controlled multicenter trial in 239 patients. *Eur Heart J* 1997;**18**:649-54.
- 74 **Azpitarte J**, Alvarez M, Baun O, *et al*. Value of single oral loading dose of propafenone in converting recent-onset atrial fibrillation: results of a randomized, double-blind, controlled study. *Eur Heart J* 1997;**18**:1649-54.
- 75 **Boriani G**, Biffi M, Capucci A, *et al*. Oral propafenone to convert recent-onset atrial fibrillation in patients with and without underlying heart disease: a randomized, controlled trial. *Ann Intern Med* 1997;**126**:621-5.
- 76 **Frost L**, Mortensen PE, Tingleff J, *et al*. Efficacy and safety of dofetilide, a new class III antiarrhythmic agent, in acute termination of atrial fibrillation or flutter after coronary artery bypass surgery. Dofetilide Post-CABG Study Group. *Int J Cardiol* 1997;**58**:135-40.
- 77 **Innes GD**, Vertesi L, Dillon EC, *et al*. Effectiveness of verapamil-quinidine versus digoxin-quinidine in the emergency department treatment of paroxysmal atrial fibrillation. *Ann Emerg Med* 1997;**29**:126-34.
- 78 **Jordaens L**, Trouerbach J, Calle P, *et al*. Conversion of atrial fibrillation to sinus rhythm and rate control by digoxin in comparison to placebo. *Eur Heart J* 1997;**18**:643-8.
- 79 **Stroobandt R**, Stiels B, Hoebrechts R. Propafenone for conversion and prophylaxis of atrial fibrillation. Propafenone Atrial Fibrillation Trial Investigators. *Am J Cardiol* 1997;**79**:418-23.
- 80 **Bianconi L**, Mennuni M. Comparison between propafenone and digoxin administered intravenously to patients with acute atrial fibrillation. PAFIT-3 Investigators. The propafenone in atrial fibrillation Italian trial. *Am J Cardiol* 1998;**82**:584-8.
- 81 **Ganau G**, Lenzi T. Intravenous propafenone for converting recent onset atrial fibrillation in emergency departments: a randomized placebo-controlled multicenter trial. FAPS Investigators Study Group. *J Emerg Med* 1998;**16**:383-7.
- 82 **Kochiadakis GE**, Igoumenidis NE, Simantirakis EN, *et al*. Intravenous propafenone versus intravenous amiodarone in the management of atrial fibrillation of recent onset: a placebo-controlled study. *Pacing Clin Electrophysiol* 1998;**21**:2475-9.
- 83 **Kochiadakis GE**, Igoumenidis NE, Solomou MC, *et al*. Conversion of atrial fibrillation to sinus rhythm using acute intravenous procainamide infusion. *Cardiovasc Drugs Ther* 1998;**12**:75-81.
- 84 **Mattioli AV**, Lucchi GR, Vivoli D, *et al*. Propafenone versus procainamide for conversion of atrial fibrillation to sinus rhythm. *Clin Cardiol* 1998;**21**:763-6.
- 85 **Reisinger J**, Gatterer E, Heinze G, *et al*. Prospective comparison of flecainide versus sotalol for immediate cardioversion of atrial fibrillation. *Am J Cardiol* 1998;**81**:1450-4.
- 86 **Tisdale JE**, Padhi ID, Goldberg AD, *et al*. A randomized, double-blind comparison of intravenous diltiazem and digoxin for atrial fibrillation after coronary artery bypass surgery. *Am Heart J* 1998;**135**:739-47.
- 87 **Volgman AS**, Carberry PA, Stambler B, *et al*. Conversion efficacy and safety of intravenous ibutilide compared with intravenous procainamide in patients with atrial flutter or fibrillation. *J Am Coll Cardiol* 1998;**31**:1414-9.
- 88 **Vos MA**, Golitsyn SR, Stangl K, *et al*. Superiority of ibutilide (a new class III agent) over DL-sotalol in converting atrial flutter and atrial fibrillation. The Ibutilide/Sotalol Comparator Study Group. *Heart* 1998;**79**:568-75.
- 89 **Blanc JJ**, Voinov C, Maarek M. Comparison of oral loading dose of propafenone and amiodarone for converting recent-onset atrial fibrillation. PARSIFAL Study Group. *Am J Cardiol* 1999;**84**:1029-32.
- 90 **Cotter G**, Blatt A, Kaluski E, *et al*. Conversion of recent onset paroxysmal atrial fibrillation to normal sinus rhythm: the effect of no treatment and high-dose amiodarone. A randomized, placebo-controlled study. *Eur Heart J* 1999;**20**:1833-42.
- 91 **Geelen P**, O'Hara GE, Roy N, *et al*. Comparison of propafenone versus procainamide for the acute treatment of atrial fibrillation after cardiac surgery. *Am J Cardiol* 1999;**84**:345-7, A8-9.
- 92 **Kochiadakis GE**, Igoumenidis NE, Parthenakis FI, *et al*. Amiodarone versus propafenone for conversion of chronic atrial fibrillation: results of a randomized, controlled study. *J Am Coll Cardiol* 1999;**33**:966-71.
- 93 **Kochiadakis GE**, Igoumenidis NE, Solomou MC, *et al*. Efficacy of amiodarone for the termination of persistent atrial fibrillation. *Am J Cardiol* 1999;**83**:58-61.
- 94 **Norgaard BL**, Wachtell K, Christensen PD, *et al*. Efficacy and safety of intravenously administered dofetilide in acute termination of atrial fibrillation and flutter: a multicenter, randomized, double-blind, placebo-controlled trial. Danish Dofetilide in Atrial Fibrillation and Flutter Study Group. *Am Heart J* 1999;**137**:1062-9.
- 95 **Vanderlugt JT**, Mattioni T, Denker S, *et al*. Efficacy and safety of ibutilide fumarate for the conversion of atrial arrhythmias after cardiac surgery. *Circulation* 1999;**100**:369-75.
- 96 **Capucci A**, Villani GQ, Aschieri D, *et al*. Oral amiodarone increases the efficacy of direct-current cardioversion in restoration of sinus rhythm in patients with chronic atrial fibrillation. *Eur Heart J* 2000;**21**:66-73.
- 97 **Peuhkurinen K**, Niemela M, Ylitalo A, *et al*. Effectiveness of amiodarone as a single oral dose for recent-onset atrial fibrillation. *Am J Cardiol* 2000;**85**:462-5.
- 98 **Vardas PE**, Kochiadakis GE, Igoumenidis NE, *et al*. Amiodarone as a first-choice drug for restoring sinus rhythm in patients with atrial fibrillation: a randomized, controlled study. *Chest* 2000;**117**:1538-45.
- 99 **Alp NJ**, Bell JA, Shahi M. Randomised double blind trial of oral versus intravenous flecainide for the cardioversion of acute atrial fibrillation. *Heart* 2000;**84**:37-40.
- 100 **Bianconi L**, Castro A, Dinelli M, *et al*. Comparison of intravenously administered dofetilide versus amiodarone in the acute termination of atrial fibrillation and flutter: a multicentre, randomized, double-blind, placebo-controlled study. *Eur Heart J* 2000;**21**:1265-73.
- 101 **Hohnloser SH**, Kuck KH, Lilienthal J. Rhythm or rate control in atrial fibrillation: pharmacological intervention in atrial fibrillation (PIAF): a randomised trial. *Lancet* 2000;**356**:1789-94.
- 102 **Joseph AP**, Ward MR. A prospective, randomized controlled trial comparing the efficacy and safety of sotalol, amiodarone, and digoxin for the reversion of new-onset atrial fibrillation. *Ann Emerg Med* 2000;**36**:1-9.
- 103 **Kumagai K**, Abe H, Hiraki T, *et al*. Single oral administration of pilsicainide versus infusion of disopyramide for termination of paroxysmal atrial fibrillation: a multicenter trial. *Pacing Clin Electrophysiol* 2000;**23**:1880-2.
- 104 **Lindeboom JE**, Kingma JH, Crijns HJ, *et al*. Efficacy and safety of intravenous dofetilide for rapid termination of atrial fibrillation and atrial flutter. *Am J Cardiol* 2000;**85**:1031-3.
- 105 **Madonia S**, De Simone M, Brai G, *et al*. Intravenous versus oral initial load of propafenone for conversion of recent-onset atrial fibrillation in the emergency room: a randomized trial. *Ital Heart J* 2000;**1**:475-9.
- 106 **Martinez-Marcos FJ**, Garcia-Garmendia JL, Ortega-Carpio A, *et al*. Comparison of intravenous flecainide, propafenone, and amiodarone for conversion of acute atrial fibrillation to sinus rhythm. *Am J Cardiol* 2000;**86**:950-3.
- 107 **Singh S**, Zoble RG, Yellen L, *et al*. Efficacy and safety of oral dofetilide in converting to and maintaining sinus rhythm in patients with chronic atrial fibrillation or atrial flutter: the symptomatic atrial fibrillation investigative research on dofetilide (SAFIRE-D) study. *Circulation* 2000;**102**:2385-90.
- 108 **Pedersen OD**, Bagger H, Keller N, *et al*. Efficacy of dofetilide in the treatment of atrial fibrillation-flutter in patients with reduced left ventricular function: a Danish investigations of arrhythmia and mortality on dofetilide (diamond) substudy. *Circulation* 2001;**104**:292-6.
- 109 **Wattanasuwan N**, Khan IA, Mehta NJ, *et al*. Acute ventricular rate control in atrial fibrillation: IV combination of diltiazem and digoxin vs. IV diltiazem alone. *Chest* 2001;**119**:502-6.
- 110 **Waldo AL**, Camm AJ, deRuiter H, *et al*. Effect of d-sotalol on mortality in patients with left ventricular dysfunction after recent and remote myocardial infarction. The SWORD Investigators. Survival with oral d-sotalol. *Lancet* 1996;**348**:7-12.

- 111 **Torp-Pedersen C**, Møller M, Bloch-Thomsen PE, *et al*. Dofetilide in patients with congestive heart failure and left ventricular dysfunction. *N Engl J Med* 1999;**341**:857–65.
- 112 **Cardiac Arrhythmia Suppression Trial (CAST) Investigators**. Preliminary report: effect of encainide and flecainide on mortality in a randomized trial of arrhythmia suppression after myocardial infarction. *N Engl J Med* 1989;**321**:406–12.
- 113 **Echt D**, Liebson P, Mitchell B, *et al*. Mortality and morbidity in patients receiving encainide, flecainide or placebo (CAST). *N Engl J Med* 1991;**324**:781–8.
- 114 **Cairns JA**, Connolly SJ, Roberts R, *et al*. Randomised trial of outcome after myocardial infarction in patients with frequent or repetitive ventricular premature depolarizations: CAMIAT. Canadian Amiodarone Myocardial Infarction Arrhythmia Trial Investigators. *Lancet* 1997;**349**:675–82.
- 115 **Burkhardt F**, Pfisterer M, Kiowski W, *et al*. Effect of antiarrhythmic therapy on mortality in survivors of myocardial infarction with asymptomatic complex ventricular arrhythmias: Basel antiarrhythmic study of infarct survival (BASIS). *J Am Coll Cardiol* 1990;**16**:1711–8.
- 116 **Cerumuzynski L**, Kleczar E, Krzeminska-Pakula M, *et al*. Effect of amiodarone on mortality after myocardial infarction: a double-blind placebo-controlled pilot study. *J Am Coll Cardiol* 1992;**20**:1056–62.
- 117 **Julian DG**, Camm AJ, Frangin G, *et al*. Randomised trial of effect of amiodarone on mortality in patients with left-ventricular dysfunction after recent myocardial infarction: EMIAT. European Myocardial Infarct Amiodarone Trial Investigators. *Lancet* 1997;**349**:667–74.
- 118 **Eckman MH**, Falk RH, Pauker SG. Cost-effectiveness of therapies for patients with nonvalvular atrial fibrillation. *Arch Intern Med* 1998;**158**:1669–77.
- 119 **Catherwood E**, Fitzpatrick WD, Greenberg ML, *et al*. Cost-effectiveness of cardioversion and antiarrhythmic therapy in nonvalvular atrial fibrillation. *Ann Intern Med* 1999;**130**:625–36.
- 120 **Moher D**, Pham B, Klassen TP, *et al*. Does the language of publication of reports of randomized trials influence the estimates of intervention effectiveness reported in meta-analyses? Working paper. Ottawa: Thomas C. Chalmers Center for Systematic Reviews, 1999.
- 121 **Altman DG**, Deeks JJ, Sackett DL. Odds ratios should be avoided when events are common. *BMJ* 1999;**317**:1318.
- 122 **Deeks JJ**. Odds ratios should be used only in case-control studies and logistic regression analyses [letter]. *BMJ* 1998;**317**:1156.
- 123 **Bracken MB**, Sinclair JC. Avoidable systematic error in estimating treatment effects must not be tolerated [letter]. *BMJ* 1999;**317**:1156.
- 124 **Blitzer M**, Costeas C, Kassotis J, *et al*. Rhythm management in atrial fibrillation: with a primary emphasis on pharmacological therapy: part 1. *Pacing Clin Electrophysiol* 1998;**21**:590–602.

IMAGES IN CARDIOLOGY

Intermittent Brugada syndrome misdiagnosed as acute myocardial infarction and unmasked with propafenone

A 40 year old man was admitted to the emergency department with chest and left shoulder pain for an hour. His admission ECG revealed 3–5 mm ST segment elevation on leads V1–3 and right bundle branch block (RBBB). His previous ECGs were normal. Acute myocardial infarction was diagnosed and thrombolytic treatment initiated. Six hours later, there was no ST segment elevation on leads V1–3 or RBBB on the ECG, and all cardiac enzymes were within normal limits. Echocardiography was normal. Coronary angiography revealed normal coronary arteries. All ECGs were reviewed and it was decided that the diagnosis of acute myocardial infarction was incorrect. Because of typical ECG findings, an intermittent form of Brugada syndrome was suspected. To confirm the diagnosis (and because of the absence of ajmaline, flecainide, and procainamide) a provocation test with propafenone was performed. The test proved strongly positive for the Brugada syndrome. ECGs of his four family members revealed similar electrocardiographic abnormalities.

While magnetic resonance imaging (MRI) of the heart was normal, MRI of the neck delineated cervical herniation, which might explain the patient's symptoms.

Brugada syndrome can present with a typical ECG but in some cases the patient can present with concealed or intermittent forms. In the presented case propafenone unmasked ST segment elevation in leads V1–3 and RBBB. The blocking of the sodium channels and β adrenoreceptors contribute to the unmasking effect of propafenone.

This case indicates that Brugada syndrome may be confused with AMI, and that propafenone is useful for unmasking the Brugada syndrome ECG pattern.

I Akdemir
V Davutoglu
M Aksoy

ilyasakdemir@hotmail.com

Admission ECG revealed ST elevation and RBBB.

ECG findings normalised six hours later.

Unmasking effect of propafenone following the provocation test (20 minutes after).