REVIEW

Off-pump coronary artery bypass graft surgery: the incidence of postoperative atrial fibrillation

R A Archbold, N P Curzen

Heart 2003;89:1134–1137

Atrial fibrillation (AF) occurs in one quarter to one third of patients after coronary artery bypass graft surgery (CABG). Conventional CABG uses cardiopulmonary bypass, a process that is itself associated with a systemic vascular inflammatory response that contributes to postoperative morbidity. The avoidance of cardiopulmonary bypass is associated with a significant reduction in the inflammatory response and in the release of markers of myocardial necrosis when compared with conventional CABG. There is speculation that off-pump CABG may reduce the incidence of postoperative AF through reduced trauma, ischaemia, and inflammation. Current data, however, do not emphatically answer the question of whether the incidence of post-CABG AF is reduced by off-pump surgery. The evidence from both observational and randomised studies is conflicting and many studies have weaknesses in design, conduct, or interpretation. It remains an attractive hypothesis that postoperative AF is reduced by off-pump CABG but more robust data are required.

change in autonomic tone, or the response to cardiopulmonary bypass (CPB), may precipitate AF in these patients.

Conventional CABG employs CPB during the period of cardiac arrest. This provides for extracorporeal oxygenation of blood before returning it to the systemic circulation, thus ensuring adequate delivery to the vital organs. CPB is itself associated with a systemic inflammatory response syndrome that contributes to postoperative morbidity.⁴ Post-pump syndrome is characterised by leucocytosis, increased capillary permeability, accumulation of interstitial fluid, and organ dysfunction. It is thought to be triggered by surgical trauma, lung reperfusion, and contact of blood with the “foreign” extracorporeal bypass circuit.

Currently only circumstantial evidence suggests a causal relation between CPB and postoperative AF, a minority of investigators having reported an association between bypass time and post-CABG AF.⁵ Nevertheless, it remains an attractive hypothesis that the avoidance of CPB will reduce the incidence of post-CABG AF.

DOES OFF-PUMP CABG REDUCE THE INCIDENCE OF POST-CABG AF? THE TRIAL EVIDENCE

There is considerable interest in beating heart surgery without CPB and minimally invasive direct coronary artery bypass (MIDCAB), both of which may reduce postoperative morbidity. Off-pump CABG is associated with a significant reduction in the inflammatory response⁶ and in the release of markers of myocardial necrosis⁷ compared with conventional CABG. Furthermore, there are now data assessing the incidence of AF in patients revascularised by these techniques (tables 1 and 2).^{8–12}

In two series, post-CABG AF occurred in 18 (5.9%) of 306 patients who underwent left internal mammary artery grafting to the left anterior descending artery by MIDCAB²³ and was recorded in only 34 (1.7%) of 2052 patients who had off-pump CABG by median sternotomy.²⁵ In a study of patients who underwent CABG by MIDCAB (n = 93) or conventional operation (n = 74), AF was less frequent (7% v 31%, respectively; p < 0.05) in the group without CPB.¹⁰ By contrast, different series have reported rates of AF

Approximately 27 000 people in the UK and 500 000 people in the USA undergo coronary artery bypass graft surgery (CABG) annually, 25–30% of whom develop atrial fibrillation (AF) postoperatively. AF is associated with increased morbidity and expenditure, and prolonged hospital stay after CABG.¹ Advances in surgical techniques have not been accompanied by a decrease in the incidence of post-CABG AF. However, the use of “off-pump” CABG may provide a novel solution to this problem. This review analyses the current evidence regarding the incidence of AF after off-pump CABG.

PATHOGENESIS OF POST-CABG AF: A ROLE FOR THE “POST-PUMP SYNDROME”?

The pathogenesis of post-CABG AF is incompletely understood. Evidence suggests that the patients at greatest risk of developing the arrhythmia have an identifiable electrophysiological substrate.² The nature of this substrate may be slow atrial conduction or dispersion of atrial refractoriness.³ A number of perioperative factors, such as inadequate myocardial protection, electrolyte imbalance, β blocker withdrawal,

See end of article for authors' affiliations

Correspondence to:
Dr R A Archbold,
Department of Cardiology,
St Bartholomew's Hospital,
Dominion House,
60 Bartholomew Close,
West Smithfield, London
EC1A 7BE, UK;
Andrew.Archbold@
bartsandthelondon.nhs.uk

Accepted 7 April 2003

Table 1 Incidence of AF after conventional and minimally invasive direct CABG

Study, year (reference)	Incidence of postoperative AF		p Value
	CPB	Non-CPB	
Allen 1997* (8)	7/12 (58%)	0/23 (0%)	0.0001
Collard 1997 (9)	NA	1/10 (10%)	NA
Allen 1998 (10)	23/74 (31%)	6/93 (7%)	<0.05
Calafiore 1998 (11)	NA	35/434 (8%)	NA
Jansen 1998 (12)	NA	5/54 (9.3%)	NA
Tamis 1998 (13)	11/33 (33%)	11/42 (26%)	NS
Arom 1999 (14)	39/243 (16%)	8/44 (18%)	NA
Cohn 1999 (15)	11/55 (20%)	13/55 (24%)	0.6
Abreu 1999 (16)	192/747 (26%)	4/34 (12%)	0.06
Turner Jr 1999 (17)	NA	5/100 (5%)	NA
Siebert 2000 (18)	7/39 (18%)	6/24 (25%)	NS
Stamou 2000 (19)	NA	40/292 (14%)	NA
Koutlas 2000† (20)	57/220 (26%)	14/53 (26%)‡	NS
d'Amato 2000 (21)	14/42 (28%)	4/96 (4%)	0.003
Tamis-Holland 2000 (22)	42/108 (39%)	23/101 (23%)	0.02
Cremer 2000 (23)	NA	18/306 (5.9%)	NA
Hravnak 2001 (24)	238/720 (33.1%)	22/94 (23.4%)	0.059

*Single vessel repeat coronary artery bypass graft surgery; †≥75 years; ‡majority by minimally invasive direct coronary artery bypass.
AF, atrial fibrillation; CPB, cardiopulmonary bypass; NA, not applicable.

of up to 37% despite no CPB.²⁶ while other observational studies have shown no difference in AF incidence post-CABG with and without CPB.^{16, 20}

Theoretically, any reduction in AF incidence after off-pump CABG may be attributed to the less invasive nature of MIDCAB procedures than of median sternotomy rather than

to the absence of CPB. In 969 patients who underwent CABG without CPB, for example, MIDCAB was associated with a lower incidence of AF (14% v 25%; odds ratio 0.4, 95% confidence interval 0.3 to 0.7; $p < 0.001$) than a median sternotomy approach.¹⁹ Specific information about the impact of CPB on postoperative AF therefore comes from studies

Table 2 Incidence of AF after conventional and off-pump CABG by median sternotomy

Study, year (reference)	Incidence of postoperative AF		p Value
	CPB	Non-CPB	
Collard 1997 (9)	NA	22/66 (33%)	NA
Jansen 1998 (12)	NA	7/46 (15.2%)	NA
Tasdemir 1998 (25)	NA	34/2052 (1.7%)	NA
Saatvedt 1999 (26)	248/685 (36%)	7/19 (37%)	NS
Spooner 1999 (27)	NA	44/331 (13.3%)*	NA
Boyd 1999 (28)	17/60 (28.3%)	3/30 (10%)†	<0.05
Ardehali 1999 (29)	NA	5/26 (19%)	NA
Arom 1999 (14)	39/243 (16%)	8/62 (13%)	NS
Cartier 2000 (30)	580/1870 (31%)	90/300 (30%)	0.8
Siebert 2000 (18)	7/39 (18%)	7/24 (29%)	NS
Stamou 2000 (19)	NA	166/677 (25%)	NA
Stamou 2000*** (31)	12/41 (29%)	13/91 (14%)‡	0.04
Ascone 2000 (32)	39/100 (39%)	8/100 (8%)	0.001
Arom 2000 (33)	721/3171 (23%)	50/350 (14%)	<0.001
Kshetry 2000 (34)	131/609 (21.5%)	24/135 (17.8%)	NS
Hart 2000 (35)	NA	236/1582 (14.9%)§	NA
Arom 2000¶ (36)	31/132 (23%)	6/45 (13.3%)	0.154
Hernandez 2000 (37)	102/445 (22.9%)	63/332 (19.0%)	NS
Omeroglu 2000 (38)	NA	5/70 (7.1%)	NA
Bedi 2000 (39)	NA	1/100 (1%)	NA
Kilger 2001 (40)	9/53 (17%)	2/50 (4%)	<0.05
Czerny 2001 (41)	7/40 (17.5%)	5/40 (12.5%)	NS
Bull 2001 (42)	7/40 (18%)	7/40 (18%)	NS
Yeatman 2001** (43)	38/312 (12.2%)	9/75 (12%)	0.95
Bowles 2001 (44)	24/70 (34%)	22/67 (33%)	NS
Calafiore 2001 (45)	111/924 (12%)	110/919 (12%)	NS
Siebert 2001 (46)	64/650 (9.8%)	12/118 (10.2%)	0.9††
Arom 2001‡‡ (47)	NA	56/342 (16.4%)	NA
Puskas 2001 (48)	158/1000 (15.8%)	25/200 (12.5%)	0.28
Demers 2001§§ (49)	268/497 (54%)	41/98 (42%)	0.05
Lund 2001 (50)	109/368 (29.6%)	37/165 (22.4%)	NS
Van Dijk 2001 (51)	29/139 (21%)	28/142 (20%)	0.79
Al-Ruzzeh 2001¶¶ (52)	35/84 (41%)	16/56 (29%)	0.15

*Mixed approaches; †>70 years old and 93% by median sternotomy; ‡56% by median sternotomy; §83% by median sternotomy; ¶ejection fraction ≤30%; **left main stenosis ≥50%; ††AF incidence on intensive care unit (mean two days); ‡‡ejection fraction >30%; §§≥70 years; ¶¶≥75 years; ***single vessel repeat coronary artery bypass graft surgery.

comparing patients undergoing median sternotomy with and without CPB. Thus, AF occurred in 23% of 3171 patients who underwent conventional CABG compared with 14% of 350 off-pump operations performed through sternotomy ($p < 0.001$).³³ Objective comparison is, however, difficult since the off-pump patients received fewer grafts, had shorter surgery, and were ventilated for a shorter period. Furthermore, in other studies, the incidence of AF was not significantly different in groups undergoing CABG by median sternotomy with and without CPB.^{26, 48}

The importance of considering differences in clinical characteristics is illustrated by a recent study in which the incidence of post-CABG AF was significantly lower (23% *v* 39%; $p = 0.02$) among 101 MIDCAB patients than in 108 on-pump patients.²² The MIDCAB patients were younger, had a lower frequency of previous cardiac surgery, had less extensive coronary disease, received fewer grafts, and had a shorter postoperative hospital stay. After adjustment for these differences, surgical method was no longer a significant predictor of AF.

To elucidate the role of CPB properly in post-CABG AF, randomised data are required. In one study, 200 low risk patients were randomly assigned to off-pump or conventional CABG.³² Sustained AF (> 10 minutes) was significantly less common (8% *v* 39%; $p = 0.001$) in the off-pump group. Indeed CPB was the only independent predictor of postoperative AF. These data, however, contrast with those from a more recent trial in which 281 patients undergoing first, elective, isolated CABG were randomly assigned to on- or off-pump surgery.⁵¹ All patients received sotalol prophylaxis. Off-pump CABG resulted in reduced creatine kinase MB fraction release, reduced use of blood products, and a shorter hospital stay compared with on-pump surgery, but nevertheless the incidence of AF (20% *v* 21%; respectively; $p = 0.79$) was similar.

DISCUSSION

Post-CABG AF continues to affect morbidity, hospital stay, and financial resources. There is speculation that off-pump CABG may reduce the incidence of AF, but current data do not answer this hypothesis. The evidence is conflicting. Interpretation of studies is confounded by patient selection, inappropriate control groups, and poorly defined arrhythmia monitoring.

Alternative strategies for reducing both the inflammatory response to CPB and the incidence of post-CABG AF exist. β Blockers are generally accepted to prevent AF.⁵³ Some studies have reported a reduction in the serum concentration of inflammatory markers after CPB using heparin coated extracorporeal circuits, or by high dose aprotinin or prednisolone.⁵⁴ Whether these findings translate into clinical benefit, specifically a decrease in the incidence of post-CABG AF, remains uncertain.

Conclusion

Off-pump techniques offer several potential advantages over conventional CABG (shorter operation times, more rapid recovery times, reduced neurocognitive impairment, reduced need for blood products). It is uncertain whether a reduction in the incidence of postoperative AF can be added to this list. More robust data are required.

Authors' affiliations

R A Archbold, Department of Cardiology, St Bartholomew's Hospital, London, UK

N P Curzen, Manchester Heart Centre, Manchester Royal Infirmary, Manchester, UK

REFERENCES

- Archbold RA, Curzen N. Atrial fibrillation after coronary artery bypass graft surgery: more than an irritation! *Clin Intensive Care* 1999;10:109-16.
- Zaman AG, Archbold RA, Helft G, et al. Atrial fibrillation after coronary artery bypass surgery: a model for pre-operative risk stratification. *Circulation* 2000;101:1403-8.
- Cox JL, Schuessler RB, Boineau JP. The surgical treatment of atrial fibrillation I: summary of the current concepts of the mechanisms of atrial flutter and fibrillation. *J Thorac Cardiovasc Surg* 1991;101:402-5.
- Butler J, Rocker GM, Westaby S. Inflammatory response to cardiopulmonary bypass. *Ann Thorac Surg* 1993;55:552-9.
- Roffman JA, Fieldman A. Digoxin and propranolol in the prophylaxis of supraventricular tachyarrhythmias after coronary artery bypass surgery. *Ann Thorac Surg* 1981;31:496-501.
- Gu YJ, Mariana MA, van Oeveren W, et al. Reduction of the inflammatory response in patients undergoing minimally invasive coronary artery bypass grafting. *Ann Thorac Surg* 1998;65:420-4.
- Kilger E, Pichler B, Weis F, et al. Markers of myocardial ischaemia after minimally invasive and conventional coronary operation. *Ann Thorac Surg* 2000;70:2023-8.
- Allen KB, Matheny RG, Robison RJ, et al. Minimally invasive versus conventional reoperative coronary artery bypass. *Ann Thorac Surg* 1997;64:616-22.
- Collard E, Ducart A, Buche M, et al. Postoperative morbidity after coronary artery bypass graft surgery without cardiopulmonary bypass [abstract]. *Br J Anaesth* 1997;78(suppl 2):43.
- Allen KB, Matheny RG, Heimansohn DA, et al. Minimally invasive direct coronary artery bypass grafting versus single vessel conventional coronary artery bypass [abstract]. *J Am Coll Cardiol* 1998;31(suppl A):70A.
- Calafiore AM, Di Giammarco G, Teodori G, et al. Midterm results after minimally invasive coronary surgery (LAST operation). *J Thorac Cardiovasc Surg* 1998;115:763-71.
- Jansen EWL, Borst C, Lahpor JR, et al. Coronary artery bypass grafting without cardiopulmonary bypass using the octopus method: results in the first one hundred patients. *J Thorac Cardiovasc Surg* 1998;116:60-7.
- Tamis JE, Vloka ME, Malhotra S, et al. Atrial fibrillation is common after minimally invasive direct coronary artery bypass graft surgery [abstract]. *J Am Coll Cardiol* 1998;31(suppl A):118A.
- Arom KV, Emery RW, Flavin TF, et al. Cost-effectiveness of minimally invasive coronary artery bypass surgery. *Ann Thorac Surg* 1999;68:1562-6.
- Cohn WE, Sirois CA, Johnson RG. Atrial fibrillation after minimally invasive coronary artery bypass grafting: a retrospective, matched study. *J Thorac Cardiovasc Surg* 1999;117:298-301.
- Abreu JE, Reilly J, Salzano RP, et al. Comparison of frequencies of atrial fibrillation after coronary artery bypass grafting with and without the use of cardiopulmonary bypass. *Am J Cardiol* 1999;83:775-6.
- Turner WF Jr. "Off-pump" coronary artery bypass grafting: the first one hundred cases of the Rose City experience. *Ann Thorac Surg* 1999;68:1482-5.
- Siebert J, Rogowski J, Jagielak D, et al. Atrial fibrillation after coronary artery bypass grafting without cardiopulmonary bypass. *Eur J Cardiothorac Surg* 2000;17:520-3.
- Stamou SC, Dangas G, Hill PC, et al. Atrial fibrillation after beating heart surgery. *Am J Cardiol* 2000;86:64-7.
- Koutlas TC, Elbeery JR, Williams JM, et al. Myocardial revascularization in the elderly using beating heart coronary artery bypass surgery. *Ann Thorac Surg* 2000;69:1042-7.
- d'Amato TA, Savage EB, Wiechmann RJ, et al. Reduced incidence of atrial fibrillation with minimally invasive direct coronary artery bypass. *Ann Thorac Surg* 2000;70:2013-6.
- Tamis-Holland JE, Homel P, Durani M, et al. Atrial fibrillation after minimally invasive direct coronary artery bypass surgery. *J Am Coll Cardiol* 2000;36:1884-8.
- Cremer JT, Wittwer T, Böning A, et al. Minimally invasive coronary artery revascularization on the beating heart. *Ann Thorac Surg* 2000;69:1787-91.
- Hravnak M, Hoffman LA, Saul MI, et al. Atrial fibrillation: prevalence after minimally invasive direct and standard coronary artery bypass. *Ann Thorac Surg* 2001;71:1491-5.
- Tasdemir O, Vural KM, Karagoz H, et al. Coronary artery bypass grafting on the beating heart without the use of extracorporeal circulation: review of 2052 cases. *J Thorac Cardiovasc Surg* 1998;116:68-73.
- Saatvedt K, Fiane AE, Sellevold O, et al. Is atrial fibrillation caused by extracorporeal circulation? *Ann Thorac Surg* 1999;68:931-3.
- Spooner TH, Hart JC, Pym J. A two-year, three institution experience with the Medtronic Octopus: systematic off-pump surgery. *Ann Thorac Surg* 1999;68:1478-81.
- Boyd WD, Desai ND, Del Rizzo DF, et al. Off-pump surgery decreases postoperative complications and resource utilization in the elderly. *Ann Thorac Surg* 1999;68:1490-3.
- Ardehali A, Kessler D, Faroushani F, et al. Multivessel coronary artery bypass without cardiopulmonary bypass. *Am Heart J* 1999;138:983-6.
- Cartier R, Brann S, Dagenais F, et al. Systematic off-pump coronary artery revascularization in multivessel disease: experience of three hundred cases. *J Thorac Cardiovasc Surg* 2000;119:221-9.
- Stamou SC, Pfister AJ, Dangas G, et al. Beating heart versus conventional single-vessel reoperative coronary artery bypass. *Ann Thorac Surg* 2000;69:1383-7.
- Ascone R, Caputo M, Calori G, et al. Predictors of atrial fibrillation after conventional and beating heart coronary surgery: a prospective, randomized study. *Circulation* 2000;102:1530-5.
- Arom KV, Flavin TF, Emery RW, et al. Safety and efficacy of off-pump coronary artery bypass grafting. *Ann Thorac Surg* 2000;69:704-10.

- 34 **Kshetty VR**, Flavin TF, Emery RW, *et al.* Does multivessel, off-pump coronary artery bypass reduce postoperative morbidity? *Ann Thorac Surg* 2000;**69**:1725-31.
- 35 **Hart JC**, Spooner TH, Pym J, *et al.* A review of 1,582 consecutive octopus off-pump coronary bypass patients. *Ann Thorac Surg* 2000;**70**:1017-20.
- 36 **Arom KV**, Flavin TF, Emery RW, *et al.* Is low ejection fraction safe for off-pump coronary bypass operation? *Ann Thorac Surg* 2000;**70**:1021-5.
- 37 **Hernandez F**, Clough RA, Klemperer JD, *et al.* Off-pump coronary artery bypass grafting: initial experience at one community hospital. *Ann Thorac Surg* 2000;**70**:1070-2.
- 38 **Omeroglu SN**, Kurali K, Güler M, *et al.* Midterm angiographic assessment of coronary artery bypass grafting without cardiopulmonary bypass. *Ann Thorac Surg* 2000;**70**:844-50.
- 39 **Bedi HS**, Suri A, Kalkat MS, *et al.* Global myocardial revascularization without cardiopulmonary bypass using innovative techniques for myocardial stabilization and perfusion. *Ann Thorac Surg* 2000;**69**:156-64.
- 40 **Kilger E**, Weis FC, Goetz AE, *et al.* Intensive care after minimally invasive and conventional coronary surgery: a prospective comparison. *Intensive Care Med* 2001;**27**:534-9.
- 41 **Czerny M**, Baumer H, Kilo J, *et al.* Complete revascularization in coronary artery bypass grafting with and without cardiopulmonary bypass. *Ann Thorac Surg* 2001;**71**:165-9.
- 42 **Bull DA**, Neumayer LA, Stringham JC, *et al.* Coronary artery bypass grafting with cardiopulmonary bypass versus off-pump cardiopulmonary bypass grafting: does eliminating the pump reduce morbidity and cost? *Ann Thorac Surg* 2001;**71**:170-5.
- 43 **Yeatman M**, Caputo M, Ascione R, *et al.* Off-pump coronary artery bypass surgery for critical left main stem disease: safety, efficacy and outcome. *Eur J Cardiothorac Surg* 2001;**19**:239-44.
- 44 **Bowles BJ**, Lee JD, Dang CR, *et al.* Coronary artery bypass performed without the use of cardiopulmonary bypass is associated with reduced cerebral microemboli and improved clinical results. *Chest* 2001;**119**:25-30.
- 45 **Calafiore AM**, Di Mauro M, Contini M, *et al.* Myocardial revascularization with and without cardiopulmonary bypass in multivessel disease: impact of the strategy on early outcome. *Ann Thorac Surg* 2001;**72**:456-63.
- 46 **Siebert J**, Anisimowicz L, Lango R, *et al.* Atrial fibrillation after coronary artery bypass grafting: does the type of procedure influence the early postoperative incidence? *Eur J Cardiothorac Surg* 2001;**19**:455-9.
- 47 **Arom KV**, Emery RW, Flavin TF, *et al.* OPCAB surgery: a critical review of two different categories of pre-operative ejection fraction. *Eur J Cardiothorac Surg* 2001;**20**:533-7.
- 48 **Puskas JD**, Thourani VH, Marshall JJ, *et al.* Clinical outcomes, angiographic patency, and resource utilization in 200 consecutive off-pump coronary bypass patients. *Ann Thorac Surg* 2001;**71**:1477-84.
- 49 **Demers P**, Cartier R. Multivessel off-pump coronary artery bypass surgery in the elderly. *Eur J Cardiothorac Surg* 2001;**20**:908-12.
- 50 **Lund O**, Christensen J, Holme S, *et al.* On-pump versus off-pump coronary artery bypass: independent risk factors and off-pump graft patency. *Eur J Cardiothorac Surg* 2001;**20**:901-7.
- 51 **van Dijk D**, Nierich AP, Jansen EWL, *et al.* for the Octopus Study Group. Early outcome after off-pump versus on-pump coronary bypass surgery. Results from a randomized study. *Circulation* 2001;**104**:1761-6.
- 52 **Al-Ruzzeh S**, George S, Yacoub M, *et al.* The clinical outcome of off-pump coronary artery bypass surgery in the elderly patients. *Eur J Cardiothorac Surg* 2001;**20**:1152-6.
- 53 **Crystal E**, Connolly SJ, Sleik K, *et al.* Interventions on prevention of postoperative atrial fibrillation in patients undergoing heart surgery: a meta-analysis. *Circulation* 2002;**106**:75-80.
- 54 **Harig F**, Feyer R, Mahmoud FO, *et al.* Reducing the post-pump syndrome by using heparin-coated circuits, steroids, or aprotinin. *Thorac Cardiovasc Surg* 1999;**47**:111-8.

IMAGES IN CARDIOLOGY.....

Life threatening total occlusion of the left coronary artery after acetylcholine infusion

A 44 year old woman was referred to the hospital with anginal complaints and ST segment changes during exercise. A coronary angiogram was performed, but no stenoses were found. As her complaints persisted the question arose as to whether this could be caused by coronary spasm. Therefore it was decided to perform an acetylcholine provocation test. Acetylcholine was infused into the left main artery at a rate of 0.22 µg/min. After two minutes, the patient experienced severe anginal pain and the ECG showed ST segment elevations. Another angiogram was taken and revealed

severe coronary spasm of both the left anterior descending (LAD) and the circumflex (CX) coronary arteries (below left). A bolus of glyceryl trinitrate was given intracoronary to achieve vasodilation. This action was repeated, but with no satisfactory result. Subsequently, an intracoronary bolus of atropine (500 µg) was given as a last resort to terminate the spasm. After one dose of atropine the CX was open again, whereas the distal LAD remained closed (below centre). A second bolus was needed to normalise the angiogram (below right) as well as the ECG. Although a slight increase in troponin was found after the procedure,

no major complications occurred and the patient was discharged from hospital two days later.

This case shows that life threatening coronary spasms can occur during acetylcholine infusion in patients with "normal" coronary arteries. More important, glyceryl trinitrate is not always capable of reversing the evoked coronary spasm. Intracoronary atropine infusion may establish a satisfying vasodilating effect in these cases.

J S Wijkema
G A J Jessurun
R A Tio
 r.a.tio@thorax.azg.nl