Statin use is associated with lower risk of atrial fibrillation in women with coronary disease: the HERS trial

C N Pellegrini,¹ E Vittinghoff,² F Lin,² S B Hulley,² G M Marcus¹

See Editorial, p 693

¹ Division of Cardiology, Electrophysiology Section, University of California, San Francisco, California, USA;

² Department of Epidemiology and Biostatistics, University of California, San Francisco, California, USA

Correspondence to: Dr G M Marcus, 500 Parnassus Avenue, Room MUE434, Box 1354, San Francisco, CA 94143-1354, USA; marcusg@medicine.ucsf.edu

Accepted 14 October 2008
Published Online First
28 January 2009

ABSTRACT

Objective: To determine the efficacy of statin treatment in atrial fibrillation (AF) prevention in women.

Design: Cohort study using data obtained in the Heart and Estrogen/Progestin Replacement Study (HERS).

Setting: Secondary analysis of a multicentre, randomised controlled clinical trial.

Patients: 2673 Postmenopausal women with coronary disease.

Main outcome measures: AF prevalence at baseline and incident AF over a mean follow-up of 4.1 years.

Results: 88 Women with AF were identified: 29 at baseline and 59 during follow-up. Women with AF were significantly less likely to be taking a statin at study enrolment than those without AF (22% vs 37%, $p = 0.003$). Baseline statin use was associated with a 65% lower odds of having AF at baseline after controlling for age, race, history of myocardial infarction or revascularisation and history of heart failure (odds ratio 0.35, 95% confidence interval (CI) 0.13 to 0.93, $p = 0.04$). The risk of developing AF during the study among those free from AF at baseline, adjusted for the same covariates, was 55% less for those receiving statin treatment (hazard ratio 0.45, 95% CI 0.26 to 0.78, $p = 0.004$).

Conclusions: Statin treatment is associated with a lower prevalence and incidence of AF after adjustment for potential confounders in postmenopausal women with coronary disease.

Atrial fibrillation (AF) is the most common arrhythmia in the USA, affecting over two million people.¹ While women have a lower incidence rate of AF than men, they comprise half of those with the condition, owing to an over-representation of women among the elderly population.^{2–4} Several studies have demonstrated differences between women and men in the natural history of AF and incidence of its complications, with women experiencing more frequent recurrences of paroxysmal AF and higher rates of thromboembolism and stroke.^{5–11}

Inflammation is increasingly being understood to play an important role in the pathophysiology of AF,^{12–15} as shown, in part, by increased levels of inflammatory markers in those with AF^{14–17} and the presence of inflammation on their atrial biopsy specimens.¹⁸ This has led to the exploration of non-antiarrhythmic agents for AF prevention, with statins among the most promising. Recent studies have examined the role of statins for AF prevention in multiple settings^{19–36}; however, the vast majority of all study participants were men and none of the studies focused specifically on women.

Given the observed differences between women and men for multiple aspects of AF and the established fact that gender may significantly influence the effect of medicines, the efficacy of statin treatment for AF prevention may not be equivalent between the sexes. The Heart and Estrogen/Progestin Replacement Study (HERS) offers a unique opportunity to examine the role of statin treatment in a large cohort of women with coronary artery disease.

METHODS

The HERS design has been previously described in detail.³⁷ Briefly, 2763 postmenopausal women <80 years of age with known coronary heart disease (CHD) were randomised to receive either 0.625 mg conjugated equine oestrogens plus 2.5 mg medroxyprogesterone acetate in one daily pill ($n = 1380$) or identical placebo ($n = 1383$). CHD was documented in the form of myocardial infarction (MI) in 39% of study participants, $\geq 50\%$ narrowing of ≥ 1 coronary artery in 82%, catheter-based coronary revascularisation in 39% and surgical coronary revascularisation in 37%. The primary outcome was death from CHD or non-fatal MI. Average follow-up duration was 4.1 years.

For this analysis, baseline characteristics were obtained by patient interview at study enrolment. The presence of AF was assessed by standard 12-lead electrocardiograms (ECG) obtained at study enrolment and subsequently at yearly follow-up visits. A total of 2673 (97%) of the 2763 women included in the HERS trial met criteria for study inclusion. Subjects were excluded from the current analysis if they did not have a baseline ECG and at least one follow-up ECG ($n = 0$). In addition, subjects with atrial flutter ($n = 11$), wandering atrial pacemaker, supraventricular tachycardia, a paced rhythm, uncertain rhythm classification, or an uncertain detection of P waves were excluded ($n = 79$ for all other arrhythmias), and a sensitivity analysis was performed to determine if these exclusions meaningfully changed any of the results, which they did not. Study personnel were specifically trained in ECG acquisition and adequate ECG quality was certified at each clinical site. ECG interpretation was performed using NOVACODE software, which produces the classic Minnesota Code Classification.³⁸

Statin use was determined by inspection of prescription bottles and discussion with subjects at the time of study enrolment and at every 4-month follow-up visit thereafter. Start and stop dates of statin use were recorded. Additionally, use of

Table 1 Baseline characteristics of subjects categorised by the presence of atrial fibrillation at baseline or during follow-up, or both

Characteristics	Atrial fibrillation (n = 88)	No atrial fibrillation (n = 2585)	p Value
Age (years), mean (SD)	69.6 (5.9)	66.5 (6.6)	<0.001
Race			0.40
White	82 (93)	2285 (88)	
Black	4 (5)	208 (8)	
Hispanic	2 (2)	52 (2)	
Other	0 (0)	40 (2)	
Prior myocardial infarction or revascularisation	39 (44)	1322 (51)	0.21
History of heart failure	31 (35)	302 (12)	<0.001
Diabetes	25 (28)	585 (23)	0.21
Hypertension	57 (67)	1516 (60)	0.16
Left ventricular hypertrophy by ECG	12 (14)	257 (10)	0.24
Exercise >3 times/week	33 (38)	1007 (39)	0.78
Any alcohol use	31 (35)	1023 (40)	0.41
Body mass index (kg/m ²), mean (SD)	28.6 (5.9)	28.6 (5.5)	0.94
Baseline statin use	19 (22)	957 (37)	0.003
ACE inhibitor use	30 (34)	438 (17)	<0.001
Antiarrhythmic drug use	4 (4.5)	35 (1.4)	0.01

Results are presented as number (%) unless stated otherwise.

Vaughan Williams class IA, IC and III antiarrhythmic drugs was recorded at study enrolment and every follow-up visit thereafter.

Statistical analysis

The baseline characteristics of participants with either prevalent or incident AF were compared with those of other HERS participants using unpaired *t* tests, the χ^2 test, or Fisher's exact test as appropriate. Statin use was examined as a predictor of both AF prevalence and incidence. A logistic model was used to estimate the independent association between statin use and prevalent AF at baseline, and a Cox proportional hazards model was used to assess the independent relationship between statin

Table 2 Baseline characteristics of subjects categorised by statin use at study enrolment

Characteristics	Statin use (n = 976)	No statin use (n = 1697)	p Value
Age (years), mean (SD)	66.8 (6.2)	66.5 (6.9)	0.14
Race			0.01
White	883 (90)	1484 (87)	
Black	57 (6)	155 (9)	
Hispanic	18 (2)	36 (2)	
Other	18 (2)	22 (1)	
Prior myocardial infarction or revascularisation	470 (48)	891 (53)	0.03
History of heart failure	104 (11)	229 (13)	0.03
Diabetes	211 (22)	399 (24)	0.27
Hypertension	570 (59)	1003 (60)	0.71
Left ventricular hypertrophy by ECG	89 (9)	180 (11)	0.25
Exercise > 3 times per week	403 (41)	637 (38)	0.06
Any alcohol use	415 (43)	639 (38)	0.01
Body mass index (kg/m ²), mean (SD)	28.5 (5.3)	28.7 (5.6)	0.59
ACE inhibitor use	180 (18)	288 (17)	0.34
Antiarrhythmic drug use	10 (1)	29 (2)	0.16

Results are presented as number (%) unless stated otherwise.

use as a time-dependent covariate and AF detected later in the study. Covariates were selected for the multivariate models based on face validity (ie, race and history of MI or revascularisation) or an association with both the outcome and predictor of interest in univariate analyses with a *p* value <0.1. Women with AF at baseline were excluded from the analysis of incident AF. Statistical analyses were performed using SAS software V9.1 and were validated by a second, independent analyst.

RESULTS

Of the 2673 women constituting the study cohort, 29 were found to be in AF at the baseline visit, most of whom had AF noted at one or more follow-up visit also; there were 59 incident cases of AF during the study in women who were not in AF at baseline. Compared with those without AF, subjects with AF were older, more likely to have a history of heart failure and reported a higher use of ACE inhibitors at the time of study enrolment (table 1). Patients with AF were significantly less likely to be taking a statin at enrolment (22% vs 37%, *p* = 0.003).

To examine potential confounders of the relationship between statin use and AF, those taking a statin drug were compared with the study group as a whole. The women taking a statin drug at study enrolment were less likely to be black, have a history of heart failure or myocardial infarction or previous revascularisation, and more likely to report some alcohol use (table 2).

Baseline statin use was associated with an approximate 65% decrease in the odds of having AF at baseline (odds ratio (OR) = 0.36, 95% confidence interval (CI) 0.14 to 0.95, *p* = 0.04), with nearly identical results after adjusting for age, race, history of MI or revascularisation and history of heart failure (OR = 0.35, 95% CI 0.13 to 0.93, *p* = 0.04) (fig 1). Inclusion of antiarrhythmic drug use as a marker for AF treatment overall did not alter the results (adjusted OR for baseline statin use 0.35, 95% CI 0.13 to 0.92, *p* = 0.03). In addition, a sensitivity analysis that included atrial flutter with the outcome of AF did not produce an appreciably different result, nor did limiting the outcome to chronic persistent AF (ie, AF present on every ECG).

Statin use was also associated with a lower incidence of AF. The risk of developing AF during the study among those free of AF at baseline was approximately 55% lower for those receiving

Figure 1 Prevalence of atrial fibrillation at baseline based on the presence of predictors. Points display odds ratios, with adjustment for potential confounders. Bars represent 95% confidence intervals; upper confidence interval for race exceeds the scale. Age, 10-year increase in age; race, Caucasian compared with all others; MI/revasc, history of myocardial infarction or revascularisation.

Figure 2 Incidence of atrial fibrillation at follow-up based on presence of predictors. Points display hazard ratios, with adjustment for potential confounders. Bars represent 95% confidence intervals. Age, 10-year increase in age; race, Caucasian compared with all others; MI/revasc, history of myocardial infarction or revascularisation.

statin treatment (hazard ratio (HR) = 0.43, 95% CI 0.25 to 0.74, $p = 0.002$), again with nearly identical results after adjustment for age, race, history of MI or revascularisation, and history of heart failure (HR = 0.45, 95% CI 0.26 to 0.78, $p = 0.004$) (fig 2). History of heart failure and age were also positively associated with development of AF, with a 7% higher incidence of AF for every additional year of age (HR = 1.07, 95% CI 1.02 to 1.11, $p = 0.003$). There was no association between randomised treatment assignment (equine oestrogens plus medroxyprogesterone acetate versus placebo) and AF incidence.

DISCUSSION

As far as we know, this study is the first to extend the exploration of statin treatment in AF prevention to women, a group generally under-represented in the study of both statins and AF. Among the large cohort of postmenopausal women with coronary disease included in the HERS trial, those taking statins had a 65% lower prevalence at baseline and a 55% lower incidence of AF during the study after adjustment for potential confounders.

Statin treatment may prevent AF through multiple pathways, but their effects on inflammation and oxidative stress are probably paramount.¹² Evidence linking inflammation with AF includes data from atrial biopsies, inflammatory markers and measurements of atrial voltage among other sources.^{16–18 39 40} Statins may combat this process by inhibiting leukocyte–endothelium interaction and attenuating leukocyte infiltration,⁴¹ as well as by antagonising oxidant pathways,⁴² thereby protecting atrial tissue from damage and resultant structural remodelling. Statins increase nitric oxide synthesis,⁴³ potentially protecting the atrial myocardium during ischaemia.⁴⁴ It has also been postulated that they prevent electrical remodelling and alter the electrophysiological milieu through antisymphathetic activity, and have direct electrophysiological effects on ion channels and potential indirect effects via the angiotensin II-dependent pathway.⁴⁵ Finally, they may simply guard against development of a ready substrate for AF by decreasing cardiovascular ischaemic events and thereby preventing structural heart disease.⁴⁶

After animal studies demonstrated that statins prevent atrial remodelling and AF promotion in the canine atrial tachycardia and sterile pericarditis models,⁴⁷ human studies began to explore the possibility of statin treatment for AF prevention.^{28 29 32} A variety of settings have been employed, including the postoperative period after cardiac and thoracic surgery,^{19–24} the

post-pacemaker setting,²⁵ in patients with AF undergoing ablation and cardioversion procedures,^{26–31} among people with coronary artery disease and depressed left ventricular function^{32–35} and in those with paroxysmal AF.³⁶ A meta-analysis and five randomised trials have been published examining the effect of statin treatment for AF prevention.^{20 21 29 30 36 48} Both the meta-analysis and four of the five trials found that statins decreased AF occurrence.^{20 21 30 36 48} Importantly, as these trials were limited to subjects already with AF or the postoperative setting, none examined prevention of spontaneous AF, as was studied here. In addition, only one of the trials performed a subgroup analysis of the effect of statin treatment on AF in women, finding a reduction in postoperative AF occurrence in the 21 women included in the treatment group²⁰; the meta-analysis did not examine the effect of statins in women. Among the observational studies, results have been mixed, with many finding an inverse association between statin use and AF,^{17 20–22 26 29 30 32 33} and others not^{23–25 29 31} probably, to a large extent, because of the heterogeneity of statin treatments, study designs and subjects.

All these studies included, on average, four men for each woman, and none focused specifically on women. Thus whether the promise of statins in preventing AF would apply to women has remained uncertain until now. This is an important question because AF affects women differently than men. Women have higher heart rates when in AF^{5 6} and more frequent recurrences of paroxysmal AF.^{6 7} They sustain more thromboembolisms when not receiving warfarin,⁸ and may be at increased risk of stroke.^{9–11} Additionally, data regarding the pathophysiology of AF in women are generally unknown, particularly for inflammation as a potential cause of AF. The sex-based differences described in studies of other cardiovascular disease processes^{49–52} call for the specific examination of disease pathophysiology and pharmacology in women.

As expected, heart failure and older age were each significantly associated with both a higher prevalence and incidence of AF. Use of ACE inhibitors was higher in the AF group, apparently in contradiction to previous data, which demonstrated a protective effect of ACE inhibitors against AF⁵³; this probably represents confounding by indication (as subjects with heart failure are more likely to be treated with ACE inhibitors and to have AF). Use of ACE inhibitors was not significantly associated with statin use in the univariate model ($p = 0.26$), making it unlikely to confound the association between statin use and AF, and therefore it was not included in the multivariate model. As those with more cardiovascular disease and hence a higher risk of AF may be more likely to receive statins than not, confounding by indication is probably not responsible for our findings related to statins and AF (which would make statins appear to be associated with *more* AF). Of note, the low overall use of statin treatment among this population of women with CHD is probably owing to the less aggressive guidelines in place before 1998 when the follow-up for the HERS trial was completed, as well as imperfect compliance with guidelines in general.

Our study has several limitations. ECG diagnoses of AF were made with the aid of computer software as opposed to solely by human adjudicators. While the NOVACODE program has been well validated and extensively used in major clinical trials, its performance in the diagnosis of arrhythmias has not been formally tested. Imperfect sensitivity and specificity in diagnosing AF may have resulted, but as statin exposure is unlikely to affect ECG interpretation, any misclassification of the presence of AF would have been non-differential—that is, occurring to

the same extent in those with and without statin use. As a result, bias due to the use of this algorithm would be expected to drive our findings towards the null hypothesis, a type II error, suggesting that the magnitude and significance level of the observed associations somewhat understate the true associations.

An additional limitation of the study is the relatively infrequent (yearly) capture of ECGs and lack of additional mechanisms of AF diagnosis, such as patient questionnaire. However, these also would probably have affected both exposure groups equally, and therefore may have limited our ability to see a difference between groups, but should not have created a spurious association.

While we cannot confirm that women receiving statin treatment were not also being treated more aggressively for AF, such as with more cardioversions, adjustment for antiarrhythmic drug use did not alter our results, suggesting that this is probably not the case.

Finally, we cannot rule out the possibility that healthcare providers were less likely to use statin treatment in women with AF—for example, owing to concern about drug–drug interaction with simvastatin and warfarin. New occurrence of AF in our study was not associated with increased likelihood of cessation of statin treatment, nor was presence of AF associated with decreased likelihood of initiating statin treatment, suggesting that this was not a common concern of study healthcare providers.

CONCLUSIONS

For the first time, we have shown that statin treatment is associated with both a reduced incidence and prevalence of AF in a cohort of women. Specifically, in postmenopausal women with coronary heart disease, AF prevalence was 65% lower and AF incidence was 55% lower in those receiving statin treatment both before and after adjustment for multiple potential confounders. Women with AF may therefore be excellent candidates for new antiarrhythmic treatments aimed at the underlying atrial substrate.

Funding: This work was made possible by a fellowship grant from the American Heart Association Western States Affiliate (CNP), grant number KL2 RR024130 from the National Center for Research Resources, a component of the NIH (GMM), and an American Heart Association Beginning Grant-in-Aid Award (GMM).

Competing interests: None.

Ethics approval: Ethics committee approval obtained.

REFERENCES

- Rosamond W, Flegal K, Friday G, *et al.* Heart disease and stroke statistics—2007 update: a report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation* 2007;**115**:e69–171.
- Lloyd-Jones DM, Wang TJ, Leip EP, *et al.* Lifetime risk for development of atrial fibrillation: the Framingham Heart Study. *Circulation* 2004;**110**:1042–6.
- Feinberg WM, Blackshear JL, Laupacis A, *et al.* Prevalence, age distribution, and gender of patients with atrial fibrillation. Analysis and implications. *Arch Intern Med* 1995;**155**:469–73.
- Larsen JA, Kadish AH. Effects of gender on cardiac arrhythmias. *J Cardiovasc Electrophysiol* 1998;**9**:655–64.
- Hnatkova K, Waktare JE, Murgatroyd FD, *et al.* Age and gender influences on rate and duration of paroxysmal atrial fibrillation. *Pacing Clin Electrophysiol* 1998;**21**:2455–8.
- Humphries KH, Kerr CR, Connolly SJ, *et al.* New-onset atrial fibrillation: sex differences in presentation, treatment, and outcome. *Circulation* 2001;**103**:2365–70.
- Suttorp MJ, Kingma JH, Koomen EM, *et al.* Recurrence of paroxysmal atrial fibrillation or flutter after successful cardioversion in patients with normal left ventricular function. *Am J Cardiol* 1993;**71**:710–3.
- Fang MC, Singer DE, Chang Y, *et al.* Gender differences in the risk of ischemic stroke and peripheral embolism in atrial fibrillation: the Anticoagulation and Risk factors In Atrial fibrillation (ATRIA) study. *Circulation* 2005;**112**:1687–91.
- Cabin HS, Clubb KS, Hall C, *et al.* Risk for systemic embolization of atrial fibrillation without mitral stenosis. *Am J Cardiol* 1990;**65**:1112–6.
- Hart RG, Pearce LA, McBride R, *et al.* Factors associated with ischemic stroke during aspirin therapy in atrial fibrillation: analysis of 2012 participants in the SPAF I–III clinical trials. The Stroke Prevention in Atrial Fibrillation (SPAF) Investigators. *Stroke* 1999;**30**:1223–9.
- Wang TJ, Massaro JM, Levy D, *et al.* A risk score for predicting stroke or death in individuals with new-onset atrial fibrillation in the community: the Framingham Heart Study. *JAMA* 2003;**290**:1049–56.
- Issac TT, Dokainish H, Lakkis NM. Role of inflammation in initiation and perpetuation of atrial fibrillation: a systematic review of the published data. *J Am Coll Cardiol* 2007;**50**:2021–8.
- Boos CJ, Lip GY. Inflammation and atrial fibrillation: cause or effect? *Heart* 2008;**94**:133–4.
- Chung MK, Martin DO, Sprecher D, *et al.* C-reactive protein elevation in patients with atrial arrhythmias: inflammatory mechanisms and persistence of atrial fibrillation. *Circulation* 2001;**104**:2886–91.
- Dernellis J, Panaretou M. C-reactive protein and paroxysmal atrial fibrillation: evidence of the implication of an inflammatory process in paroxysmal atrial fibrillation. *Acta Cardiol* 2001;**56**:375–80.
- Marcus GM, Whooley MA, Glidden DV, *et al.* Interleukin-6 and atrial fibrillation in patients with coronary artery disease: data from the Heart and Soul Study. *Am Heart J* 2008;**155**:303–9.
- Kallergis EM, Manios EG, Kanoupakis EM, *et al.* The role of the post-cardioversion time course of hs-CRP levels in clarifying the relationship between inflammation and persistence of atrial fibrillation. *Heart* 2008;**94**:200–4.
- Frustaci A, Chimenti C, Bellocci F, *et al.* Histological substrate of atrial biopsies in patients with lone atrial fibrillation. *Circulation* 1997;**96**:1180–4.
- Marin F, Pascual DA, Roldan V, *et al.* Statins and postoperative risk of atrial fibrillation following coronary artery bypass grafting. *Am J Cardiol* 2006;**97**:55–60.
- Patti G, Chello M, Candura D, *et al.* Randomized trial of atorvastatin for reduction of postoperative atrial fibrillation in patients undergoing cardiac surgery: results of the ARMYDA-3 (Atorvastatin for Reduction of MYocardial Dysrhythmia After cardiac surgery) study. *Circulation* 2006;**114**:1455–61.
- Chello M, Patti G, Candura D, *et al.* Effects of atorvastatin on systemic inflammatory response after coronary bypass surgery. *Crit Care Med* 2006;**34**:660–7.
- Ozaydin M, Dogan A, Varol E, *et al.* Statin use before by-pass surgery decreases the incidence and shortens the duration of postoperative atrial fibrillation. *Cardiology* 2007;**107**:117–21.
- Mariscalco G, Lorusso R, Klersy C, *et al.* Observational study on the beneficial effect of preoperative statins in reducing atrial fibrillation after coronary surgery. *Ann Thorac Surg* 2007;**84**:1158–64.
- Amar D, Zhang H, Heerd PM, *et al.* Statin use is associated with a reduction in atrial fibrillation after noncardiac thoracic surgery independent of C-reactive protein. *Chest* 2005;**128**:3421–7.
- Amit G, Katz A, Bar-On S, *et al.* Association of statin therapy and the risk of atrial fibrillation in patients with a permanent pacemaker. *Clin Cardiol* 2006;**29**:249–52.
- Richter B, Derntl M, Marx M, *et al.* Therapy with angiotensin-converting enzyme inhibitors, angiotensin II receptor blockers, and statins: no effect on ablation outcome after ablation of atrial fibrillation. *Am Heart J* 2007;**153**:113–9.
- Al Chekakie MO, Akar JG, Wang F, *et al.* The effects of statins and renin-angiotensin system blockers on atrial fibrillation recurrence following atrial pulmonary vein isolation. *J Cardiovasc Electrophysiol* 2007;**18**:942–6.
- Siu CW, Lau CP, Tse HF. Prevention of atrial fibrillation recurrence by statin therapy in patients with lone atrial fibrillation after successful cardioversion. *Am J Cardiol* 2003;**92**:1343–5.
- Tveit A, Grundtvig M, Gundersen T, *et al.* Analysis of pravastatin to prevent recurrence of atrial fibrillation after electrical cardioversion. *Am J Cardiol* 2004;**93**:780–2.
- Ozaydin M, Varol E, Aslan SM, *et al.* Effect of atorvastatin on the recurrence rates of atrial fibrillation after electrical cardioversion. *Am J Cardiol* 2006;**97**:1490–3.
- Humphries KH, Lee M, Sheldon R, *et al.* Statin use and recurrence of atrial fibrillation after successful cardioversion. *Am Heart J* 2007;**154**:908–13.
- Young-Xu Y, Jabbour S, Goldberg R, *et al.* Usefulness of statin drugs in protecting against atrial fibrillation in patients with coronary artery disease. *Am J Cardiol* 2003;**92**:1379–83.
- Adabag AS, Nelson DB, Bloomfield HE. Effects of statin therapy on preventing atrial fibrillation in coronary disease and heart failure. *Am Heart J* 2007;**154**:1140–5.
- Ramani G, Zahid M, Good CB, *et al.* Comparison of frequency of new-onset atrial fibrillation or flutter in patients on statins versus not on statins presenting with suspected acute coronary syndrome. *Am J Cardiol* 2007;**100**:404–5.
- Hanna IR, Heeke B, Bush H, *et al.* Lipid-lowering drug use is associated with reduced prevalence of atrial fibrillation in patients with left ventricular systolic dysfunction. *Heart Rhythm* 2006;**3**:881–6.
- Dernellis J, Panaretou M. Effect of C-reactive protein reduction on paroxysmal atrial fibrillation. *Am Heart J* 2005;**150**:1064.
- Grady D, Applegate W, Bush T, *et al.* Heart and Estrogen/progestin Replacement Study (HERS): design, methods, and baseline characteristics. *Control Clin Trials* 1998;**19**:314–35.
- Rautaharju PM, Calhoun HP, Chaitman BR. NOVAECODE serial ECG classification system for clinical trials and epidemiologic studies. *J Electrocardiol* 1992;**24**:179–87.
- Aviles RJ, Martin DO, Apperson-Hansen C, *et al.* Inflammation as a risk factor for atrial fibrillation. *Circulation* 2003;**108**:3006–10.
- Marcus GM, Yang Y, Varosy PD, *et al.* Regional left atrial voltage in patients with atrial fibrillation. *Heart Rhythm* 2007;**4**:138–44.

41. **Ross R.** Atherosclerosis—an inflammatory disease. *N Engl J Med* 1999;**340**:115–26.
42. **Wassmann S,** Laufs U, Muller K, *et al.* Cellular antioxidant effects of atorvastatin in vitro and in vivo. *Arterioscler Thromb Vasc Biol* 2002;**22**:300–5.
43. **Laufs U,** La Fata V, Plutzky J, *et al.* Upregulation of endothelial nitric oxide synthase by HMG CoA reductase inhibitors. *Circulation* 1998;**97**:1129–35.
44. **Cai H,** Li Z, Goette A, *et al.* Downregulation of endocardial nitric oxide synthase expression and nitric oxide production in atrial fibrillation: potential mechanisms for atrial thrombosis and stroke. *Circulation* 2002;**106**:2854–8.
45. **Savelieva I,** Camm J. Statins and polyunsaturated fatty acids for treatment of atrial fibrillation. *Nat Clin Pract Cardiovasc Med* 2008;**5**:30–41.
46. **Kjekshus J,** Pedersen TR. Reducing the risk of coronary events: evidence from the Scandinavian Simvastatin Survival Study (4S). *Am J Cardiol* 1995;**76**:64C–8C.
47. **Shiroshita-Takeshita A,** Schram G, Lavoie J, *et al.* Effect of simvastatin and antioxidant vitamins on atrial fibrillation promotion by atrial-tachycardia remodeling in dogs. *Circulation* 2004;**110**:2313–9.
48. **Fauchier L,** Pierre B, de Labriolle A, *et al.* Antiarrhythmic effect of statin therapy and atrial fibrillation: a meta-analysis of randomized controlled trials. *J Am Coll Cardiol* 2008;**51**:828–35.
49. **Rathore SS,** Wang Y, Krumholz HM. Sex-based differences in the effect of digoxin for the treatment of heart failure. *N Engl J Med* 2002;**347**:1403–11.
50. **Hill MD,** Kent DM, Hinchey J, *et al.* Sex-based differences in the effect of intra-arterial treatment of stroke: analysis of the PROACT-2 study. *Stroke* 2006;**37**:2322–5.
51. **Berger JS,** Roncaglioni MC, Avanzini F, *et al.* Aspirin for the primary prevention of cardiovascular events in women and men: a sex-specific meta-analysis of randomized controlled trials. *JAMA* 2006;**295**:306–13.
52. **Andreotti F,** Marchese N. Women and coronary disease. *Heart* 2008;**94**:108–16.
53. **Healey JS,** Baranchuk A, Crystal E, *et al.* Prevention of atrial fibrillation with angiotensin-converting enzyme inhibitors and angiotensin receptor blockers: a meta-analysis. *J Am Coll Cardiol* 2005;**45**:1832–9.

Images in cardiology

Use of multislice CT coronary angiography to assess degree of left main stent overhang into the aorta

A 57-year-old symptomatic woman underwent emergency stenting of an isolated ostial lesion (video 1) in an unprotected left main coronary artery (LMCA) after becoming hypotensive and ischaemic during routine angiography. A 3.5×12 Taxus stent was deployed using the stent overhang technique (intentional protrusion of one to two stent struts into the aorta) and its placement and apposition of stent struts were confirmed by intravascular ultrasound.

Six months later, she was asymptomatic and had routine surveillance of her coronary arteries with routine coronary angiography and a 64-slice cardiac computed tomography angiography (CTA). Both angiographic studies confirmed the absence of significant in-stent restenosis (panels A and B, video 1). However, CTA clearly demonstrated a significant protrusion (measuring 8.3 mm) of proximal stent edge in the aorta (panel C, video 2), a finding not appreciated on angiographic images.

Isolated stenosis of the LMCA, specifically of the ostial segment, is an uncommon clinical entity predominantly affecting young women.¹ Despite controversy about the optimum revascularisation strategy of isolated LMCA stenosis, angioplasty with a drug-eluting stent is the preferred approach in emergent situations

(A) Anterior-posterior view; (B) left anterior oblique caudal view.

requiring prompt intervention. To achieve adequate lesion coverage at aorto-ostial locations, the stent is placed with slight overhang into the aorta and proximal stent struts flared using a post-dilatation balloon.² Excessive protrusion in the aorta, however, can make coaxial catheter

alignment difficult with risk of coronary dissection or stent deformation during subsequent interventions. Cardiac CTA may be preferable to coronary angiography for follow-up of patients with stents placed at aorto-ostial locations.

P Dehghani, D Marcuzzi, A N Cheema

Cheemaa@smh.toronto.on.ca

► Additional videos are published online only at <http://heart.bmj.com/content/vol95/issue9>

Heart 2009;**95**:708. doi:10.1136/hrt.2008.157362

REFERENCES

1. **Topaz O,** Warner M, Lanter P, *et al.* Isolated significant left main coronary artery stenosis: angiographic, hemodynamic, and clinical findings in 16 patients. *Am Heart J* 1991;**122**:1308–14.
2. **Hoffmann R,** Mintz GS, Mehran R, *et al.* Intravascular ultrasound predictors of angiographic restenosis in lesions treated with Palmaz-Schatz stents. *J Am Coll Cardiol* 1998;**31**:43–9.