and that PPAR α is an essential regulator of metabolism in hypoxia. PPAR α -/- mice (n=18) and wild-type (wt) controls (n=20) were exposed to 3 weeks of normobaric hypoxia. Control activated receptor α -/- (n=17) and wt mice (n=21) were housed in normoxic conditions within the same room. Oxygen content was reduced incrementally in the first week of housing, followed by 2 weeks at 11% oxygen. In-vivo cardiac function was measured using multislice cardiac magnetic resonance imaging. Hearts were perfused in the Langendorff mode to measure palmitate oxidation and glycolysis using 3H-labelled substrates. Cardiac output was unchanged in hypoxic wt and normoxic PPAR α -/- mice, but was reduced by 31% by hypoxia in PPAR α -/- mice ($p<0.02$). Late-stage ventricular filling was 46% lower in hypoxic PPAR α -/- mice ($p<0.01$). Hypoxia reduced palmitate oxidation by 27% in mouse wt hearts, but did not affect PPAR α -/- hearts. Hypoxia increased net lactate efflux 2.4-fold in hearts from wt animals ($p<0.01$), but lactate efflux from PPAR α -/- hearts was unchanged with hypoxia. Hypoxia increased basal glycolytic flux 2.4-fold in wt hearts but did not alter glycolytic flux in PPAR α -/- mouse hearts ($p<0.01$), which was already 3.7-fold greater than wt hearts. Thus PPAR α -/- hearts lack the metabolic flexibility essential for adaptation to chronic hypoxia, and their inability to upregulate glycolysis probably impairs cardiac function.

005 NOX2-DEFICIENT BONE MARROW-DERIVED MACROPHAGES EXHIBIT DEFECTS IN CELL SPREADING AND MIGRATION

doi:10.1136/hrt.2009.191049e

¹S Chaubey, ²C M Wells, ³G E Jones, ¹A M Shah, ¹A C Cave. ¹King's College London BHF Centre of Excellence, Cardiovascular Division, London, UK; ²King's College London, Division of Cancer Studies, London, UK; ³King's College London, Randall Division of Cell and Molecular Biophysics, London, UK

Introduction Our previous studies have demonstrated that the Nox2 isoform of NADPH oxidase is essential for the development of cardiac fibrosis in response to either angiotensin II or aldosterone. These in-vivo studies could not, however, identify the critical cell type responsible for this protective effect. As monocytes/macrophages are known to play an important role in the initiation of fibrosis, the current studies investigated whether Nox2 influences the ability of macrophages to migrate in response to the chemo-attractant, colony stimulating factor-1 (CSF-1).

Methods CSF-1-dependent bone-marrow derived macrophages (BMM) were isolated from femurs of wild-type (wt) and Nox2 knockout (Nox2^{-/-}) mice. BMM were seeded onto glass coverslips, starved of CSF-1, re-stimulated with CSF-1, fixed and stained for F-actin. Cell images were analysed for shape (elongation) and spread area using Image J software. Mean cell speed and chemotactic potential in a CSF-1 gradient were analysed using the Dunn Chemotaxis chamber.

Results A number of endpoints were assessed both under growing (basal) conditions and following CSF-1 stimulation. Under growing conditions Nox2^{-/-} BMM had a significantly increased spread area (proportion of the cell in contact with the glass coverslip) compared with wt cells (0.261 ± 0.03 vs 0.237 ± 0.03 ; $p<0.05$). However, while CSF-1 stimulation increased the spread area in wt cells above growing conditions (0.261 ± 0.01) Nox2^{-/-} BMM returned only to their initial spread area (0.256 ± 0.01). Nox2^{-/-} BMM were, however, significantly more elongated than wt cells following CSF-1 stimulation (0.670 ± 0.01 vs 0.624 ± 0.01). Importantly, Nox2^{-/-} BMM had a significantly reduced migration speed (0.46 ± 0.03 vs 0.6116 ± 0.02 $\mu\text{m}/\text{minute}$, $p<0.0004$), a significant reduction in their persistence of direction (0.15 ± 0.02 vs 0.38 ± 0.02 $p<0.0005$) and failed to migrate positively towards CSF-1. Nox2^{-/-} cells also had significantly lower increases in Akt and ERK1/2 phosphorylation following CSF-1 stimulation compared with wtT cells.

Conclusions Nox2^{-/-} BMM display marked abnormalities in morphological and migratory behaviour that may contribute significantly to the ability of the monocyte to differentiate and migrate in vivo in response to pathological stimuli. This phenotype could underlie the protection against fibrosis observed in vivo in Nox2^{-/-} mice.

006 NORMOBARIC HYPOXIA IMPAIRS CARDIAC ENERGETICS IN NORMAL HUMAN VOLUNTEERS

doi:10.1136/hrt.2009.191049f

^{1,2}C Holloway, ^{1,2}L Cochlin, ^{1,2}I Codreanu, ¹E Bloch, ¹M Fatemah, ²C Szmigielski, ¹H Atherton, ¹L Heather, ²J Francis, ²S Neubauer, ¹P Robbins, ¹K Clarke. ¹Department of Physiology, Anatomy and Genetics, University of Oxford, UK; ²University of Oxford Centre for Clinical Magnetic Resonance Research, John Radcliffe Hospital, Oxford, UK

Background In the first few days of hypoxic exposure, left ventricular dysfunction is consistently observed in the human heart, yet the cellular mechanisms underlying the dysfunction are poorly understood.

Objective Our hypothesis was that normobaric hypoxia impairs cardiac energetics, leading to cardiac dysfunction in healthy subjects.

Methods Normal healthy volunteers underwent 20 h of normobaric hypoxia in purpose-built hypoxia chambers. The partial pressure of oxygen during end tidal expiration (PETO₂) was kept between 50 and 60 mm Hg, while keeping peripheral oxygen saturation (SpO₂) above 80%. Cardiac function was measured using magnetic resonance imaging and echocardiography. High-energy phosphate metabolism was measured as the ratio of phosphocreatine to ATP (PCr/ATP) by ³¹P magnetic resonance spectroscopy before and after 20 h of hypoxia. Healthy men (n=12, aged 24 ± 2 years) were recruited from the University of Oxford.

Results During hypoxia, PETO₂ and SpO₂ averaged 55 ± 1 mm Hg and $83.6\pm0.4\%$, respectively. There was a 15% reduction in cardiac PCr/ATP, from 2.0 ± 0.1 to 1.7 ± 0.1 after hypoxia ($p<0.01$) and reduced diastolic function, measured as E/E', from 6.1 ± 0.4 to 7.5 ± 0.7 , $p<0.01$.

Conclusion Short-term normobaric hypoxia led to rapid changes in cardiac metabolism and alterations in diastolic function in normal human hearts. Impaired high-energy phosphate metabolism may explain the cardiac dysfunction observed after hypoxic exposure, whether in health or disease.

007 ROLES OF P47PHOX S303/S304 PHOSPHORYLATION IN TNF α -INDUCED ENDOTHELIAL REACTIVE OXYGEN SPECIES PRODUCTION AND MITOGEN-ACTIVATED PROTEIN KINASE ACTIVATION

doi:10.1136/hrt.2009.191049g

L Teng, J-M Li. Cardiovascular Research Theme, Faculty of Health and Medical Sciences, University of Surrey, UK

Endothelial cells express constitutively a Nox2 oxidase, which by generating reactive oxygen species (ROS) plays an important role in TNF α signalling. The Nox2 has at least four regulatory subunits and p47phox is a major regulatory subunit of this enzyme. It has been reported that phosphorylation of double serines (S303/S304) in p47phox is a key step for Nox2 activation. In this study, we investigated the role of p47phox S303/S304 phosphorylation in TNF α -induced ROS production and mitogen-activated protein kinase (MAPK) activation in endothelial cells. Serines 303/304 (human p47phox cDNA) were replaced to alanines by site-directed mutagenesis and the wild-type and mutated p47phox were used to transfect a mouse microvascular endothelial cell line (SVEC4-10). Forty-eight hours after transfection, cells were stimulated with or

without TNF α (100 U/ml for 30 minutes) and examined for ROS production (lucigenin emiluminescence) and MAPK activation (Western blot). Compared with cells transfected with an empty vector control) or cells transfected with wild-type p47phox (overexpression), S303/304A mutation had no significant effect on the basal level (without TNF α) of ROS production. When cells were stimulated with TNF α , there was a twofold increase in ROS production by control and p47phox overexpressed cells. TNF α -induced ROS production was significantly reduced (~60%) in cells transfected with the S303/304A mutation. However, there was no significant difference in TNF α -induced phosphorylation of ERK1/2, p38MAPK and JNK between controls and the cells transfected with the S303/304A mutation. In conclusion, p47phox phosphorylation at S303/S304 plays an important role in TNF α -induced ROS production, but is not involved in TNF α -induced MAPK activation.

008 NEURONAL NITRIC OXIDE SYNTHASE-MEDIATED REGULATION OF MYOCARDIAL REDOX STATE AND β 3 ADRENERGIC RECEPTOR RESPONSES

doi:10.1136/hrt.2009.191049h

W Idigo, Y H Zhang, S Reilly, M H Zhang, M Crabtree, N Alp, B Casadei. *Department of Cardiovascular Medicine, University of Oxford, John Radcliffe Hospital, Oxford, UK*

Recent evidence suggests that increased superoxide production may decrease the bioavailability of endothelial nitric oxide synthase (eNOS)-derived nitric oxide in the myocardium of neuronal nitric oxide synthase (nNOS)^{-/-} mice. To test the functional relevance of this observation, we compared the eNOS-dependent inotropic and [Ca²⁺]_i transient effect of beta3-adrenergic receptor (β 3-AR) stimulation in left ventricular myocytes from eNOS^{-/-} and nNOS^{-/-} mice and their wild-type littermates. β 3-AR stimulation (with BRL 37344, 10 μ mol/l in the presence of the β 1 and β 2-AR blocker nadolol, 10 μ mol/l) resulted in a significant reduction in cell shortening and [Ca²⁺]_i transient amplitude in wild-type myocytes. In contrast, BRL plus nadolol had no effect in myocytes from eNOS^{-/-} and nNOS^{-/-} myocytes (in the absence of changes in β 3-AR messenger RNA levels or in eNOS protein expression in the latter) or in nNOS^{+/+} myocytes pretreated with the nNOS-specific inhibitor, SMTC (0.1 μ mol/l). Inhibition of xanthine oxidoreductase (XOR) or NADPH oxidases caused a similar reduction in basal superoxide production in nNOS^{-/-} myocytes; however, whereas apocynin had no effect, XOR inhibition restored the negative inotropic response to β 3-AR stimulation in nNOS^{-/-} myocytes but not in eNOS^{-/-} myocytes. eNOS activity was uncoupled in the myocardium of nNOS^{-/-} mice in the absence of changes in BH4 and biopterin levels or arginase activity. However, L-arginine transport was impaired in nNOS^{-/-} myocytes and excess L-arginine (10 mmol/l) restored the response to β 3-AR stimulation. In summary, increased superoxide production by XOR (but not by NADPH oxidase) selectively abolished the negative inotropic effect of β 3-AR stimulation in nNOS^{-/-} myocytes by decreasing L-arginine transport and the bioavailability of eNOS-derived nitric oxide. These findings demonstrate that the source and subcellular localisation of superoxide production can account for the diverse and specialised actions of reactive oxygen species in the heart.

009 ENDOTHELIAL-SPECIFIC OVEREXPRESSION OF NOX2 ENHANCES ANGIOTENSIN II-INDUCED CARDIAC DYSFUNCTION AND FIBROSIS

doi:10.1136/hrt.2009.191049i

¹C E Murdoch, ¹A Brewer, ¹M Zhang, ²D Vanhoutte, ²S Heymans, ¹A M Shah. ¹King's College, London, UK; ²University Hospital, Maastricht, The Netherlands

Nox2-containing NADPH oxidases are reported to be involved in the development of cardiac fibrosis in response to chronic angiotensin II

infusion, but the cellular source(s) of Nox2 involved in fibrosis remains unclear. We investigated the role of endothelial Nox2 in angiotensin II-induced left ventricular hypertrophy (LVH). Male transgenic mice with endothelial-specific overexpression of Nox2 were compared with matched wild-type (wt) littermates after angiotensin II (1.1 mg/kg per day) or saline infusion for 14 days. Basal blood pressure and left ventricular NADPH oxidase activity were similar in wt and transgenic mice. After angiotensin II infusion, both wt and transgenic groups developed similar hypertension (170.2 $\pm$ 11.6 vs 170.4 $\pm$ 12.3 mm Hg; n=10) and hypertrophy (left ventricular/body weight ratio 4.8 $\pm$ 0.2 vs 4.7 $\pm$ 0.2 mg/g; and echocardiographic septal thickness increased by 34% wt and 37% transgenic mice; n>10). NADPH oxidase activity was higher in angiotensin II-infused transgenic compared with wt left ventricles (p<0.05). Interestingly, transgenic hearts showed significantly more interstitial cardiac fibrosis (2.3 $\pm$ 0.5 vs 1.3 $\pm$ 0.1% by Sirius red staining; p<0.05; n=6). In concurrence with a greater increase in inflammatory cell infiltration in transgenic left ventricles compared with wt (55% increase in CD45 staining; p<0.05; n=6). Left ventricular systolic function assessed in vivo by left ventricular pressure-volume analysis was similar in both angiotensin II groups (ESPVR, 8.9 $\pm$ 0.7 in wt vs 8.9 $\pm$ 1.3 mm Hg/ μ l in transgenic mice; p=ns; n=10), whereas diastolic stiffness increased significantly only in the angiotensin II transgenic group (EDPVR increased from 0.2 $\pm$ 0.0 to 0.5 $\pm$ 0.1 mm Hg/ μ l in transgenic mice; p<0.05; and from 0.2 $\pm$ 0.0 to 0.3 $\pm$ 0.0 mm Hg/ μ l in wt; n=10). These results indicate that endothelial-specific overexpression of Nox2 significantly enhances the development of angiotensin II-induced cardiac fibrosis and left ventricular diastolic dysfunction, independent of changes in hypertrophy.

Funding Supported by the British Heart Foundation.

010 CARDIOMYOCYTE-SPECIFIC OVEREXPRESSION OF NOX4 ATTENUATES ADVERSE CARDIAC REMODELLING AFTER MYOCARDIAL INFARCTION

doi:10.1136/hrt.2009.191049j

M Zhang, Y H Looi, A C Brewer, A M Shah. *King's College London, BHF Centre of Excellence, The James Black Centre, London, UK*

Background Reactive oxygen species (ROS) production is implicated in the development of cardiac remodelling after myocardial infarction (MI). NADPH oxidases are major sources of cardiovascular ROS, with cardiomyocytes expressing both Nox2 and Nox4 isoforms. We previously showed that Nox2 contributes significantly to the processes underlying adverse cardiac remodelling and contractile dysfunction post-MI, but the effect of Nox4 remains unclear. The aim of this study was to investigate the role of Nox4 in cardiac remodelling after MI.

Methods and Results We generated transgenic mice with cardiomyocyte-specific overexpression of Nox4, expressing the mouse full-length Nox4 complementary DNA under control of the α -MHC promoter. Transgenic mice were backcrossed onto a C57BL/6/J background and wild-type littermates (wt) used as controls. Nox4 transgenic mice were grossly similar to wt in terms of body weight, growth, activity, fertility and basal cardiac function. Transgenic mice displayed heart-specific expression of the Nox4 transgene, had significantly higher Nox4 protein expression, over 60% higher heart NADPH oxidase activity and 42% increased hydrogen peroxide production (all p<0.05). Compared with wt, p22phox protein expression was increased by 2.4-fold, which may be related to an increased protein stability. There was no change in the protein expression level of Nox2 or eNOS. MI was achieved by permanent left coronary ligation for 4 weeks. Compared with wt/MI, transgenic/MI groups displayed significantly high survival rate (72% vs 95%) and less hypertrophy at 4 weeks post-MI in terms of heart/body weight ratio (6.95 $\pm$ 0.16 vs 6.44 $\pm$ 0.14, p<0.01), although the