(CrCl) was calculated using Cockcroft-Gault formula. Patients were subgrouped into 5 grades based on preoperative CrCl; Group I CrCl ≥ 90 ml/min; II 60–89; III 30–59; IV 15–29; V < 15 or haemodialysis. Late Kaplan–Meier survival data (compared by log rank method), censored at 1/10/2009 were obtained from the UK CCAD. Surgical morbidity outcomes included re-exploration for bleeding, stroke (type 1 deficit) and low cardiac output state (LCOS) requiring inotropes $\pm$ intra-aortic balloon counterpulsation were compared using Fisher's Exact tests.

Results 1215 patients (921 males) with a mean age of 64 years (31–89 years) underwent CABG; 742 on OH and 472 on IN. Preoperative renal status in the groups were Group I -209(17%), II-584 (48%), III-387(32%), IV-26(2%) and V (8(1%). Similar percentages in each group had ≥ 1 grade deterioration of renal function post-operatively 19%, 18%, 16% and 23% (grades I–IV respectively; $p=0.470$). When examined as a continuous variable, higher preoperative CrCl correlated with a better postoperative improvement in CrCl ($r=0.073$, $p=0.012$ Spearman Rank). Overall 30-day mortality was 3.33% (CI 2.32 to 4.34%) and was not different by group I-3.37% (CI 0.92 to 5.82), II-2.09% (CI 0.92 to 3.26%), III 4.92% (CI 2.76 to 7.08%), IV 8% (CI 0 to 18.6%) and Stage V 0% (CI 0 to 0.4%; $p=0.101$) or by therapy type; ($p=0.411$). IN patients had similar preoperative renal function (median CrCl 66.8 vs 68.6; $p=0.828$) but a higher rate of postoperative renal deterioration (53.3 vs 46.7%, $p<0.001$). Stroke ($p=1.000$), bleeding ($p=0.755$) and LCOS ($p=0.335$) incidence were not different between therapy type. Overall mean survival was 9 years (CI 8.7 to 9.2 years) and was not different by renal function grade ($p=0.612$). However, IN patients had shorter mean survival 8.7 (8.3 to 9.0) vs OH 9.1(8.8 to 9.4) years; $p=0.03$.

Conclusions In T2DM-CABG, 36% of patients have CrCl < 60 ml/min. Higher CrCl protects against postoperative renal deterioration. Renal dysfunction does not appear to affect hospital outcome or survival. However, preoperative IN requirement increases the risk of renal dysfunction and is associated with worse longer-term survival.

58 TEMPORAL EVALUATION OF REFERRAL FOR AND LONG-TERM SURVIVAL FROM CARDIAC REHABILITATION FOR ACUTE MYOCARDIAL INFARCTION

doi:10.1136/heartjnl-2011-300198.58

¹C L Lewinter, ¹M B Bland, ²P D Doherty, ¹B L Lewin, ³A S H Hall, ⁴C P G Gale. ¹University of York, York, UK; ²York St John University, York, UK; ³Yorkshire Heart Centre, Leeds, UK; ⁴University of Leeds, Leeds, UK

Background Cardiac rehabilitation (CR) is a cost-effective, evidence-based approach to managing heart disease. Rates of uptake have and continue to vary despite recommendations from the NSF for CHD and NICE. The Evaluation of the Management and Methods of Acute Coronary Events (EMMACE) 1 and 2 studies are 2 large prospective multi-centre registries of care of acute coronary syndromes (ACS) in Yorkshire undertaken in 1995 and 2003 in respectively. We studied the temporal changes in referral for and long-term survival from CR in patients who were admitted to hospital with an acute myocardial infarction (AMI).

Methods Baseline characteristics were described as numbers (%) or as means with IQRs. For continuous variables, the Kruskal Wallis test was used for comparisons. Discrete variables were assessed by the χ^2 test. Unadjusted relative risk ratios (RRR) were calculated to assess mortality after referral for CR. Kaplan–Meier (KM) curves compared unadjusted survival stratified by CR referral and EMMACE study. Log rank tests compared the survival estimates. Sex, age, STEMI, heart failure, diabetes, COPD and mini-GRACE score, revascularisation, reperfusion, ACE-inhibitors, α -blockers, statins, anti-platelet agents and admitting cardiologist were regressed (backward logistic, $p<0.10$ and goodness of fit with a group of 10) on CR referral and represented as 95% CI OR. A Cox proportional model (Model 1: mini-GRACE score, Model 2: sex, age, STEMI, heart failure, diabetes, COPD,

EMMACE risk score, revascularisation, reperfusion, ACE-inhibitor, α -blocker, statins, anti-platelet agent, admitting cardiologist) was used to compare the temporal long-term survival estimates (all cause mortality) by CR referral.

Results 4341 had AMI. CR referral was 44% in 1995 and 59 % in 2003 ($p<0.001$). CR referral was associated with reduced mortality in 2003 (RRR, 95%CI: 0.54; 0.50 to 0.60), but was not in 1995 (1.02; 0.96 to 1.09). Unadjusted survival for patients not referral for CR in 1995 was similar to that for patients referred for CR in 1995; (Abstract 58 figure.1). For those referred for CR, the mean mini-GRACE score for CR referrals was lower in 2003 than 1995; 0.53 and 0.72, $p<0.001$. After adjustment using the min-GRACE score (Model 1), the impact (HR, 95% CI) of CR referral was 0.63, 0.55 to 0.73 in 2003 and 1.07, 0.92 to 1.3 in 1995. After adjustment using Model 2, the impact (HR, 95% CI) of CR referral was 0.57, 0.48 to 0.66 in 2003 and 1.31, 1.04 to 1.60 in 1995.

Conclusion Between 1995 and 2003, referral for CR increased and became a significantly important factor contributing to reduced mortality rates post-AMI. This is despite the differences in patient and treatment factors between the 2 studies periods. Even so, rate of referral for CR remain sub-optimal.

Abstract 58 Figure 1 Kaplan–Meier survival estimates.

59 SHORT TERM ELEVATION OF CHOLESTEROL LEVEL IN NEONATAL LIFE AND LONG TERM CHANGES IN AORTIC STIFFNESS: INSIGHTS FROM USE OF INTRAVENOUS LIPIDS

doi:10.1136/heartjnl-2011-300198.59

¹A J Lewandowski, ¹M Lazdam, ¹E Davis, ¹R Poole, ¹J Diesch, ¹J Francis, ¹S Neubauer, ²A Lucas, ²A Singhal, ¹B Kelly, ¹P Leeson. ¹Cardiovascular Medicine, University of Oxford, Oxford, UK; ²Institute of Child Health, University College London, London, UK

Introduction Offspring born to hypercholesterolaemic mothers have increased fatty streak formation in the fetal aorta, which persists into adolescence. To understand whether exposure to elevated cholesterol in early life, independent of a maternal history of hypercholesterolaemia, also has a long-term impact on the cardiovascular system we studied the vascular phenotype of adults in whom cholesterol levels were artificially elevated for a short period postnatally.

Methods We prospectively followed-up 102 subjects born premature now aged 23 to 28 years. Individuals exposed to maternal hypercholesterolaemia were excluded. 18 received intravenous (IV) lipids during the first nine weeks of life and were matched 2:1 for pregnancy and early life complications, age, sex, birthweight and gestational age with controls that did not receive IV lipids. Aortic pulse wave velocity (aPWV), regional aortic distensibility, left ventricular mass and ejection fraction were determined by cardiovascular

magnetic resonance. Detailed lifestyle information and anthropometric measurements were collected during childhood and adolescence. Metabolic parameters were measured multiple times per week for the first 9 weeks of life and again at follow-up visits.

Results Individuals that received IV lipids achieved significantly higher maximum cholesterol levels during the first 9 weeks of life than those that did not (mean $\pm$ SD=4.38 $\pm$ 1.65 vs 3.12 $\pm$ 0.78 mmol/l, $p=0.006$). Dose given and number of days on IV lipids also associated with maximum cholesterol level during this period ($r=0.557$, $p<0.001$ and $r=0.567$, $p<0.001$, respectively). There was a graded relation between the maximum elevation in circulating cholesterol postnatally and aortic stiffness (aPWV) in young adulthood ($r=0.596$, $p<0.001$). The greatest increase in stiffness was seen in the abdominal aorta, where distensibility was significantly reduced in the group that received IV lipids (mean $\pm$ SD=9.74 $\pm$ 4.27 vs 12.91 $\pm$ 4.11/mm Hg $\times 10^3$, $p=0.012$). There were no differences between the groups in other vascular or left ventricular measures. In a stepwise regression model, maximum cholesterol level achieved in the first few weeks of life was an independent predictor of aPWV in young adulthood ($\beta=0.596$, $p<0.001$) and accounted for 30.9% of the variance in hierarchical multiple regression ($\beta=0.584$, $p<0.001$).

Conclusions Brief artificial elevation of cholesterol level in immediate postnatal life is associated with long term changes in aortic function independent of later cholesterol levels. The association is graded depending on the degree of elevation of circulating cholesterol. High cholesterol exposure during sensitive periods of early postnatal life may have long term impacts on the cardiovascular system.

60

ETHNIC DIFFERENCES IN REPOLARISATION PATTERNS AND LEFT VENTRICULAR REMODELLING IN HIGHLY TRAINED MALE ADOLESCENT (14–18 YEARS) ATHLETES

doi:10.1136/heartjnl-2011-300198.60

¹N Sheikh, ¹M Papadakis, ²F Carre, ²G Kervio, ¹J Rawlins, ³V Panoulas, ¹N Chandra, ¹H Raju, ¹R Bastiaenen, ¹E Behr, ¹S Sharma. ¹St. George's University of London, London, UK; ²French Institute of Health and Medical Research, University of Rennes, Rennes, France; ³University Hospital Lewisham, London, UK

Purpose Studies in adult, black athletes (BA) demonstrate a high prevalence of ECG repolarisation changes and echocardiographic left ventricular hypertrophy (LVH) that may overlap with hypertrophic cardiomyopathy (HCM). The prevalence of ECG repolarisation changes and echocardiographic LVH in adolescent BA, the group most vulnerable to exercise-related sudden death from HCM, is unknown.

Methods This study evaluated 219 male adolescent BA (14–18 years, inclusive) with 12-lead ECG and 2-D echocardiography. Results were compared with 1440 male adolescent WA. Athletes with T wave inversions and morphological LVH were invited for further investigation with exercise stress test, 24 h Holter and CMR.

Results ST segment elevation was common in both groups but more frequent in BA (63.5% vs 14.9%, $p<0.001$), while ST segment depression was exceedingly rare. Both T wave inversions (21.5% vs 2.9%, $p<0.001$) and deep T wave inversions (11% vs 0.3%, $p<0.001$) were commoner in BA. Black athletes demonstrated greater left ventricular wall thickness (10.4 $\pm$ 1.6 vs 9.4 $\pm$ 1.2 mm, $p<0.001$) compared to WA. Twenty-three (10.5%) BA exhibited a left ventricular wall thickness >12 mm vs only 6 (0.4%) WA ($p<0.001$). None of the athletes exhibited the broader phenotype of HCM on further investigation. In multivariable analysis black ethnicity was the strongest independent predictor for the presence of T wave inversions (OR 3.56, 95% CI 1.56 to 8.13, $p=0.003$) and LVH (OR 3.17, 95% CI 1.77 to 5.71, $p<0.001$).

Conclusions As with adult athletes, T wave inversions and LVH were more prevalent in adolescent BA compared to WA. These findings have important implications in the pre-participation screening era, particularly in countries with a high proportion of BA competing at

elite level, since extrapolation of ECG and echocardiographic criteria, solely derived from Caucasian cohorts, would result in 25.6% of BA requiring further investigations for cardiac pathology.

Abstract 60 Figure 1 Bar chart depicting the distribution of left ventricular wall thickness in black and white adolescent athletes.

61

FIVE-MIN HEART RATE VARIABILITY CAN PREDICT OBSTRUCTIVE ANGIOGRAPHIC CORONARY DISEASE

doi:10.1136/heartjnl-2011-300198.61

¹D Kotecha, ²G New, ¹M Flather, ³D Eccleston, ³H Krum. ¹Royal Brompton Hospital, London, UK; ²Box Hill Hospital, Melbourne, UK; ³Monash University, Melbourne, UK

Background Obstructive coronary artery disease (CAD) is evident in only half of patients referred for diagnostic coronary angiography. Five-minute heart rate variability (HRV) is a marker for autonomic control of the vasculature, which we hypothesised could be used to risk-stratify cardiac patients (the Alternative Risk Markers in Coronary Artery Disease (ARM-CAD) study).

Methods Resting HRV prior to elective coronary angiography was analysed in 470 participants with predominantly normal cardiac rhythm, regardless of comorbidity. The presence of obstructive CAD ($\geq 50\%$ stenosis) was regressed in a multivariate model including risk factors, ECG variables and medications.

Results Mean age was 65 years (SD 11), 67% were male, 21% had diabetes, mean blood pressure was 144/79 mm Hg (SD 21/10) and 16% had impaired left-ventricular function. Patients with obstructive CAD had significantly reduced HRV, particularly in the low frequency (LF) range (median 180 vs 267 ms² without CAD; $p<0.001$). There was a linear trend according to the severity of CAD; the median LF power (IQR) in patients with normal coronaries was 275 (612), with minor coronary irregularities 255 (400),

Abstract 61 Figure 1