[gw22-e0410]

DIFFERENT VESSEL RESPONSE BETWEEN CHRONIC TOTAL OCCLUSIONS AND NON-CTO LESION AFTER SIROLIMUS-ELUTING STENT IMPLANTATION: A SERIAL OPTICAL COHERENCE TOMOGRAPHY STUDY

Jia Haibo¹, Li Ning¹, Gao Guangren¹, Wang Xuedong¹, Sun Meng², Wu Jian¹, Liu Fang¹, Yin Li¹, Zhang Maomao¹, Hu Sining¹, Hou Jingbo¹, Yu Bo¹, Zhang Shuo¹, Zhang Shaosong³, Jang Ik-kyung⁴ ¹Department of Cardiology, 2nd Affiliated Hospital of Harbin Medical University, Key Laboratories of Education Ministry for Myocardial Ischemia Mechanism and Treatment, Harbin, China; ²Department of Biostatistics Harbin Medical University, Harbin, China; ³Visiting Professor of Harbin Medical University, Light lab Imaging St Jude Medical, Westford, Maryland, United States; ⁴Cardiology Division, Massachusetts General Hospital, Harvard Medical School, Boston, United States

Objectives To investigate the neointimal coverage and malapposition of sirolimus-eluting stent in treatment of CTOs and non-CTO lesions (including lipid-rich plaque and non-lipid-rich plaque) by optical coherence tomography.

Methods We enrolled 64 patients (CTO: n=29 stents/21 patients, lipid-rich plaque: n=33 stents/26 patients, and non-lipid-rich plaque: n=21 stents/17 patients) with 68 target vessels and examined lesion characteristics by using OCT at pre- and post-SES implantation and 6 months follow-up. Lipid rich plaque was defined as the plaque with lipid content in ≥ 2 quadrants. Non-lipid-rich plaque consists of fibrous, fibrocalcific plaque, and lipid plaque with less than two quadrants lipid content.

Results The incidence of incomplete stent apposition was higher in CTO and lesion with lipid rich plaque compared with those with non-lipid rich plaque (3.03%, 2.58% vs 0.64%, $p=0.0219$). The tissue protrusion (14.99 % vs 11.00% vs 6.41%, $p<0.001$) and intra-stent thrombus (3.81% vs 2.39% vs 1.07%, $p=0.0119$) after percutaneous coronary intervention were observed most frequently in lipid-rich plaque group then followed by those in CTO group and non-lipid rich plaque group. At 6 months follow-up, the incidence of malapposition (5.0% vs 1.04% and 0.4%, $p=0.002$), the cross section with uncovered struts (23.4% vs 8.2% and 6.6%, $p<0.001$), and protrusion (30.8% vs 9.6% and 9.6%, $p<0.001$) was significantly greater in patients with CTO than those with lipid rich plaque and non-lipid rich plaque. Although the incidence of in-stent thrombus was non-significantly higher in CTO than LRP and non-LRP (9.4% vs 3.1% vs 0%, $p=0.26$), there was no adverse event in these patients with CTO.

Conclusions Initial lesion morphology is contributable to coronary artery response to DES after stent implantation. CTOs and lipid rich plaque is associated with higher incidence of malapposition, tissue protrusion, and mural thrombus after stenting. At 6 months follow-up, the vessels with CTO lesions were characterised by delayed healing and higher incidence of malapposition.