[gw22-e0005]

THE ANTI-HYPOXIA EFFECT OF GINSENOSES-RBL ON HYPOXIA CARDIOMYOCYTES MEDIATED BY THE APELIN-APJ SYSTEM EX VIVO

Hongliang Kong, Zhanquan Li, Long Yuan, Ru-ming Guan Cardiology Center, The People's Hospital Of Liaoning Province, Shenyang, China

10.1136/heartjnl-2011-300867.130

Objective On the basis of hypoxia induced by CoCl_2 , to elucidate whether Ginsenosides-Rbl (Gs-Rb1) inhibited the apoptosis of hypoxia cardiomyocytes by up-regulating Apelin-APJ system and whether the system was adjusted by Hif-1 α .

Materials and Methods Neonatal rat cardiomyocytes were randomly divided into control group, simple CoCl_2 group, simple Gs-Rb1 group, CoCl_2 and Gs-Rb1 hypoxia group, CoCl_2 and YC-1 group, CoCl_2 and YC-1 and Gs-Rb1 group; The concentration of CoCl_2 , Gs-Rb1 and YC-1 is 500 μM , 500 μM and 5 $\mu\text{mol/L}$, respectively; Apoptosis ratio (AR) was analysed by flow cytometer (FCM, Annexin V-FITC/PI), and Hif-1 α , Apelin and APJ were assayed by immunocytochemistry, Rt-PCR and Western blot.

Results (1) Gs-Rb1 significantly down-graded AR of hypoxia cardiomyocytes, which was significantly inhibited by YC-1; (2) Gs-Rb1 significantly increased the expression of hypoxia-induced factor 1 α (Hif-1 α), which was completely suppressed by YC-1; 3. Hypoxia significantly up-graded the expression of both mRNA and protein of Apelin, which were further reinforced by Gs-Rb1 and significantly inhibited by YC-1; 4. Gs-Rb1 further strengthen the expression of both mRNA and protein of APJ once hypoxia, which was significantly inhibited by YC-1; 5. There was a significant positive correlation between the AR and Apelin (including mRNA and protein), between the AR and APJ (including mRNA and protein).

Conclusion Apelin-APJ system, which was adjusted by Hif-1 α , plays an important role in Gs-Rb1 anti-hypoxia apoptosis of cardiomyocytes.