occurs in 20%–40% of adult patients harbouring the common m.3243A>G mutation, usually with a hypertrophic phenotype. Pathogenetic mechanisms are unclear yet no detailed study of myocardial structure, function and bioenergetics has been performed in m.3243A>G mutation carriers to identify early markers of cardiac involvement.

Methods Cardiac MRI was performed in 20 adult patients (10 males, mean age 38.7 ± 13.1 years) harbouring the m.3243A>G mutation, without clinical evidence of cardiac involvement on routine ECG and echocardiographic screening, and 20 age- and gender-matched healthy controls (10 males, 38.4 ± 14.2 years): (1) phosphorus-31 magnetic resonance spectroscopy, (2) cine imaging (3), cardiac tagging, and (4) late gadolinium enhancement (LGE) imaging on a Philips Intera Achieva 3 Tesla scanner. Clinical disease burden was determined using: (1) the Newcastle Mitochondrial Disease Adult Scale (NMDAS), a 29-domain validated scoring system with a single cardiac domain, and (2) urinary mutation load, the best predictor of overall clinical outcome in m.3243A>G mutation carriers.

Results Compared to control subjects and following Bonferroni correction for multiple comparisons, patients had increased left ventricular mass index (LVMI), LV mass to end-diastolic volume ratio (M/V ratio), LV radial wall thicknesses (all $p < 0.01$), peak torsion and torsion to endocardial circumferential strain ratio (both $p < 0.05$). Longitudinal shortening was decreased in patients ($p < 0.0001$) and correlated with increased LVMI ($r = -0.59$, $p = 0.01$). These findings are consistent with a reduction in contractile function in the subendocardium compared to the subepicardium, and similar to those reported in familial hypertrophic cardiomyopathy, but there were no differences in diastolic function in our study. Nine patients had diabetes mellitus and 3 had treated hypertension. Among patients, there was no correlation of LVMI or M/V ratio with age, blood pressures, fasting blood glucose or HbA1C but urinary mutation load and NMDAS score correlated strongly with LVMI ($r = 0.72$, $p < 0.001$ and $r = 0.85$, $p < 0.0001$ respectively). Phosphocreatine (PCr)/adenosine triphosphate (ATP) ratio was significantly decreased ($p = 0.003$) in patients (1.57 ± 0.34) compared to controls (1.97 ± 0.22) but there were no associations between PCr/ATP ratio and markers of myocardial function, or mass. No patients displayed evidence of focal myocardial fibrosis on LGE imaging.

Conclusions Concentric remodelling and subendocardial dysfunction are prevalent in m.3243A>G mutation carriers without clinical cardiac disease. Assessment of myocardial deformation may be useful in monitoring disease progression or response to early intervention. Patients with higher urinary mutation loads and disease burden may be at increased risk of cardiac involvement.

072 FRAGMENTED QRS: A PREDICTOR OF MYOCARDIAL SCAR AND FIBROSIS IN HYPERTROPHIC CARDIOMYOPATHY

doi:10.1136/heartjnl-2012-301877b.72

N Sheikh,* M Papadakis, R Bastiaenen, L Millar, N Emmanuel, S Ghani, A Zaidi, S Gati, N Chandra, E Behr, S Sharma. *St. George's University of London, London, UK*

Background It is well-established that fragmented QRS complexes (fQRS) on the 12-lead ECG are a predictor of delayed gadolinium enhancement (DGE) on Cardiac MRI (CMR) and indicate myocardial scar or fibrosis in patients with coronary artery disease and dilated cardiomyopathy. Moreover, fQRS appear to correlate well with arrhythmic events and mortality in these cohorts. However the significance of fQRS in hypertrophic cardiomyopathy (HCM) is yet to be established. We sought to determine whether the presence of fQRS is a predictor of delayed gadolinium enhancement (DGE) on CMR in patients with HCM.

Methods The 12-lead ECGs of 82 consecutive patients with HCM who underwent CMR with gadolinium were analysed for the

presence of fQRS by two independent readers blinded to the CMR findings. Patients with documented myocardial infarction ($n=3$) were excluded from further analysis. The ECGs were correlated to CMR findings, and patients separated into DGE positive (DGE+ve; $n=44$) and negative (DGE-ve; $n=35$) groups. ECG territories of fQRS were correlated with myocardial segments of DGE on CMR, in order to determine whether areas of fQRS predicted areas of DGE.

Results Patients from the DGE+ve and DGE-ve groups were of similar gender (75% vs 77% male respectively, $p=1.00$) and age (54 ± 19 vs 57 ± 11 years respectively, $p=0.41$). Fragmented QRS complexes were significantly more prevalent in the DGE+ve group than in the DGE-ve group (68.2% vs 14.3%, $p < 0.001$). The positive predictive value (PPV) of fQRS for DGE on CMR was 85.7%, with a specificity of 85.7%, sensitivity of 68.2% and negative predictive value of 68.2%. In the DGE+ve group with fQRS ($n=30$), fQRS ECG lead territory was predictive of regions of DGE on CMRI in 73.3% ($n=22$) of patients.

Conclusions The presence of fQRS on 12-lead ECG correlates with DGE on CMR in patients with HCM, with good specificity and PPV. Electrocardiographic territories containing fragmentation also correlate with myocardial segments of DGE on CMR. This simple, inexpensive method may therefore be valuable for predicting scar or fibrosis in patients with HCM. Future work should focus on correlating fQRS with risk factors and events to determine its use in risk stratification.

073 EVALUATION OF CLINICAL MARKERS OF EARLY DISEASE EXPRESSION AND THE ABILITY TO PREDICT GENOTYPE IN FAMILIES WITH HCM AND MUTATIONS IN CARDIAC MYOSIN BINDING PROTEIN C

doi:10.1136/heartjnl-2012-301877b.73

¹S P Page,* ¹S Kounas, ¹P Syrris, ²M I Christiansen, ²F Rune-Hansen, ²P S Andersen, ¹P M Elliott, ¹W J McKenna. ¹The Heart Hospital, UCLH, UK; ²Statens Serum Institut, Denmark

Introduction Familial evaluation for the presence of left ventricular hypertrophy (LVH) is an important part of management in hypertrophic cardiomyopathy. However due to incomplete penetrance, the presence of LVH does not reliably identify all mutation carriers, with consequences for cascade screening. Previous studies in small genotyped populations have suggested that reduced myocardial tissue Doppler velocities in unaffected relatives may predate the development of LVH and may therefore be useful in identifying at risk relatives. The value of such techniques in a large cohort of genotyped relatives remains unknown however. This study sought to prospectively evaluate ECG and Echo markers of early disease expression in a large genotyped cohort of families with mutations in myosin binding protein C (MYBPC3).

Methods Relatives of index cases with HCM related to mutations in MYBPC3 (4 insertion/deletion, 7 missense, 4 nonsense, 5 intronic, 2 double heterozygotes) were evaluated. Clinical examination, ECG and transthoracic Echo (operator blinded to genetic status) were performed and combined with genetic predictive testing. The clinical value of ECG and Echo derived indices in predicting genotype were assessed.

Results Of 95 relatives, 40 did not carry the family mutation (Group 1, 22 males, 38.5 ± 16.7 years), 39 were unaffected mutation carriers (Group 2, 17 males, 37.4 ± 16.8 years) and 16 were clinically affected and excluded from the study. ECG evidence of left atrial enlargement (21 vs 3%, $p=0.01$) and non-pathological Q waves (64 vs 28%, $p=0.001$) were more common in group 2 than group 1. S wave amplitude in lead V2 was greater in group 2 than group 1 (16.6 ± 7.9 vs 12.4 ± 8.2 mV, $p=0.02$) and also lead V3 (12.7 ± 6.9 vs 9.3 ± 6.1 mV, $p=0.02$). ECG criteria for left ventricular hypertrophy were specific

but not sensitive for identifying mutation carriers (Abstract 073 table 1). Lateral annular late diastolic velocities were increased in group 2 but all other tissue Doppler indices were similar between the two groups (Abstract 073 table 2). The proposed cut-off values (from previous studies) were tested in our larger study and found to be clinically unhelpful; lateral Ea velocity <14 (sensitivity 40%, specificity 50%), lateral Sa velocity <13 (sensitivity 50%, specificity 50%), septal Sa <12 (sensitivity 100%, specificity 0%) and septal Ea <13 (sensitivity 70%, specificity 30%).

Abstract 073 Table 1

	Group 1 (mutation negative)	Group 2 (unaffected mutation carriers)	p Value
Sa lateral annulus (cm/s)	9.8±2.6	10.1±2.4	0.60
Ea lateral annulus (cm/s)	13.4±4.9	13.3±3.9	0.97
Aa lateral annulus (cm/s)	7.5±1.9	8.9±2.3	0.005
E/Ea ratio lateral annulus	6.0±1.8	5.8±1.2	0.62
Isovolumic relaxation time (lateral annulus, ms)	75.8±17.5	81.3±16.3	0.17
Sa septum (cm/s)	8.1±1.4	7.9±1.6	0.72
Ea septum (cm/s)	10.1±2.7	9.0±2.8	0.33
Aa septum (cm/s)	8.1±2.1	7.3±1.3	0.25
Isovolumic relaxation time (septum, ms)	92.0±21.1	92.5±13.8	0.95

Abstract 073 Table 2

	Sensitivity	Specificity
Sokolow-Lyon (%)	3	95
Cornell (%)	18	95
Romhilt-Estes ≥ 4 (%)	8	95
Abnormal T wave inversion (%)	21	93

Conclusions In previous small studies, tissue Doppler indices have been reported to be both sensitive and specific for identifying unaffected mutation carriers. However in our larger study, tissue Doppler imaging was unhelpful in predicting genotype and with current imaging techniques it is not possible to reliably identify unaffected mutation carriers before the development of ventricular hypertrophy.

074 THE PREVALENCE AND SIGNIFICANCE OF LEFT VENTRICULAR HYPERTRABECULATION IN HIGHLY TRAINED ATHLETES

doi:10.1136/heartjnl-2012-301877b.74

¹S Gati, ¹N Chandra, ¹R L Bennett, ¹M Reed, ²K Gaele, ¹M Papadakis, ¹V F Panoulas, ³L Chen, ²F Carrie, ¹S Sharma. ¹St. George's, University of London, London, UK; ²French Institute of Health and Medical Research, Paris, France; ³University Hospital Lewisham, London, UK

Introduction Left ventricular non-compaction (LVNC) cardiomyopathy is characterised by increased myocardial trabeculation and recesses. Clinical manifestations of the disorder include progressive left ventricular dilatation, systolic impairment, predilection to fatal arrhythmias and thrombo-embolic events. Studies in heart failure patients demonstrate a high prevalence (up to 30%) of myocardial

trabeculations irrespective of the criterion used, and raise the potential diagnosis of LVNC. Given the high prevalence compared with other primary cardiomyopathies, it is unclear whether the myocardial morphology is representative of LVNC or merely an epiphenomenon associated with increased cardiac preload. The large cardiac preload associated with regular participation of intensive exercise results in physiological cardiac remodelling including increased left ventricular wall thickness and cavity size. Isolated case findings have also revealed increased trabeculations in some athletes but the significance of the anomaly is unclear. The distinction between cardiac remodelling from athletic training and LVNC is important when one considers that primary cardiomyopathies are the most commonly implicated cause of exercise related sudden cardiac death in young athletes. The aim of this study was to identify the prevalence and significance of hypertrabeculation in highly trained young athletes.

Method Between 2003 and 2011, 1146 athletes, aged 14–35 years, underwent 12-lead ECG and echocardiography. Echocardiograms were analysed in accordance with ASE guidelines and hypertrabeculation was defined as >3 localised protrusions of the ventricular wall >3 mm in thickness associated with intertrabecular recesses. Results were compared with 415 healthy controls of similar age.

Results Athletes displayed a higher prevalence of LV HTC compared with controls (18.3% vs 9.0%; $p<0.0001$). Of the athletes, 10.1% fulfilled conventional criteria for LVNC. African/Afro-Caribbean athletes exhibited a higher prevalence of LV HTC compared with Caucasians (28.8% vs 16.3%; $p=0.002$). Left ventricular hypertrabeculation was associated with T-wave inversion and lower indices of systolic function, however, assessment with 48 h ECG, exercise stress test and cardiac MRI failed to identify broader features of LVNC phenotype. Follow-up during the ensuing 48.6 ± 14.6 months did not reveal adverse events.

Conclusion The high prevalence of LV HTC in athletes, particularly among African/Afro-Caribbean athletes, suggests that the morphological anomaly represents an ethnically determined physiological epiphenomenon secondary to increased cardiac preload and afterload. Associated marked repolarisation changes and lower LV fractional shortening cannot exclude a myocardial disorder in a small minority. Prolonged longitudinal follow-up in a larger cohort of athletes should identify the precise significance of LV HTC.

075 EFFECT OF AORTIC VALVE REPLACEMENT ON LV REMODELLING AND MATRIX METALLOPROTEINASES AND THEIR TISSUE INHIBITORS IN ISOLATED SEVERE AORTIC STENOSIS

doi:10.1136/heartjnl-2012-301877b.75

C D Steadman, ^{*}L L Ng, B Grundy, I B Squire, N J Samani, G P McCann. University of Leicester, Leicester, UK

Introduction Surveillance of valvular heart disease accounts for a significant proportion of outpatient attendances in cardiology. A biomarker that correlates with disease severity and helps guide the optimum timing for intervention would be of great interest. NT-proBNP has been extensively examined in cardiac disease but focus has turned to enzymes that control collagen turnover, specifically matrix metalloproteinases (MMPs) and their tissue inhibitors (TIMPs). There is little data, other than from tissue samples, in aortic stenosis. We sought to examine the relationship between LV geometry and remodelling and serum MMPs and TIMPs in isolated severe aortic stenosis.

Methods 46 patients with isolated severe AS (peak velocity >4 m/s or mean pressure gradient >40 mm Hg or aortic valve area <1 cm²) without obstructive coronary artery disease (on angiography) were