GW23-e2598

CLINICAL ANALYSIS OF 64 CASES OF ATRIAL MYXOMA SURGERY UNDER SUPERFICIAL HYPOTHERMIA AND EXTRACORPOREAL CIRCULATION WITHOUT HEART BEATING CEASE

doi:10.1136/heartjnl-2012-302920a.284

¹Ding Kai, ²Zhou Hua-fu, ²Zhen Bao-shi, ¹Ding Kai. ¹*Department of Cardiothoracic Surgery, Wanbei Coal-electricity Group General Hospital, SuZhou;* ²*Department of Cardiothoracic Surgery, The First Affiliated Hospital of Guangxi Medical University, Nanning*

Objectives To investigate means and effect on surgical treatment of atrial myxoma under superficial hypothermia and cardiopulmonary bypass (CPB) without heart beating cease.

Methods Clinical datas and surgical treatment of 64 patients with primary atrial myxoma from January 2003 to March 2012 were retrospectively reviewed. All of atrial myxoma excised under general anaesthesia with superficial low temperature extracorporeal circulation on beating heart.

Results The CPB time was 19–84 min (mean 44.6 ± 15.2 min). All of 53 cases of left atrial myxoma and 7 cases of right atrial myxoma and four cases of double atrial myxoma were respected successfully and without cerebral embolism or gas embolism etc postoperative complications occurred, and without death in period of operation, they were all myxoma recovered. The patients were followed up for 1 month–9 years with a follow-up rate of 85.9%. All of cases didn't relapsed after operation nine patients were lost to follow-up.

Conclusions Atrial myxoma should be early performed after diagnosis results suggested that intracardiac procedures on pump beating-heart with superficial hypothermia cardiopulmonary bypass is safe and available in patients with atrial myxoma.