post-procedural troponin T elevation. According to the results of multivariate analysis, we found that age, total cholesterol, number of stenosed vessels and number of implanted stents were independent risk factors of post-procedural troponin T elevation.

Conclusions Age, serum total cholesterol, number of stenosed vessels and number of implanted stents could be independent risk factors of troponin T elevation after elective DES implantation.

GW23-e0061

RISK FACTORS OF CARDIAC TROPONIN T ELEVATION IN PATIENTS WITH STABLE CORONARY ARTERY DISEASE AFTER ELECTIVE CORONARY DRUG-ELUTING STENT IMPLANTATION

doi:10.1136/heartjnl-2012-302920e.3

陈章伟, 钱菊英, 马剑英, 葛均波, 葛均波. 复旦大学附属中山医院

Objectives Cardiac troponin T elevation after coronary intervention has been demonstrated to be associated with the prognosis of coronary artery disease (CAD). However, there were few studies about comprehensive risk factors analysis of troponin T elevation after elective drug-eluting stent (DES) implantation.

Methods From March to December in 2010, patients with stable CAD were admitted for elective coronary intervention in our hospital. They were divided into elevated troponin T group and normal troponin T group by post-procedural troponin T. Clinical factors, laboratory-test factors and angiographic factors (such as gender, age, cholesterol, Gensini score and so on) were analysed.

Results A total of 209 patients with an average age of 64.0 ± 9.9 years were enrolled in the study: 70 patients with elevated troponin T (≥ 0.03 ng/ml) after DES implantation and 139 patients with normal troponin T (< 0.03 ng/ml). After univariate analysis, we found that age, hypertension, total cholesterol, LDL-C, Gensini score, number of stenosed vessels and total implanted stents were associated with