GW23-e1096

THE RELEVANCE AMONG SERUM CRP LEVEL, CRP GENE C+1444T POLYMORPHISM AND THE RISK WITH ACUTE MYOCARDIAL INFARCTION

doi:10.1136/heartjnl-2012-302920k.20

He Guo-Ping, Yang Hai-Tao, Qi Chuan-Ping, Xu Lian-Hong, Qian Yi-Chao, He Guo-Ping.
Affiliated Wujin Hospital of Jiangsu University, Changzhou 213002, China

Objectives To investigate the relationship between serum CRP level, CRP gene C+1444T polymorphism and the risk with AMI from Sunan Chinese population.

Methods The serum CRP level was measured by enzyme linked immunosorbent assay (ELISA) and the CRP gene C+1444T polymorphism was genotyped by Polymerase reaction restriction-fragment length polymorphism (PCR-RFLP) between 227 patients with AMI (AMI group) and 161 control subjects.

Results

1. No differences were found in genotype distribution between AMI group and controls (CC 82.38%, CT 17.62%, TT 0 vs 86.96%, 13.04%, 0) ($p>0.05$).
2. The serum CRP level in AMI group was significantly higher than controls ($p<0.01$).
3. There were no differences in the serum levels between any genotypes of the CRP gene C+1444T ($p<0.05$).

Conclusions The CRP gene C+1444T polymorphism was not associated with increased risk of AMI, and it did not yet effect the serum levels in Chinese Han population of Sunan region.