GW23-e2389

EFFECT OF VALSARTAN ON EXTRACELLULAR MATRIX REMODELLING IN RATS WITH HEART FAILURE AFTER MYOCARDIAL INFARCTION

doi:10.1136/heartjnl-2012-302920o.19

Zhao Juan, Tian Ye. *The First Clinical Hospital of Harbin Medical University*

Objectives To investigate the effect of Valsartan on the change of Disintegrin metalloproteinases (ADAMs10,17) expressions in the heart failure (HF) process after myocardial infarction (MI).

Methods Based on UCG (LVEF<45%) results, the successfully MI-operated Wistar rats were divided into three groups: HF group, placebo group and Val group, which were given Valsartan by gavage. After 16 weeks, all rats were assessed by hemodynamic evaluation and serum TNF- α from LV was measured by ELISA (R & D, USA). In addition, their left ventricular (LV) muscle samples were extracted from the ischaemic segments, and then the ADAMs 10,17 expression were measured by immunoblotting.

Results In this study, heart failure (LVdp/dtmax, LVdp/dtmin and LVSP) was significantly elevated in Val group than the others ($p=0.006$, $p=0.015$, $p=0.003$), and the LVEDP level was decreased ($p=0.002$). At the same time, the TNF- α level in Val group was lower than HF groups ($p=0.023$). The ADAM17 and TNF-R₁ expressions in Val group was lower compared with those in HF group ($p=0.011$, $p=0.022$). However, ADAM10 expression is unchangeable in the four groups.

Conclusions Valsartan may reduce the ADAM17 and TNF-R₁ expression in the ischaemic zone of myocardium, decrease the TNF- α concentration in LV so as to inhibit cardiac remodelling and improve the heart function after MI.