Methods The levels of serum cardiac troponin I (cTnI) of 174 patients with chronic heart failure (CHF) were detected. The patients were divided into two groups: cTnI -positive group with serum cTnI ≥ 0.16 ng/ml (n=78) and cTnI-negative group with serum cTnI < 0.16 ng/ml (n=96). Ventricular arrhythmia was assessed by 24 h Holter monitoring in two groups. The incidence of adverse cardiac events, re-hospitalisation mortality during the period of in-hospital and the 6 months of out-of-hospital were observed and followed up.

Results Mean hourly number of single ventricular premature beats, Mean hourly ventricular pairs, and the frequency of ventricular tachycardia episodes per 24 h in cTnI-positive group were significantly higher than those in cTnI-negative group (all $p < 0.01$); during the period of observation and follow-up, the incidence of adverse cardiac events, re-hospitalisation and mortality of patients in cTnI-positive group were significantly higher than those in cTnI-negative group (all $p < 0.05$).

Conclusions Serum cTnI levels correlated with ventricular arrhythmia and late prognosis, which could be used as a prognosis predicting for patients with CHF.

GW23-e1629

CORRELATION OF CARDIAC TROPONIN I WITH VENTRICULAR ARRHYTHMIA AND LATE PROGNOSIS IN THE PATIENTS WITH CHRONIC HEART FAILURE

doi:10.1136/heartjnl-2012-302920v.9

Ma Cai-yun, Ren Feng-xue, Li Yuan-zheng, Ma Cai-yun. *Aviation General Hospital*

Objectives TO investigate correlation of cardiac troponin I (cTnI) with ventricular arrhythmia and late prognosis in the patients with chronic heart failure (CHF).