Results The peak I_{Na} densities did not differ between WT and mutant channels containing R800L-SCN5A, A261V-SNTA1 or R800L-SCN5A plus A261V-SNTA1. However, late I_{Na} for either mutant channel was moderately increased 2–3 fold compared to WT. The combined mutations of R800L-SCN5A plus A261V-SNTA1 significantly enhanced the I_{Na} late/peak ratio by 5.6-fold compared with WT. The time constants of current decay of combined mutant channel were markedly increased. The ‘gain-of-function’ effect could be blocked by the NG-monomethyl-L-arginine (L-NMMA), a nNOS inhibitor.

Conclusions We conclude that novel mutations in SCN5A and SNTA1 synergistically exert a nNOS dependent ‘gain-of-function’ on SCN5A channels, which may consequently prolong the action potential duration (APD) and lead to LQTS phenotype.

GW23-e1704

SYNERGISTIC EFFECT OF THE NOVEL MUTATIONS IN SCN5A AND SNTA1 ON LATE I_{Na} CONTRIBUTING TO LQT SYNDROME

doi:10.1136/heartjnl-2012-302920a.211

Rou-Mu Hu, Jieli Pu. Center for Arrhythmia Diagnosis and Treatment, Fu Wai Cardiovascular Hospital, Peking Union Medical College and Chinese Academy of Medical Sciences

Objectives SCN5A and SNTA1 are reported susceptible genes for long QT syndrome (LQTS). This study was designed to elucidate a plausible pathogenic arrhythmia mechanism for the combined novel mutations R800L-SCN5A and A261V-SNTA1 on cardiac sodium channels.

Methods A Caucasian family with syncope and marginally prolonged QT interval was screened for LQTS-susceptibility genes and found to harbour the R800L mutation in SCN5A and A261V mutation in SNTA1. The mutations were engineered into the most common splice variant of human SCN5A and SNTA1 cDNA respectively and sodium current (I_{Na}) was characterised in HEK293 cells co-transfected with neuronal nitric oxide synthase (nNOS) and the cardiac isoform of the plasma membrane Ca-ATPase (PMCA4b).