B

CHROMOSOME 9p21 LOCUS AND CORONARY ARTERY DISEASE – COLLABORATIVE META-ANALYSIS ON ANGIOGRAPHIC BURDEN AND MOLECULAR FUNCTION ANALYSIS

K Chan,¹ A Motterle,¹ R S Patel,^{1,2,3} X Pu,¹ S Ye¹ ¹William Harvey Research Institute, Barts and the London School of Medicine and Dentistry, Queen Mary University London, London, UK; ²Department of Medicine, Cardiff University, Cardiff, Wales, UK; ³Department of Medicine, Emory University School of Medicine, Atlanta, GA, USA

doi:10.1136/heartjnl-2013-304019.275

Background Chromosome 9p21 variants showed the strongest association with coronary heart disease in genome-wide association studies. But questions remain on the mechanism underlying the risk rendered by this locus, and clinically whether the locus contributes to coronary atheroma burden or plaque instability. We investigated the relationship of 9p21 locus with (1) angiographic coronary artery disease (CAD) burden and progression to myocardial infarction (MI); and (2) biological function of vascular smooth muscle cell (VSMC).

Methods and results We established a collaboration of 21 studies consisting of 33,673 patients with information on both CAD (clinical or angiographic) and MI status along with 9p21 genotype. We first confirmed an association between 9p21 and CAD using angiographically defined cases and controls (pooled odds ratio (OR)=1.31 (95% CI 1.20–1.43)). Among subjects with angiographic CAD (n=20,987), random-effects model identified an association with multi-vessel CAD, compared to those with single-vessel disease (OR=1.10 (95% CI 1.04–1.17) per copy of risk allele). However, there was no significant association between 9p21 and prevalent MI when both cases (n=17,791) and controls (n=15,882) had underlying CAD (OR=0.99 (95% CI 0.95–1.03) per risk allele). Immunohistochemical staining of human atherosclerotic plaque showed co-localization of VSMC with the cell-cycle regulator proteins p16^{INK4a}, p14^{ARF} and p15^{INK4b}, which are encoded by the genes *CDKN2A* and *CDKN2B* genomically located nearby the 9p21 locus. The 9p21 risk genotype confers reduced p15^{INK4b} levels ($p=3.7 \times 10^{-2}$) and higher VSMC content ($p=5.6 \times 10^{-4}$) in the plaques. We further examined the influence of 9p21 genotype on primary cultures of VSMC isolated from human umbilical cord. The risk genotype was associated with reduced expression of p16^{INK4a}, p15^{INK4b} ($p=1.2 \times 10^{-5}$, 1.4×10^{-2}), and increased VSMC proliferation ($p=1.6 \times 10^{-2}$).

Conclusions The 9p21 locus shows convincing association with greater burden of CAD, but not with MI in the presence of underlying CAD, indicating that the 9p21 locus primarily mediates an atherosclerotic phenotype. This is supported by the observations from atherosclerotic plaques and primary cell cultures indicating its influence on VSMC proliferation, a process featured in atherosclerotic pathogenesis.