phenomenon in asymptomatic white individuals without a relevant family history. Conversely, ATWI beyond V2 is rare, particularly in males and may warrant investigation. The results of this study will have a significant impact on ECG interpretation in young white athletes and non-athletes.

BCS Affiliates — Allied Professionals (BACPR/BANCC)

150 THE INCIDENCE OF LONG QTc IN ACUTE MEDICAL ADMISSIONS: AN EVALUATION OF FIVE METHODS

¹Rahel Mahmud*, ²Adam Grey, ²Adam Nabeebaccus, ²Martin Whyte. ¹Kings College London; ²Kings College Hospital; *Presenting Author

10.1136/heartjnl-2016-309890.150

Aim Long QTc is associated with poor outcomes on CCU and acute stroke units. Its incidence and outcome in unselected medical admissions is unknown. A range of drug classes and electrolyte abnormalities may lead to acquired long QTc. Given the frequency with which these are encountered in medical admissions, the incidence of long QTc requires study. We have calculated the incidence of long QTc encountered in acute medical admissions. As inaccuracies in the correction formula may give misleading data, this study compared five formulas commonly used to correct QT interval duration by heart rate (HR): Bazett (B), Fridericia (Fa), Framingham (Fm), Hodges (H), Van der Water's (VdW) as well as the automated QTc correction provided by the Welch Allyn (WA) ECG recorder.

Methods 1000 records of consecutive patients admitted to acute medical wards at a tertiary teaching hospital were reviewed by two physicians. Admission ECGs were examined. QTc was calculated using the five formulas specified. The Pearson correlation coefficient (R^2) was used to estimate the relationship between QTc and HR. Patient age and gender were recorded. QTc of ≥ 500 ms was considered significantly prolonged in both genders. ECGs with missing patient identifiers or poor trace quality were excluded. Cohen's kappa statistic, for inter-observer agreement, was made from a subgroup of 200 randomly chosen ECGs evaluated by a third observer.

Results 715 ECGs were included. 356 males, 359 females. Age 61 ± 20 years (mean $\pm$ SD). Kappa inter-observer agreement was 0.65 indicating substantial agreement. Incidence of QTc ≥ 500 ms was 1.7% (B), 0.4% (Fa), 0.4% (Fm), 0.3% (H), 0.4% (VdW) but was 6.2% with WA. Pearson correlation between QTc and HR was 0.02 (B), 0.06 (Fa), 0.5 (Fm), 0.001 (H), 0.5 (VdW) 0.009 (WA).

Conclusion The incidence of significant QT prolongation in unselected medical admissions varied by a factor of ten depending on the method used. Automated QTc software gave a far higher incidence of long QTc than manual interpretation. Framingham and Van der Water's equations show bias by HR. The clinical sequelae of long QT in unselected medical admissions needs further study.

Valve Disease/Pericardial Disease/ Cardiomyopathy

151 CARDIAC MYOSIN-BINDING PROTEIN C IS A NOVEL MARKER OF MYOCARDIAL INJURY AND FIBROSIS IN PATIENTS WITH AORTIC STENOSIS

¹Atul Anand*, ²Calvin Chin, ²Anoop Shah, ³Jacek Kwietniński, ²Alex Vesey, ⁴Joanna Cowell, ⁵Ekkehard Weber, ⁶Thomas Kaier, ⁷David Newby, ⁷Marc Dweck, ⁶Michael Marber, ⁷Nicholas Mills. ¹University of Edinburgh, BHF Centre for Cardiovascular Science; ²University of Edinburgh; ³University of Edinburgh and Poznan University of Medical Sciences; ⁴Department of Geriatric Medicine, NHS Lothian; ⁵Institute of Physiological Chemistry, Martin Luther University Halle-Wittenberg; ⁶King's College London BHF Centre; ⁷BHF Centre for Cardiovascular Science, University of Edinburgh; *Presenting Author

10.1136/heartjnl-2016-309890.151

Background Cardiac myosin binding protein C (cMyC) is an abundant sarcomeric protein and a novel highly specific marker of myocardial injury. Given myocyte death characterises the transition from hypertrophy to replacement myocardial fibrosis in advanced aortic stenosis, we hypothesised that serum cMyC concentrations would be associated with cardiac structure and outcomes in patients with aortic stenosis.

Methods cMyC was measured in two cohorts: a mechanism cohort of patients with aortic stenosis ($n = 161$) and healthy controls ($n = 46$) who underwent cardiac magnetic resonance imaging, and an outcomes cohort with aortic stenosis ($n = 104$) followed for a median of 11.3 years. True cut myocardial biopsies were obtained from 10 patients in the mechanism

Abstract 151 Figure 1 Linear regression correlation of cMyC with indexed LV mass and fibrosis volume in aortic stenosis