LEFT VENTRICULAR LONGITUDINAL STRAIN-VOLUME RELATIONSHIPS IN ELITE ATHLETES

¹David Oxborough, ²Annemieke Heemels, ¹Lynsey Forsythe*, ³Gavin McClean, ¹Punit Mistry, ¹Rachel Lord, ¹Victor Utomi, ⁴Nigel Jones, ²Dick Thijssen, ⁵John Somauroo, ⁶Sanjay Sharma, ⁷Rebecca Osborne, ⁸Nicholas Sculthorpe. ¹Liverpool John Moores University; ²Radboud University; ³Aspetar Orthopaedic and Sports Medicine Hospital; ⁴Aintree University Hospitals NHS Trust; ⁵Countess of Chester NHS Foundation Trust; ⁶St Georges University; ⁷Cardiac Risk in the Young; ⁸University of West UK; *Presenting Author

10.1136/heartjnl-2016-309890.120

Introduction It is well established that left ventricular (LV) adaption occurs in response to chronic physiological conditioning. There is also evidence highlighting functional differences in myocardial strain imaging between athletes from sporting disciplines. This difference may be a consequence of the vague classification of sport i.e. not taking into account relative static and dynamic components and/or merely a consequence of chamber enlargement. We sought to utilise a novel *simultaneous* assessment of longitudinal strain and LV volume in athletes classified in the 4 corners of Mitchell's classification of sporting disciplines. The primary aim was to determine relative longitudinal strain throughout the cardiac cycle and its specific contribution to LV volume change in these athletes.

Methods 92 elite male athletes were studied and sub classified based on sporting discipline in accordance with the Mitchell's classification. (Group IA *low static-low dynamic* n = 20, Group IC *low static-high dynamic* n = 25, Group IIIA *high static-low dynamic* n = 21, Group IIIC *high static-high dynamic* n = 26). Conventional echocardiography of the LV was undertaken. The raw temporal global longitudinal strain values were exported and divided into 5% time increments across the cardiac cycle. Concomitant LV volumes were traced at each 5% time increment to provide simultaneous strain-volume loops. The strain-volume relationship was assessed by applying a polynomial regression analysis for each systolic and diastolic curve to derive absolute values for% end diastolic volumes (EDV).

Results Conventional and peak strain indices are presented in table 1. Athletes in group IC and IIIC had larger LV end diastolic volumes (EDV) compared to athletes in groups IA and IIIA (50 ± 6 and 54 ± 8 ml/(m²)^{1.5} vs. 42 ± 7 and 43 ± 2 ml/(m²)^{1.5} respectively). Group IIIC also had significantly larger mean wall thickness (MWT) compared to all groups. Peak strain was variable between groups but once normalised for EDV all groups, with exception of IIIC, required similar strain to generate the same% reduction in EDV (see Figures 1 and 2). Conversely group IIIC required greater longitudinal strain for any given% volume which correlated to MWT ($r = 0.4$, $p < 0.0001$).

Abstract 120 Table 1 Echocardiographic Parameters

PARAMETER	GROUP IA	GROUP IC	GROUP IIIA	GROUP IIIC
LVDd index (mm/(m ²) ^{0.5})	37 ± 3†	39 ± 3	37 ± 3†	40 ± 2*‡
LVEDV index (ml/(m ²) ^{1.5})	42 ± 7^†	50 ± 6*‡	43 ± 2^†	54 ± 8*‡
EF (%)	60 ± 7	58 ± 7	59 ± 5	59 ± 7
MWT index (mm/(m ²) ^{0.5})	6.0 ± 0.4†	6.3 ± 0.6	6.3 ± 0.6	6.7 ± 0.7*
MaxWT index (mm/(m ²) ^{0.5})	6.6 ± 0.7†	7.0 ± 0.7†	7.1 ± 0.7	7.6 ± 0.9*^
LV Mass Index (g/(m ²) ^{2.7})	33 ± 8†	37 ± 8	35 ± 9†	42 ± 9*‡
LVMass/LVEDV (g/ml)	1.4 ± 0.2	1.4 ± 0.3	1.4 ± 0.3	1.5 ± 0.3
Longitudinal Strain (%)	-20 ± 3^†	-16 ± 2*†	-18 ± 2*†	-20 ± 3^†

Symbol denotes P > 0.05 to IA=*, IC=^, IIIA=‡, IIIC=†

Abstract 120 Figure 1 Temporal Assessment of Simultaneous Strain and Volume

Conclusion There are physiological differences between athletes with the largest LV demonstrated in athletes from group IIIC. These athletes also have greater resting longitudinal contribution to volume change which, in part, is related to an increased wall thickness. The variance in peak strain seen in the other athlete groups was solely related to chamber size with no intrinsic differences in contractility or relaxation.

Abstract 120 Figure 2 Derived Strains for% EDV in the EF range 10 to 70%

Cardiac Rhythm Management

121 THE 12-LEAD ELECTROCARDIOGRAM OF THE ELITE RUGBY FOOTBALL LEAGUE PLAYER

¹Lynsey Forsythe*, ¹Keith George, ²John Somauroo, ¹Mohammad Qasem, ¹Benjamin Brown, ³Michael Papadakis, ¹David Oxborough. ¹Liverpool John Moores University; ²Countess of Chester Hospital; ³St Georges University of London; *Presenting Author

10.1136/heartjnl-2016-309890.121

Introduction Specific ECG criteria have been proposed to differentiate physiology (athlete's heart) from pathology that may increase the risk of sudden cardiac death. Although the current European Society of Cardiology (ESC) guidelines have clear criteria for physiological adaptation to intense training, their sensitivity and specificity are sub-optimal and hence the more recent Seattle and Refined criteria have attempted to improve diagnostic accuracy of the athlete's ECG.

Rugby Football League (RFL) is a moderate dynamic and moderate static sport with recent high profile fatalities linked to inherited cardiomyopathy. This study utilises resting 12-lead ECG, in elite RFL players to compare 3 sets of diagnostic ECG criteria in the athlete and specifically false positive rates.

Methods 103 consecutive, male, elite RFL players (mean age 25.4 years) underwent pre-participation cardiac screening. Participants were predominantly white Caucasian (n = 81) with a minority of ethnic backgrounds being represented (Pacific Islander, n = 16; mixed race, n = 5; African-Caribbean, n = 1). All athletes had resting blood pressure assessed and a standard resting 12-lead ECG. In addition *all* athletes had a standard transthoracic echocardiogram and where indicated further investigations were undertaken. Standard ECG parameters were measured and the ESC recommendations for interpretation of 12-lead ECG in the athlete were used to define normal training (Group 1) and abnormal (Group 2) changes. All ECG's were also assessed for normality using the Seattle and Refined criteria. False-positive rates were presented for each ECG criteria.

Results Based on ECG, echocardiography and/or follow-up investigation all athletes were considered normal with no evidence of underlying cardiac disease. The continuous ECG

Abstract 121 Table 2 % Training unrelated ECG changes for ESC, Seattle and Refined Criteria (% False Positive Rate)

ECG Abnormality	False Positive Rate (%)		
	ESC Criteria	Seattle Criteria	Refined Criteria
Long QTc Interval	1	0	0
Short QTc Interval	44	0	Not Relevant
Intraventricular conduction delay (IVCD)	5	0	1
Right atrial enlargement (RAE)	1	0	1
Right ventricular hypertrophy (RVH)	3	0	3
Right axis deviation (RAD)	1	0	1
Left axis deviation (LAD)	3	3	3
T wave Inversion	1	1	1
Y2 PVC's per 10s tracing	Not Relevant	1	1
False Positive Rate according to Criteria	58	5	3

measurements are shown in Table 1. According to ESC criteria 95% of athletes had at least one group 1 "normal training-related" ECG change. 58% of athletes had at least one Group 2 "abnormal training-related" change. This is therefore a false positive rate of 58%. Following application of the revised Seattle and Refined criteria the false-positive rates reduced to 5% and 3% respectively (Table 2).

Conclusion A significant proportion of RFL players present with an abnormal, non-training related, 12-lead ECG according to ESC criteria. In the cardiac screening setting this would result in significant extra cardiac investigation burden. Application of the Seattle and refined criteria decreased the false-positive rate and hence improves ECG specificity in this population.

Imaging

122 INTRODUCTION OF STRESS ONLY MYOCARDIAL PERFUSION SCINTIGRAPHY: A QUALITY IMPROVEMENT PROGRAMME

¹Eleanor Wicks*, ²Isabella Ceravolo, ³Rayjanah Allie, ³Anna Barnes, ²Perry Elliott, ²Leon Menezes. ¹University College London; ²UCL; ³UCLH; *Presenting Author

10.1136/heartjnl-2016-309890.122

Abstract 121 Table 1 Continuous ECG variables of 103 elite RFL players

Continuous ECG variables	Mean $\pm$ SD(Range)
Heart Rate (bpm)	54 $\pm$ 9 (39–89)
P Duration (ms)	105 $\pm$ 16 (58–176)
PR Interval (ms)	174 $\pm$ 26 (129–309)
QRS Duration (ms)	95 $\pm$ 12 (11–123)
QT Corrected (ms) (Bazett)	381 $\pm$ 20 (338–442)
QRS Axis (degrees)	61 $\pm$ 32 (-46–117)

Introduction Myocardial perfusion scintigraphy (MPS) is one of the most commonly performed non-invasive imaging procedures for the diagnosis of Coronary Artery Disease (CAD). Two large trials evaluating over 21,000 patients with a normal SPECT study have demonstrated the safety of stress-only imaging as compared to traditional stress/rest imaging. Comparably low all-cause and cardiac mortality rates were observed with both imaging protocols. This was true irrespective of patient age, gender, cardiac risk factor profile, or stressor employed with SPECT. These results are consistent with earlier studies evaluating patient outcome following normal stress-only imaging. The best use of a stress-only imaging strategy is likely to be in the selected low or intermediate risk population, which forms the referral basis for myocardial perfusion imaging. In this study, we describe the introduction of stress only MPS in comparison to traditional stress/rest MPS.