Abstract 011 Figure 1 A. Comparison of T1 reactivity profiles for 'Unaffected' myocardium versus regions with abnormal perfusion versus LGE; B. ROC curve depicting sensitivity and specificity of stress T1 reactivity to predict perfusion deficits in HCM.

with LGE and perfusion defects compared to the remaining 'unaffected' myocardium (Figure 1). Using a diagnostic threshold of 3.5% on ROC analysis for detecting impaired MBF in HCM, T1 reactivity had a sensitivity and specificity of 82% and 88% respectively (AUC 0.894, $p < 0.01$) (Figure 1). Adenosine induced rise in MBF, T1 reactivity, oxygenation and global longitudinal strain (GLS) were significantly reduced in HCM compared with controls ($p < 0.01$ for all comparisons). Stress T1 reactivity moderately correlated with MBF reactivity ($r = 0.55$, $p < 0.01$) and GLS ($r = -0.44$, $p < 0.01$).

Conclusions

1. Stress T1 mapping is capable of distinguishing regions with perfusion defects versus normal perfusion in HCM without the need of contrast.
2. Stress T1 reactivity, MBF reactivity, oxygenation and GLS were significantly reduced in HCM compared to controls.

REFERENCES

1. Liu A, Wijesurendra RS, Francis JM, et al. Adenosine Stress and Rest T1 Mapping Can Differentiate Between Ischaemic, Infarcted, Remote, and Normal Myocardium Without the Need for Gadolinium Contrast Agents. *JACC Cardiovasc Imaging* 2016;**9**(1):27–36.
2. Mahmod M, Piechnik SK, Levelt E, et al. Adenosine stress native T1 mapping in severe aortic stenosis: evidence for a role of the intravascular compartment on myocardial T1 values. *J Cardiovasc Magn Reson* 2014;**16**:92.
3. Jerosch-Herold M. Quantification of myocardial perfusion by cardiovascular magnetic resonance. *J Cardiovasc Magn Reson* 2010;**12**(1):57.
4. McCommis KS, Zhang H, Goldstein TA, et al. Myocardial Blood Volume Is Associated with Myocardial Oxygen Consumption: An Experimental Study with CMR in a Canine Model. *JACC Cardiovascular imaging* 2009;**2**(11):1313–1320.

012 LONGITUDINAL CARDIAC MAGNETIC RESONANCE ASSESSMENT OF DIFFUSE AND REPLACEMENT MYOCARDIAL FIBROSIS IN AORTIC STENOSIS

RJ Everett, CWL Chin, J Kwiecinski, WJ Jenkins, S Mirsadraee, A White, SK Prasad, S Semple, DE Newby, MR Dweck.

10.1136/heartjnl-2017-311399.12

Objective To investigate whether cardiac magnetic resonance (CMR) can assess the progression and regression of myocardial fibrosis in patients with aortic stenosis (AS).

Methods Sixty-three patients with aortic stenosis were followed up for 2 years with serial CMR and echocardiography: 28 patients (63 ± 13 years, 68% male, 50% asymptomatic severe AS) did not undergo intervention (*natural history cohort*) and 35 patients (67 ± 8 years, 73% male, 83% symptomatic severe AS) underwent aortic valve replacement (AVR) within the first year (*AVR cohort*). Replacement and diffuse myocardial fibrosis were assessed using the late gadolinium enhancement (LGE) and T1 mapping techniques respectively. Annualised change was calculated for all measures.

Results In the natural history cohort, left ventricular mass index (LVMI) increased over time ($6\% \pm 1\%$, $p < 0.0001$) and was accompanied by a fall in longitudinal systolic function ($-4\% \pm 2\%$, $p = 0.03$). The indexed extracellular volume (iECV, a measure of diffuse fibrosis) also increased over time ($7\% \pm 2\%$, $p < 0.0001$). However, no changes were observed in native T1, post-contrast T1 or extracellular volume (ECV) fraction on serial imaging. Mid-wall LGE was observed in nine patients (32%). Absolute LGE mass increased by 3.8 ± 0.8 g ($p < 0.0001$) amongst all the patients with baseline mid-wall LGE. Four patients with mid-wall LGE at baseline developed new areas of LGE in different myocardial segments. No patients without pre-existing mid-wall LGE developed new LGE during follow up.

The AVR cohort displayed a fall in LVMI following surgery ($-15\% \pm 2\%$, $p < 0.0001$) mirrored by a reduction in diffuse fibrosis (iECV, $-9\% \pm 2\%$, $p < 0.0001$). The ECV fraction however was observed to increase (5% [2, 11], $p < 0.0001$), with no change in native T1. Mid-wall LGE was present in 9 patients (26%). No patient went on to develop new LGE nor did existing LGE resolve in any patient and LGE mass did not change. Longitudinal systolic function increased following AVR ($16\% \pm 6\%$, $p = 0.01$).

Conclusion Changes in diffuse fibrosis can be tracked using iECV and increase with time in patients with AS alongside LV mass and replacement fibrosis. Following AVR there is a fall in LV mass driven predominantly by regression of cellular

Abstract 012 Figure 1 Main Figure: Serial cardiac magnetic resonance imaging in a patient with severe aortic stenosis.

hypertrophy with more gradual reduction in diffuse fibrosis burden but no change in replacement fibrosis.

Mid-wall LGE is present on baseline scan (white arrow) and new areas are seen on 1 year scan (red arrows). Following AVR there is no change in mid-wall LGE (white arrows). Annualised change in imaging measures is presented below in the natural history (black) and AVR (red) groups.

013

FREE-BREATHING MOCO LGE LEADS TO BETTER IMAGE QUALITY AND FASTER SCANNING TIMES IN CLINICAL PRACTICE

¹Ilaria Lobascio, ^{1,2}Gabriella Captur, ¹Veronica Culotta, ³Redha Boubertakh, ⁴Andras Eke, ⁵Hui Xue, ^{1,6}Charlotte Manisty, ⁵Peter Kellman, ^{1,6}James C Moon, ¹Barts Heart Centre, The Cardiovascular Magnetic Resonance Imaging Unit, St Bartholomew's Hospital, West Smithfield, London, UK; ²UCL Biological Mass Spectrometry Laboratory, Institute of Child Health and Great Ormond Street Hospital, London and NIHR University College London Hospitals Biomedical Research Centre, London W1T, UK; ³Cardiovascular Biomedical Research Unit, Barts and the London School of Medicine and Dentistry, Queen Mary University of London, UK; ⁴Department of Physiology, Semmelweis University, Budapest, Hungary and Diagnostic Radiology, Yale University, New Haven, CT, USA; ⁵National Heart, Lung, and Blood Institute, National Institutes of Health, DHHS, Bethesda, MD, USA; ⁶University College London UCL Institute of Cardiovascular Science, London WC1E 6BT, UK

10.1136/heartjnl-2017-311399.13

Objectives Late gadolinium enhancement (LGE) sequences have evolved. Free-breathing, motion-corrected LGE (MOCO-

LGE)^{1,2} has several potential advantages over breath held LGE (bh-LGE)³ including minimal user input for the LGE short axis (SA) stack and no need for breath-holds. We hypothesised that the use of MOCO-LGE would be faster, cheaper and easier for clinical scanning, increasing throughput.

Methods 200 consecutive clinical patients underwent bh-LGE or MOCO-LGE at 1.5T. Image quality (Figure 1), scan time, patient throughput (change-over time) and reader confidence were compared. LGE image quality was evaluated qualitatively (adaption of previously reported method)⁴ and quantitatively (assessing image texture heterogeneity using grayscale lacunarity, λ).

Results MOCO-LGE image quality was better than bh-LGE qualitatively (lower score better: 0.56 ± 1.2 vs 1.93 ± 0.83 , $p < 0.0001$) especially in clinically vulnerable patients eg. atrial fibrillation, poor breath-holding, low ejection fraction (0.59 vs 3.05 , $p = 0.0001$). Excellent image quality (score=0) was also more common (78% vs 27%, $p < 0.0001$). Quantitative image quality was superior with MOCO-LGE (lower score better: blood pool λ bh-LGE 0.38 ± 0.11 vs MOCO-LGE 0.28 ± 0.08 , $p < 0.0001$). MOCO-LGE led to greater diagnostic confidence (blinded review: basic analytic, retained diagnostic and "Omny" correction methods, respectively $p = 0.005$; $p = 0.018$, $p < 0.001$). Although patient change-over time did not differ significantly between scan protocols, total LGE imaging time was 1.6 times shorter with MOCO-LGE compared to bh-LGE (5.23 vs 8.84 minutes, $p < 0.0001$).