infarct size was observed when WT-MI hearts were compared to their PMCA1^{Ht} counterparts 1 week post-surgery. Interestingly, whilst both WT and PMCA1^{Ht} MI-treated mice showed a significant deterioration in cardiac function 1 week post-surgery there were differences seen in cardiac structure between the two groups. For example, echocardiography revealed that the WT-MI hearts were significantly more hypertrophic when compared to their PMCA1^{Ht} counterparts shown by a significant increase in the LV diameters alongside a significant difference in the normalised heart weight to tibia length ratios. ECG revealed significantly longer QT intervals and considerably more extra-systolic events among the WT-MI mice. Whilst histological analysis of cardiomyocyte size showed a significantly exacerbated hypertrophic response in the remote regions of WT-MI mice compared to PMCA1^{Ht}-MI mice.

Conclusion Heterozygous deletion of PMCA1 might serve a protective role in the heart following both moderate and severe MI. The protective mechanism most likely develops in the early post-operative phase, as an MI of similar extent is associated with higher mortality rates in WT mice. Future work will aim to elucidate the mechanism producing this phenotype mainly focusing on the potential role that PMCA1 might serve in the process.

141

HERITABILITY AND FAMILY-BASED GWAS ANALYSES TO DISCOVER NOVEL LIPIDOMIC BIOMARKERS OF CARDIOVASCULAR DISEASE

¹Kathryn McGurk*, ²Anna Nicolaou, ¹Bernard Keavney. ¹Division of Cardiovascular Sciences, University of Manchester; ²Division of Pharmacy and Optometry, University of Manchester

10.1136/heartjnl-2017-311726.140

Introduction Lipids have key roles in CVD but heritability studies focus on few species bar lipoproteins. Lipids play key roles in cell signalling, immunity, inflammation, vasodilation and cell death. Although not encoded, their activities are tightly linked to DNA-encoded entities (e.g. enzymes and other proteins) and those with a strong genetic influence (high heritability) may identify novel pathways in CVD. Purpose: Analysing eicosanoid, endocannabinoid and sphingolipid profiles in 250 British Caucasian families with GWAS data will identify particularly heritable lipid biomarkers for the discovery of causative genetic variation of CVD.

Methods An array of 79 eicosanoids and related species, 33 endocannabinoids and congeners, 63 ceramides and related species, from 204 plasma samples (31 families of 1+ individuals with hypertension) were extracted and analysed by liquid chromatography coupled to tandem mass spectrometry (LC-MS/MS). Quality control was accessed and estimates of heritability of each lipid species was estimated using QTDT software. The concentration of each lipid species were assessed for normal distribution, outliers, and adjusted for age and sex. The resulting lipid concentrations will be analysed using FaST-LMM software for associations with 557,124 SNPs.

Results 19 species of eicosanoids were identified (mean concentrations 19 pg/ml – 7600 pg/ml); the species at highest abundance in plasma (HODEs) are derivatives of the omega-6 fatty acid linoleic acid. 8 species passed quality control assessments and 3 species were estimated to be significantly heritable (9-OxoODE 24%, 12-HETE 44%, 12(13)-EPOME 78%). 15 congeners of endocannabinoids were identified (mean concentrations 20 pg/ml – 4000 pg/ml); the species at highest

abundance in plasma are glycerol derivatives. 9 species passed quality control assessments and all were estimated to be significantly heritable (27%–73%)%. 57 species of ceramides were identified (mean concentrations 0.01 pmol/ml – 190 pmol/ml); the species at highest abundance in plasma are NS ceramides involved in apoptosis. 36 species passed quality control assessments and all species were estimated to be significantly heritable (10%–63%)%.

Conclusions We demonstrate for the first time estimates of heritability for an array of bioactive lipids. Although found at concentrations million times lower than cholesterol, many of the heritability estimates are similar to plasma genetic biomarkers of CVD (ACE h²>45%). The association between heritable biomarkers, cardiovascular phenotypes and risk will be determined. Particularly heritable species and those with highly significant hits at GWAS will be profiled in the extended cohort (1400 samples, 250 families) and replication will take place in the UK biobank, or other. A subset of the most heritable species and those with hits at GWAS resulting from the full lipidomic profile of 3 lipidomic classes will be used to identify causal metabolic pathways, novel diagnostics and drug targets for CVD intervention.

142

INTRAVITAL IMAGING OF LEUKOCYTE, PLATELET AND STEM CELL TRAFFICKING IN VIVO IN THE CARDIAC MICROCIRCULATION FOLLOWING MYOCARDIAL ISCHAEMIA-REPERFUSION INJURY

Dean Kavanagh*, Adam Lokman, Neena Kalia. University of Birmingham Stem cells, Reperfusion injury, Imaging

10.1136/heartjnl-2017-311726.141

Background The clinical success of stem cell (SC) therapy for myocardial infarction is compromised by poor cardiac homing following systemic delivery. As such therapy may depend on beneficial paracrine effects, it is a further hindrance that little is known about the inflammatory response dynamics within myocardial microcirculation *in vivo*.

Methods 3D-printed stabilisers were bonded to the beating heart of anaesthetised (ketamine/xylazine) mice to enable confocal intravital imaging of ventricular microcirculation. PE-anti-Gr-1 and APC-anti-CD41 antibodies were injected to label neutrophils and platelets respectively with FITC-BSA enabling blood flow visualisation. In some mice, haematopoietic SCs (HSCs; HPC-7s) were introduced intra-arterially. IR injury was induced by 45 min (reversible) ligation of the LAD artery.

Results Neutrophil adhesion and platelet accumulation were both significantly ($p<0.001$) and rapidly increased in injured microvessels with platelet accumulation increasing with time. No difference in number or velocity of free-flowing neutrophils was observed. A significant ($p<0.05$) decrease in functional capillary density was also observed in injured hearts. Although HSC adhesion was not significantly enhanced following injury, a time-dependent increase in adhesion was observed in sham and injured hearts. No significant change in number or velocity of free-flowing HSCs was observed following injury. Interestingly, despite reduced capillary perfusion, approximately 10–20 HSCs were observed trafficking through the heart at each time point throughout reperfusion.

Discussion Intravital microscopy has allowed successful visualisation of the microvascular inflammatory response and HSC homing events in the beating mouse heart post-reperfusion.

Subsequent experiments will allow us to assess whether HSC administration can confer vasculoprotection.

143 THE ROLES AND MECHANISMS OF PORE-FORMING TOXINS IN CARDIAC INJURY

¹Guozheng Wang*, ²Yasir Alhamdi, ²Cheng-Hock Toh. ¹Sepsis Research Group, Institute of Infection and Global Health, University of Liverpool, Liverpool, L69 7ZB; ²University of Liverpool

10.1136/heartjnl-2017-311726.142

Cardiac muscles and conduction system are very sensitive to interruption of plasma membrane integrity due to uncontrollable ion diffusion. There are many types of toxins that could bind to cell membrane and form pores to allow ion diffusion cross the membrane. However, the effects and underlying mechanisms of those pore-forming toxins on cardiac functions have not been systematically investigated.

In this work, we investigated the effects of extracellular histones (endogenous) and pneumolysin (exogenous) on cardiomyocytes and underlying molecular mechanisms using cultured cardiomyocytes and mouse models. We found that both toxins could bind to cardiomyocyte membrane and caused large inward currents which altered membrane potentials. This led arrhythmia of cultured cardiomyocytes and mice infused with isolated toxins. In addition, circulating troponins, a marker of cardiac injury, were significantly elevated and the contractile performance of cardiomyocytes was also reduced.

Further investigation, we found that the uncontrollable calcium influx mainly increased diastolic intracellular calcium concentration and resulted in calcium transient reduction. However, the calcium-dependent protein kinase C, particularly PKC $\pm$, was significantly activated by calcium influx. Consequently, the phosphorylation of Troponin I was enhanced, which is known to correlate to contractile reduction. In addition, the phosphorylation levels of P38 and JNK kinases, and endoplasmic reticulum (ER) stress levels were also increased. Those changes may contribute to the suppression of cardiomyocyte contractility too.

In conclusion, the endogenous pore-forming toxins, such as circulating histones released after extensive tissue injury, and exogenous pore-forming toxins released during bacterial infection, may play important roles in the development of cardiac complications in many types of critical illnesses. In certain scenario, such as sepsis, both endogenous and exogenous toxins may synergistically harm the heart and cause severe complications to lead to early death of patients. Therefore more intensive investigation should be carried out to clarify their effects in patients as well as their molecular mechanisms in order to develop effective tools to diagnose, prevent and treat those cardiac complications induced by pore-forming toxins.

144

SELECTIVE INHIBITION OF PLASMA MEMBRANE CALCIUM ATPASE 4 IMPROVES VEGF-MEDIATED ANGIOGENESIS

¹Sathishkumar Kurusamy*, ^{2,3}Dolores López-Maderuelo, ⁴Robert Little, ¹David Cadagan, ⁵Aaron M Savage, ¹Jude C Ihugba, ¹Rhiannon R Baggott, ⁶Farjana B Rowther, ²Sara Martinez-Martinez, ^{2,3,7}Pablo Gomez del Arco, ¹Clare Murcott, ⁸Weiguang Wang, ⁴Delvac Oceandy, ^{4,9}Ludwig Neyses, ⁵Robert N Wilkinson, ⁴Elizabeth J Cartwright, ^{2,3}Juan Miguel Redondo, ¹Angel Luis Armesilla. ¹Cardiovascular Molecular Pharmacology Laboratory, School of Pharmacy; ²Gene Regulation in Cardiovascular Remodelling and Inflammation Group, Centro Nacional de Investigaciones Cardiovasculares, Madrid, Spain; ³CIBERCV; ⁴Division of Cardiovascular Sciences, University of Manchester, Manchester Academic Health Sciences Centre, Manchester, UK; ⁵Department of Infection, Immunity and Cardiovascular Disease and Bateson Centre, University of Sheffield, UK; ⁶Brain Tumour UK Neuro-oncology Research Centre; ⁷Department of Molecular Biology, Universidad Autonoma de Madrid, Madrid, Spain; ⁸Oncology Laboratory, Research Institute in Healthcare Science, Faculty of Science and Engineering, University of Wolverhampton, Wolverhampton, UK; ⁹University of Luxembourg

10.1136/heartjnl-2017-311726.143

Ischaemic cardiovascular disease is the leading cause of death worldwide. Therapeutic angiogenesis aims to stimulate the growth of new blood vessels from pre-existing ones to reperfuse ischaemic tissues. Our laboratory is characterising the molecular mechanisms that regulate activation of the calcineurin/NFAT pathway during VEGF-induced angiogenesis. We recently showed that the Plasma Membrane Calcium ATPase 4 (PMCA4) negatively regulates angiogenesis by establishing a molecular interaction with calcineurin. The identification of aurointricarboxylic acid (ATA) as an inhibitor of PMCA4 prompted us to hypothesise that ATA will enhance VEGF-induced angiogenesis.

Consistent with this hypothesis, we demonstrate in this work that inhibition of PMCA4 by treatment with ATA significantly increases the activity of calcineurin/NFAT signalling, and the subsequent expression of the NFAT-dependent, pro-angiogenic protein RCAN1.4 in VEGF-stimulated endothelial cells. Targeting PMCA4 with ATA significantly reduces the level of membrane-associated calcineurin, and the amount of calcineurin co-precipitated with PMCA4 in immunoprecipitation assays, indicating that ATA promotes disruption of the PMCA4/calcineurin interaction. ATA robustly enhances endothelial cell motility, and *in vitro* and *in vivo* blood vessel formation, with no harmful effects to the cells. Interestingly, incubation of HUVECs with low concentration of ATA had no effect on the viability of the cells or the development of zebra fish embryos. However, higher ATA concentrations were associated with cellular and embryo toxicity.

Our study provides evidence for the therapeutic potential of targeting endothelial PMCA4 to improve VEGF-based pro-angiogenic interventions, and highlights possible clinical applications for PMCA4 inhibitors in the treatment of ischaemic cardiovascular disease.