OPEN ACCESS

ORIGINAL RESEARCH ARTICLE

Cardiac myosin-binding protein C is a novel marker of myocardial injury and fibrosis in aortic stenosis

Atul Anand,¹ Calvin Chin,¹ Anoop S V Shah,¹ Jacek Kwiecinski,^{1,2} Alex Vesey,¹ Joanna Cowell,³ Ekkehard Weber,⁴ Thomas Kaier,⁵ David E Newby,¹ Marc Dweck,¹ Michael S Marber,⁵ Nicholas L Mills¹

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/heartjnl-2017-312257>).

¹BHF Centre for Cardiovascular Science, University of Edinburgh, Edinburgh, UK

²First Department of Cardiology, Poznan University of Medical Sciences, Poznan, Poland

³Department of Geriatric Medicine, Royal Victoria Building, Edinburgh, UK

⁴Institute of Physiological Chemistry, Martin Luther University Halle-Wittenberg, Halle, Germany

⁵King's College London BHF Centre, The Rayne Institute, St Thomas' Hospital, London, UK

Correspondence to

Dr Atul Anand, BHF Centre for Cardiovascular Science, University of Edinburgh, Edinburgh EH16 4SB, UK; atul.anand1@nhs.net

Received 2 August 2017

Revised 15 November 2017

Accepted 16 November 2017

Published Online First

1 December 2017

Listen to Podcast
www.goo.gl/EivKMP

ABSTRACT

Objective Cardiac myosin-binding protein C (cMyC) is an abundant sarcomeric protein and novel highly specific marker of myocardial injury. Myocyte death characterises the transition from hypertrophy to replacement myocardial fibrosis in advanced aortic stenosis. We hypothesised that serum cMyC concentrations would be associated with cardiac structure and outcomes in patients with aortic stenosis.

Methods cMyC was measured in two cohorts in which serum had previously been prospectively collected: a mechanism cohort of patients with aortic stenosis (n=161) and healthy controls (n=46) who underwent cardiac MRI, and an outcome cohort with aortic stenosis (n=104) followed for a median of 11.3 years.

Results In the mechanism cohort, cMyC concentration correlated with left ventricular mass (adjusted $\beta^2=11.0$ g/m² per log unit increase in cMyC, $P<0.001$), fibrosis volume (adjusted $\beta^2=8.0$ g, $P<0.001$) and extracellular volume (adjusted $\beta^2=1.3\%$, $P=0.01$) in patients with aortic stenosis but not in controls. In those with late gadolinium enhancement (LGE) indicative of myocardial fibrosis, cMyC concentrations were higher (32 (21–56) ng/L vs 17 (12–24) ng/L without LGE, $P<0.001$). cMyC was unrelated to coronary calcium scores. Unadjusted Cox proportional hazards analysis in the outcome cohort showed greater all-cause mortality (HR 1.49 per unit increase in log cMyC, 95% CI 1.11 to 2.01, $P=0.009$).

Conclusions Serum cMyC concentration is associated with myocardial hypertrophy, fibrosis and an increased risk of mortality in aortic stenosis. The quantification of serum sarcomeric protein concentrations provides objective measures of disease severity and their clinical utility to monitor the progression of aortic stenosis merits further study.

Clinical trial registration NCT1755936; Post-results.

INTRODUCTION

Aortic stenosis is the most common valvular disease in the Western world and the incidence is rising in keeping with an ageing population.¹ The response of the myocardium to aortic stenosis is variable, with heterogeneity in the development of ventricular hypertrophy and in how this process ultimately decompensates.^{2–4} This results in a poor correlation between the severity of stenosis and the development of symptoms. Decompensation of the hypertrophic response in aortic stenosis is driven by two processes: progressive myocyte cell death and myocardial fibrosis.⁵

Biomarkers of myocardial injury are therefore an attractive addition to current imaging markers of disease progression, perhaps providing critical evidence of early decompensation that may identify patients who would benefit from early valve replacement. We have previously demonstrated that cardiac troponin I concentration is associated with advanced hypertrophy, replacement fibrosis and poor long-term outcomes in patients with aortic stenosis,⁶ suggesting that myocardial injury in advanced aortic stenosis is common, detectable and of prognostic importance.

Cardiac myosin-binding protein C (cMyC) is a cardiac-restricted sarcomeric protein located on the thick filament.^{7,8} Recent advances in assay technology have allowed measurement of cMyC with high precision at extremely low concentrations.⁹ It is more abundant in myocardial tissue and the circulation than cardiac troponins and has an important role in the assembly and function of the cardiac sarcomere.^{10–12} Indeed, mutations in the MYBPC3 gene-encoding cMyC are the most common known genetic cause of hypertrophic cardiomyopathy.¹³ Furthermore, the proteolytic cleavage of cMyC is highly regulated by a variety of myocardial kinases¹⁴ and the resultant peptide is cardiotoxic.¹⁵ Hence, there is growing interest in the protein as both a biomarker and a determinant of myocardial injury.¹⁶ Given myocyte death characterises the transition from hypertrophy to replacement myocardial fibrosis in advanced aortic stenosis, we hypothesised that serum cMyC concentrations would be associated with cardiac structure and outcomes in patients with aortic stenosis.

METHODS

We evaluated cMyC in two cohorts: a *mechanism* cohort of patients with aortic stenosis and healthy controls with cardiac MRI (CMR), and an *outcome* cohort of patients with aortic stenosis with more than 10 years of clinical follow-up. These groups were derived from existing studies in patients with stable aortic stenosis recruited from cardiology clinics across the South East of Scotland, where serum had been prospectively collected and frozen at the time of study inclusion. Additionally, a subgroup of patients from the mechanism cohort underwent myocardial biopsy at the time of subsequent aortic valve replacement (AVR), providing exploratory histological correlation with cMyC concentrations. The studies were conducted in

To cite: Anand A, Chin C, Shah ASV, et al. *Heart* 2018;**104**:1101–1108.

accordance with the Declaration of Helsinki. Written informed consent was obtained from all participants.

Mechanism cohort

The mechanism cohort consisted of 161 patients with mild to severe aortic stenosis and 46 healthy volunteers without evidence of significant valvular heart disease enrolled in an observational study assessing the role of myocardial fibrosis in aortic stenosis (NCT:01755936 post-results). Exclusion criteria comprised significant (moderate or severe) non-aortic valvular disease or any cardiomyopathy (acquired or inherited). The imaging protocol undertaken in all participants has been described in detail previously.^{6 17–19} Briefly, CMR was performed using a 3T scanner (MAGNETOM Verio, Siemens AG, Healthcare Sector, Erlangen, Germany). Dedicated software was used to assess left ventricular (LV) volume and mass indexed to body surface area and to calculate ejection fraction. Diffuse myocardial fibrosis was determined by fibrosis volume and the extracellular volume (ECV) fraction in keeping with current evidence of reproducibility from T1 mapping.^{19 20} Focal myocardial replacement fibrosis was determined by the late gadolinium enhancement (LGE) technique, with its presence determined visually and independently by two experienced assessors. Comprehensive echocardiography was performed on all subjects to classify markers of aortic stenosis severity according to European Association of Echocardiography/American Society of Echocardiography guidelines.²¹

Outcome cohort

This cohort was derived from the Scottish Aortic Stenosis and Lipid Lowering Trial, Impact of Regression (SALTIRE) study. The study design, recruitment and findings have been reported previously.^{22 23} Briefly, between March 2001 and April 2002, a total of 155 patients with asymptomatic moderate to severe aortic stenosis were randomised to receive either atorvastatin or placebo. Stored sample remained for cMyC analysis in 104 patients. In addition to comprehensive echocardiography, CT calcium scoring of the coronary arteries was performed (Twin II Flash, Philips Medical Systems).

Outcomes data were obtained by two independent investigators who were blinded to cMyC results. The General Register of Scotland was searched for all deaths. The cause of death was adjudicated for a cardiac cause using additional information from electronic health records if necessary. Disagreements were resolved by consensus. Electronic health records were also reviewed in all cases for evidence of surgical AVR.

Blood sampling and analysis

Serum cMyC concentrations were measured in duplicate on the single molecule counting Erenna platform (Singulex/Merck Millipore, CA, USA). The assay has a lower limit of detection (LoD) of 0.4 ng/L, a lower limit of quantification (LLoQ) of 1.2 ng/L (at 20% coefficient of variation, CV) and reasonable recovery (107.1%±3.7%; mean±SD), dilutional linearity (101.0%±7.7%), and intraseries (CV 11%±3%) and interseries (CV 13%±3%) precision. Cardiac troponin was determined using a high-sensitivity assay (Abbott ARCHITECT STAT, Abbott Diagnostics, IL, USA) as previously described. This assay has an LoD of 1.2 ng/L and based on our previous work an LoQ of 1.5 ng/L (at 20% CV).⁶

Myocardial biopsy and histological analysis

Tru-cut myocardial biopsies were obtained from 10 patients in the mechanism cohort who underwent AVR and also had

cMyC measured. For autophagy and oncosis assessment, formalin-fixed, paraffin-embedded 4 µm thick tissue sections were cut and dehydrated. Further details of the histological analysis are provided in the online supplementary material.

Statistical analysis

Statistical analysis was performed using the statistical software R V.3.3.2 (<http://www.r-project.org>). Continuous variables are presented as mean (SD) or median (IQR) for non-parametric data. We used analysis of variance to compare continuous parametric data and the Kruskal-Wallis test for non-parametric data across tertiles of cMyC. Categorical variables are presented as absolute numbers (percentage) and were compared using χ^2 test. Due to the positive skewing in the sample, cMyC concentrations were log transformed prior to inclusion in modelling. Multivariate linear regression modelling was used to assess the change in markers of aortic stenosis severity with serum cMyC concentration. Receiver operating characteristic (ROC) curve analysis and multivariate logistic regression modelling were used to assess the relationship between cMyC concentrations and LGE. Survival analysis was performed with a Kaplan-Meier analysis using time to any cause of death, with significance calculated by log-rank test. This relationship was also assessed using Cox proportional hazard modelling. AVR was included as a time-varying covariate as survival could be expected to improve in those who received surgery during the follow-up period. A value of $P<0.05$ was considered statistically significant.

RESULTS

The mechanism cohort consisted of 161 individuals with aortic stenosis (mean age 69 years, AV_{max} 3.8±0.9 m/s, 70% male) and 46 healthy control participants (mean age 58 years, AV_{max} 1.4±0.2 m/s, 63% male). The outcome cohort consisted of 104 patients (mean age 68 years, AV_{max} 3.4±0.7 m/s, 68% male) with a median follow-up period of just more than 11 years (4067 (3882–4161) days).

Baseline characteristics by tertile of cMyC are presented in tables 1 and 2 for the mechanism and outcome cohorts. cMyC was measurable above the LLoQ (1.2 ng/L) in all but one subject, giving an overall detection rate of 99.7%, including all healthy controls (online supplementary figure S1). Similar to observations with cardiac troponin, cMyC concentrations were positively skewed across all cohorts (online supplementary figure S2) and weakly correlated with renal function (online supplementary figure S3). The median cMyC concentration was similar among patients with aortic stenosis in the mechanism (20.5 (13.7–33.2) ng/L) and outcome (18.2 (12.2–30.1) ng/L) cohorts ($P=0.21$), with both being higher than in controls (9.5 (7.6–15.1) ng/L, $P<0.001$ for both). Samples were tested in duplicate, with a CV between repeated measurements of 5.9%±5.1% in the mechanism cohort, 6.1%±4.9% across the outcome cohort and 8.5%±4.5% in controls.

Cardiac troponin concentrations were above the LLoQ in 92.3% of subjects, but in only 65.2% of healthy controls. A strong correlation was observed between cMyC and cardiac troponin across all samples tested ($r=0.74$, 95% CI 0.69 to 0.79, $P<0.0001$, online supplementary figure S1).

cMyC and mechanistic imaging markers

In adjusted multiple regression analyses, cMyC concentration was associated with indexed LV mass in those with aortic stenosis who underwent CMR imaging ($\beta=11.0\text{ g/m}^2$ per log unit increase in cMyC after adjustment for age, sex, renal

Table 1 Baseline characteristics of the mechanism cohort by tertile of cMyC and controls

	All (n=161)	Tertile 1 3–16 ng/L (n=54)	Tertile 2 17–28 ng/L (n=55)	Tertile 3 29–171 ng/L (n=52)	P value	Controls (n=46)
Age (years)	68.5 (11.4)	64.3 (13.0)	69.4 (9.3)	71.8 (10.5)	0.002	57.9 (20.5)
Sex (male), n (%)	112 (69.6)	29 (53.7)	42 (76.4)	41 (78.8)	0.008	29 (63.0)
BMI (kg/m ²)	28.9 (4.8)	29.1 (5.2)	28.7 (4.0)	29.0 (5.1)	0.88	26.8 (3.8)
BSA (m ²)	1.9 (0.2)	1.9 (0.2)	1.9 (0.2)	1.9 (0.2)	0.71	1.9 (0.2)
Comorbidity						
Diabetes mellitus, n (%)	24 (14.9)	9 (16.7)	6 (10.9)	9 (17.3)	0.59	0 (0.0)
Hypertension, n (%)	109 (67.7)	30 (55.6)	42 (76.4)	37 (71.2)	0.06	12 (26.1)
Hyperlipidaemia, n (%)	72 (44.7)	20 (37.0)	22 (40.0)	30 (57.7)	0.07	8 (17.4)
IHD, n (%)	61 (37.9)	12 (22.2)	23 (41.8)	26 (50.0)	0.01	2 (4.3)
Previous PCI, n (%)	25 (15.5)	6 (11.1)	8 (14.5)	11 (21.2)	0.35	2 (4.3)
Previous CABG, n (%)	7 (4.4)	2 (3.7)	1 (1.9)	4 (7.7)	0.33	0 (0.0)
Medications						
Antiplatelet, n (%)	79 (49.1)	17 (31.5)	29 (52.7)	33 (63.5)	0.004	7 (15.2)
ACE-I/ARB, n (%)	67 (41.6)	16 (29.6)	28 (50.9)	23 (44.2)	0.07	9 (19.6)
Beta blocker, n (%)	55 (34.2)	16 (29.6)	17 (30.9)	22 (42.3)	0.32	5 (10.9)
Diuretic, n (%)	52 (32.3)	16 (29.6)	21 (38.2)	15 (28.8)	0.51	4 (8.7)
Blood tests						
Creatinine (μmol/L)	79.1 (17.1)	73.5 (11.1)	78.6 (15.5)	85.2 (21.5)	0.002	72.5 (15.3)
eGFR (mL/min/1.73 m ²)	86.3 (18.5)	89.0 (15.9)	88.0 (18.8)	81.7 (20.1)	0.09	97.7 (19.8)
Troponin I (ng/L)	6.6 (3.8–12.3)	3.3 (2.3–4.5)	6.7 (4.8–9.5)	15.5 (10–30.2)	<0.001	2.7 (1.1–5.4)
Echo parameters						
AV _{max} (m/s)	3.8 (0.9)	3.4 (0.8)	3.9 (0.8)	4.3 (0.9)	<0.001	1.4 (0.2)
AV area (cm ²)	1.0 (0.4)	1.1 (0.4)	0.9 (0.3)	0.9 (0.4)	0.006	2.4 (0.6)
AV MPG (mm Hg)	34.4 (18.6)	25.9 (13.5)	34.5 (16.6)	43.2 (21.3)	<0.001	4.2 (1.4)
Indexed LV mass (g/m ²)	122.7 (32.0)	102.3 (22.8)	127.6 (27.3)	139.4 (33.7)	<0.001	93.2 (24.4)
E/e'	14.6 (7.6)	12.4 (4.8)	13.1 (5.0)	18.3 (10.4)	<0.001	8.6 (2.5)
MRI parameters						
Indexed LV mass (g/m ²)	88.8 (21.5)	74.2 (15.2)	90.1 (16.6)	102.5 (22.1)	<0.001	65.8 (13.7)
Ejection fraction (%)	66.8 (7.3)	67.5 (5.6)	65.8 (8.3)	67.2 (7.8)	0.44	64.4 (4.4)
Indexed SV (mL/m ²)	47.6 (10.2)	44.3 (8.4)	47.1 (9.6)	51.4 (11.2)	0.001	46.1 (8.2)
Indexed ESV (mL/m ²)	24.2 (9.8)	21.6 (6.6)	25.5 (11.5)	25.6 (10.2)	0.055	26.0 (7.4)
Indexed EDV (mL/m ²)	71.8 (16.9)	65.9 (13.3)	72.6 (17.8)	77.0 (17.8)	0.003	72.1 (14.6)
ECV (%)	27.8 (2.6)	27.1 (2.2)	27.4 (2.1)	28.9 (3.2)	0.001	26.6 (1.7)
Fibrosis volume (g)	44.6 (15.3)	36.0 (8.9)	44.2 (11.1)	54.5 (18.5)	<0.001	31.0 (7.6)
Outcomes						
AVR, n (%)	41 (25.5)	8 (14.8)	20 (36.4)	13 (25.0)	0.04	0 (0.0)
Cardiac death, n (%)	1 (0.6)	0 (0.0)	0 (0.0)	1 (1.9)	0.35	0 (0.0)
All-cause death, n (%)	6 (3.7)	1 (1.9)	1 (1.8)	4 (7.7)	0.19	0 (0.0)

Values are number (%), mean (SD) or median (IQR). Results in bold are statistically significant.

ACE-I, ACE inhibitor; ARB, angiotensin receptor blocker; AV, aortic valve; AVR, aortic valve replacement; BMI, body mass index; BSA, body surface area; CABG, coronary artery bypass grafting; cMyC, cardiac myosin-binding protein C; ECV, extracellular volume fraction; EDV, end-diastolic volume; eGFR, estimated glomerular filtration rate; ESV, end-systolic volume; IHD, ischaemic heart disease; PCI, percutaneous coronary intervention; LV, left ventricle; MPG, mean pressure gradient; SV, stroke volume.

function, AV_{max}, cardiac troponin and comorbidity; 95% CI 4.7 to 17.3, $P < 0.001$, figure 1A). This relationship was numerically positive, but did not approach statistical significance among healthy controls ($\beta = 2.7 \text{ g/m}^2$, 95% CI -4.8 to 10.1 , $P = 0.47$, online supplementary figure S4). Similarly, fibrosis volume was related to cMyC in those with aortic stenosis (adjusted $\beta = 8.0 \text{ g}$ per log unit increase in cMyC; 95% CI 3.5 to 12.6, $P < 0.001$, figure 1B) but not in controls ($\beta = 1.8 \text{ g}$, 95% CI -1.6 to 5.2 , $P = 0.28$, online supplementary figure S4). ECV as a marker of diffuse myocardial fibrosis was associated with cMyC in patients with aortic stenosis (adjusted $\beta = 1.3\%$, 95% CI 0.3% to 2.3%, $P = 0.01$), but not in healthy controls (adjusted $\beta = 0.5\%$, 95% CI -0.5% to 1.5% , $P = 0.32$). Detailed modelling is shown in online supplementary tables S1–S3. cMyC concentrations were related

to severity of aortic stenosis across the range of AV_{max} measures in the mechanism cohort (adjusted $\beta = 0.80 \text{ m/s}$, 95% CI 0.50 to 1.10, $P < 0.001$, figure 1C), although this was not observed in the narrower range included in the outcome cohort (online supplementary figure S4). An association was further noted between cMyC and diastolic function, measured by echocardiography E/e' in the mechanism cohort (adjusted $\beta = 3.76$, 95% CI 1.21 to 6.32, $P = 0.004$, figure 1D). There was no relationship between cMyC and objective measures of coronary disease by CT calcium scoring (adjusted $\beta = 0.03$, 95% CI -0.17 to 0.23 , $P = 0.76$, online supplementary figure S4).

LGE was present in 57 (35.4%) of patients with aortic stenosis in the mechanism cohort. cMyC concentrations were almost double in those with evidence of LGE compared with

Table 2 Baseline characteristics of the outcome cohort by tertile of cMyC

	All patients (n=104)	Tertile 1 5–14 ng/L (n=35)	Tertile 2 15–26 ng/L (n=35)	Tertile 3 27–132 ng/L (n=34)	P value
Age (years)	68.2 (9.8)	64.4 (10.5)	70.1 (9.0)	70.0 (9.1)	0.02
Sex (male), n (%)	71 (68.3)	21 (60.0)	19 (54.3)	31 (91.2)	0.002
BMI (kg/m ²)	27.9 (4.5)	27.6 (4.1)	28.1 (4.9)	27.8 (4.7)	0.91
BSA (m ²)	1.9 (0.2)	1.9 (0.2)	1.9 (0.2)	2.0 (0.2)	0.19
Comorbidity					
Diabetes mellitus, n (%)	2 (1.9)	1 (2.9)	0 (0.0)	1 (2.9)	0.60
Hypertension, n (%)	57 (54.8)	16 (45.7)	24 (68.6)	17 (50.0)	0.13
Hyperlipidaemia, n (%)	7 (6.7)	0 (0.0)	5 (14.3)	2 (5.9)	0.06
IHD, n (%)	17 (16.3)	2 (5.7)	10 (28.6)	5 (14.7)	0.03
Medications					
Aspirin, n (%)	42 (40.4)	13 (37.1)	16 (45.7)	13 (38.2)	0.74
Anticoagulant, n (%)	10 (9.6)	1 (2.9)	4 (11.4)	5 (14.7)	0.23
ACE-I/ARB, n (%)	20 (19.2)	5 (14.3)	7 (20)	8 (23.5)	0.63
Beta blocker, n (%)	23 (22.1)	8 (22.9)	11 (31.4)	4 (11.8)	0.14
Blood tests					
Creatinine (μmol/L)	91.3 (21.3)	85.3 (18.1)	89.3 (17.9)	100.1 (25.6)	0.01
eGFR (mL/min/1.73 m ²)	73.5 (17.4)	77.2 (13.9)	71.7 (16.4)	71.6 (21.3)	0.31
Troponin I (ng/L)	7.2 (5.4–12.6)	4.9 (3.9–6.0)	8.3 (6.7–10.6)	14.3 (12.1–17.9)	<0.001
Echo parameters					
AV _{max} (m/s)	3.4 (0.7)	3.4 (0.7)	3.4 (0.8)	3.5 (0.6)	0.90
AV area (cm ²)	1.0 (0.4)	1.0 (0.4)	1.0 (0.4)	1.0 (0.4)	0.89
AV MPG (mm Hg)	48.8 (19.2)	47.6 (18.7)	49.7 (21.8)	49.2 (17.1)	0.89
Indexed LV mass (g/m ²)	184.4 (52.2)	170.0 (56.8)	175.7 (38.8)	208.7 (52.0)	0.004
Ejection fraction (%)	66 (31)	69 (22)	66 (25)	61 (44)	0.64
CT coronary calcium score	382.6 (611.1)	370.3 (608.3)	491.0 (783.2)	283.6 (360.2)	0.37
Outcomes					
AVR, n (%)	48 (46.2)	19 (54.3)	15 (42.9)	14 (41.2)	0.49
Cardiac death, n (%)	16 (15.4)	4 (11.4)	7 (20.0)	5 (14.7)	0.61
All-cause death, n (%)	36 (34.6)	7 (20.0)	14 (40.0)	15 (44.1)	0.08

Values are number (%), mean (SD) or median (IQR). Results in bold are statistically significant.

ACE-I, ACE inhibitor; ARB, angiotensin receptor blocker; AV, aortic valve; AVR, aortic valve replacement; BMI, body mass index; BSA, body surface area; cMyC, cardiac myosin-binding protein C; eGFR, estimated glomerular filtration rate; IHD, ischaemic heart disease; LV, left ventricle; MPG, mean pressure gradient.

those without (32.3 (21.3–56.3) ng/L vs 17.2 (11.5–24.2) ng/L, $P<0.001$). By ROC analysis, cMyC improved discrimination for LGE beyond age and gender (area under the curve (AUC) 0.77, 95% CI 0.70 to 0.85, [figure 2A](#)). A sensitivity analysis restricted to those with mid-wall fibrosis only (77% of those with any LGE) demonstrated similar discrimination (AUC 0.77, 95% CI 0.70 to 0.84). Using logistic regression modelling adjusted for age and sex, the predicted probability of LGE was seen to progressively increase with cMyC concentration ([figure 2B](#)). Univariate correlations between cMyC and baseline variables used for covariate adjustment in the mechanism cohort are presented in online supplementary table S4.

cMyC and myocyte death

Clear differences were observed in staining patterns for oncosis and autophagy between subjects with low and high cMyC concentrations ([figure 3](#)). Exploratory analysis suggested a relationship between cMyC and the rate of myocyte death (expressed as the sum of apoptosis, oncosis and autophagy counts) in 10 subjects with myocardial biopsy tissue taken at the time of AVR ($r=0.67$, 95% CI 0.08 to 0.92, $P=0.03$, online supplementary figure S5).

cMyC and long-term outcomes

Subjects in the outcome cohort were stratified by tertile of cMyC ([table 2](#)). As with CMR in the mechanism cohort, indexed LV mass by echocardiography increased across tertiles of cMyC (170.0±56.8 g in lowest tertile vs 208.7±52.0 g in highest tertile, $P=0.004$). There were no consistent differences in cardiac risk factors across the tertiles. During the follow-up period, 36 (34.6%) subjects died, of which 16 (15.4%) were adjudicated as cardiac deaths. Forty-eight patients (46.2%) within the cohort underwent surgical AVR, with no cases of transcatheter aortic valve implantation undertaken. There was a trend towards poorer survival over the period of follow-up with increasing tertile of cMyC ([figure 4](#), log-rank test for difference $P=0.07$).

In Cox proportional hazards analysis, cMyC concentration was associated with an increased risk of all-cause mortality over the follow-up period after inclusion of AVR as a time-varying covariate (HR 1.49 per log unit increase of cMyC, 95% CI 1.11 to 2.01, $P=0.009$). However, following adjustment for age, sex, AV_{max} or CT coronary calcium scores, statistical significance was lost. Age was significantly associated with cMyC in all models (online supplementary table S5).

Figure 1 Relationships between cMyC and markers of disease severity in the mechanism cohort. β values (95% CI) represent change in the progression variable for each log unit change in cMyC concentration after adjustment in multivariate modelling. Markers of disease severity are: (A) indexed LV mass (g/m²) determined by MRI; (B) fibrosis volume (g) by MRI; (C) AV_{max} (m/s) determined by echocardiography; (D) diastolic function from echocardiographic measures of E/e' ratio. *Adjusted for age, sex, glomerular filtration rate, AV_{max}, cardiac troponin, history of ischaemic heart disease, diabetes and hypertension. +As above plus body surface area. ‡As above excluding AV_{max}. AV, aortic valve; cMyC, cardiac myosin-binding protein C; LV, left ventricle.

DISCUSSION

In a comprehensive series of clinical assessments, we report the relationship between serum cMyC concentration, a novel marker of myocardial injury, and cardiac structure and outcomes

in patients with aortic stenosis. We have made several important observations. First, serum cMyC concentrations can be reliably quantified in the vast majority of patients with aortic stenosis. Second, cMyC strongly associates with markers of both diffuse

Figure 2 cMyC and late gadolinium enhancement. (A) Receiver operating characteristic (ROC) curve analysis for outcome of LGE. Addition of cMyC improves the prediction of age and gender for LGE, $P = 0.002$ (DeLong's method with bootstrapping). (B) Logistic regression modelling for the predicted probability of LGE with serum cMyC concentration (adjusted for age and sex). Shaded area represents 95% CI. AUC, area under the curve; cMyC, cardiac myosin-binding protein C; LGE, late gadolinium enhancement.

Figure 3 Patterns of myocyte death in patients with aortic stenosis. Images showing differing patterns of *oncosis* and *autophagy* visualised using a 3,3'-diaminobenzidine-based detection kit in individuals with aortic stenosis. (A) *Oncosis* and (B) *autophagy* in a patient with a low (9.4 ng/L) cMyC concentration. (C) *Oncosis* and (D) *autophagy* in a patient with a high (87.2 ng/L) cMyC concentration. There is a marked difference in staining intensity by cMyC concentration for both oncosis and autophagy. cMyC, cardiac myosin-binding protein C.

interstitial and focal replacement fibrosis as well as indexed LV mass, a marker of the hypertrophic response to aortic stenosis. These observations were independent of the severity of

valve stenosis and cMyC concentrations were unrelated to coronary artery disease. Third, cMyC concentrations are strongly correlated with cardiac troponin, and in histological analyses are associated with myocyte cell death, suggesting that sarcomeric protein release is a direct consequence of the maladaptive myocardial response to aortic stenosis. We believe these novel observations demonstrate the utility and validity of cMyC as a marker of sustained chronic myocardial injury.

Assay performance

The release profile of cMyC has been previously studied after acute injury in myocardial infarction, alcohol ablation for hypertrophic cardiomyopathy and coronary artery bypass grafting.^{7 16} These models have shown cMyC to be more abundant, to be released earlier following injury and to decline more rapidly when compared with cardiac troponin. These observations suggest that cMyC has enhanced potential for dynamic monitoring of myocardial injury and disease. However, previous generations of the cMyC assay had limited sensitivity at the low levels that would be expected in the presence of chronic myocardial injury such as in aortic stenosis. For example, cMyC was only detectable in 2 out of 20 patients with hypertrophic cardiomyopathy.²⁴ Here we report cMyC measurements from a high-sensitivity assay using magnetic nanoparticle and single molecule counting technology, and demonstrate near universal quantification in patients with aortic stenosis and healthy controls.

cMyC as a biomarker of disease

Our observations suggest cMyC is a sensitive marker of the hypertrophic and subsequent fibrotic myocardial response to

Figure 4 Survival by tertiles of cMyC in the outcome cohort. cMyC, cardiac myosin-binding protein C.

aortic stenosis. ECV fraction on CMR T1 mapping is a marker of diffuse interstitial fibrosis, which may precede the focal mid-wall replacement fibrosis detected by LGE.^{19,20} Histological studies have suggested that myocyte death drives this transition.⁵ Our exploratory analysis in myocardial biopsy tissue relates combined measures of autophagy, oncosis and apoptosis to serum cMyC concentrations, providing histological correlation of biomarker release as a result of myocyte death. This combination of autophagy and oncosis in failing myocardial tissue has been independently associated with mortality in larger studies.²⁵

Our observations are consistent with our previous findings using a high-sensitivity cardiac troponin I assay. Using two independent sarcomeric proteins, we have now demonstrated that serum markers of myocardial injury mirror important pathological changes in the myocardium in response to aortic stenosis.^{6,26,27} Similar to cardiac troponin, although cMyC was related to AV_{max} by echocardiography in the mechanism cohort, its association with markers of hypertrophy, fibrosis and diastolic function was independent of the severity of aortic stenosis. This suggests that these novel markers of the myocardial response to stenosis will add to conventional clinical assessment in aortic stenosis. While CMR imaging provides important detail on the extent of the progression to fibrosis and heart failure, it is a resource-intensive investigation.^{2,28} As such, there is great potential for simple blood measures of myocardial injury to improve decision-making in aortic stenosis, to better target detailed imaging or even help determine the timing of surgery. Current guidelines advocate AVR in the presence of LV impairment or symptoms, but these are challenging to define objectively and may signify that irreversible myocardial fibrosis has already developed. A precise and early marker of myocardial injury such as cMyC may have great clinical utility in patients with aortic stenosis.

We demonstrate an increased mortality risk with rising cMyC concentrations, but this association was not independent of other variables. Our outcome cohort benefits from extensive follow-up over 11 years after sampling, but the cohort size was restricted by limited remaining stored samples. This analysis is likely to be underpowered and further work is required in larger studies to clarify whether cMyC will provide additional prognostic information in the assessment of patients with aortic stenosis.

Limitations

We acknowledge some limitations with the current study in addition to sample size. There is only limited overlap for comparison of imaging endpoints between our two cohorts, although the magnitude of associations between cMyC and LV mass are similar. The clinical utility of biomarkers such as cMyC is more likely to be of interest in severe cases of aortic stenosis approaching the need for valve replacement, where myocardial injury becomes more pronounced. Our mechanism and outcome cohorts contained only 84 and 52 patients with severe aortic stenosis, respectively. While this represents half of our study population, further evaluation in a larger population with severe disease would be informative. The histological analysis presented is exploratory and the Tru-cut technique used may provide limited volumes of myocardium compared with novel scalpel methods.²⁹

Our findings add to our previous observations with cardiac troponin.^{6,26,27} Compared with those assays, the latest generation cMyC assay is still in the early stages of development, with no definitive healthy population studies to derive a normal reference range. However, our study suggests that the biomarker will

be measurable in virtually all individuals and is likely to meet the criteria for a high-sensitivity assay.³⁰ Larger cohort studies are necessary to assess the relationship and potential differences in cMyC and cardiac troponin release, and to determine whether the association of cMyC and long-term outcomes is independent of other patient factors and cardiac biomarkers. Further validation with prospective serial sampling of cMyC in patients with aortic stenosis prior to clinical decompensation is now required.

CONCLUSIONS

Serum cMyC concentration is associated with myocardial hypertrophy, fibrosis and an increased risk of mortality in aortic stenosis. The quantification of serum sarcomeric protein concentrations provides objective measures of disease severity and their clinical utility to monitor the progression of aortic stenosis merits further study.

Key messages

What is already known on this subject?

The progression of aortic stenosis is unpredictable and irreversible myocardial fibrosis may develop before symptoms are apparent. Cardiac troponin I as a biomarker of myocardial injury is associated with advanced hypertrophy, replacement fibrosis and poor long-term outcomes in patients with aortic stenosis.

What might this study add?

Cardiac myosin-binding protein C (cMyC) is a cardiac sarcomeric protein and novel highly sensitive biomarker of acute myocardial injury. In two cohorts of patients with aortic stenosis, increasing concentrations of cMyC were associated with greater myocardial hypertrophy, fibrosis and death, even after adjustment for the severity of aortic stenosis and cardiac troponin concentrations. This is the first study to measure cMyC using a high-precision assay in patients with aortic stenosis.

How might this impact on clinical practice?

Objective blood biomarkers are not currently included in clinical guidelines for the management of aortic stenosis. This study adds to evidence that quantification of serum sarcomeric proteins has major potential to objectively track disease progression in this condition. Further validation with prospective serial sampling of cMyC in patients with aortic stenosis prior to clinical decompensation is now required.

Acknowledgements We thank Audrey White for obtaining echocardiographic measures in all patients and Edwin Carter for his assistance in preparing these samples.

Contributors AA conducted the analysis presented and drafted the submission. CC conducted the study for the mechanism cohort. ASVS contributed to analysis of data. JK performed the histological analysis. JC conducted the study for the outcomes cohort. MD, DEN, NLM and MSM contributed to the planning, conduct and reporting of the studies included in the manuscript. The above named authors and all other authors named provided further critical review of the manuscript and further input on revisions.

Funding This research is supported by research fellowships from Chest Heart and Stroke Scotland (15/A163) and the British Heart Foundation (FS/10/024/28266 and FS/15/13/31320). The development of the cMyC biomarker is supported by grants from the MRC UK (G1000737), Guy's and St Thomas' Charity (R060701, R100404), British Heart Foundation (TG/15/1/31518) and the UK Department of Health through the National Institute for Health Research Biomedical Research Centre award to Guy's and St Thomas' National Health Service Foundation Trust. DEN is supported by the British Heart Foundation (CH/9/002) and the Wellcome Trust (WT103782AIA).

Competing interests AA has received speaker fees from Abbott Laboratories. ASVS has acted as a consultant for Abbott Laboratories. NLM has acted as a consultant for Abbott Laboratories, Beckman-Coulter, Roche and Singulex. Singulex was contracted to undertake the analyses of cMyC on a fee-for-service basis and holds no commercial interest. MSM is named as an inventor on a patent held by King's College London for the detection of cardiac myosin-binding protein C as a biomarker of myocardial injury. All other authors have no conflicts of interest to declare.

Ethics approval The studies were approved by the South East Scotland Research Ethics Committee.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Osnabrugge RLI, Mylotte D, Head SJ, *et al.* Aortic stenosis in the elderly. *J Am Coll Cardiol* 2013;62:1002–12.
- Dweck MR, Joshi S, Murigu T, *et al.* Midwall fibrosis is an independent predictor of mortality in patients with aortic stenosis. *J Am Coll Cardiol* 2011;58:1271–9.
- Dweck MR, Joshi S, Murigu T, *et al.* Left ventricular remodeling and hypertrophy in patients with aortic stenosis: insights from cardiovascular magnetic resonance. *J Cardiovasc Magn Reson* 2012;14:50.
- Cioffi G, Faggiano P, Vizzardi E, *et al.* Prognostic effect of inappropriately high left ventricular mass in asymptomatic severe aortic stenosis. *Heart* 2011;97:301–7.
- Hein S, Arnon E, Kostin S, *et al.* Progression from compensated hypertrophy to failure in the pressure-overloaded human heart: structural deterioration and compensatory mechanisms. *Circulation* 2003;107:984–91.
- Chin CW, Shah AS, McAllister DA, *et al.* High-sensitivity troponin I concentrations are a marker of an advanced hypertrophic response and adverse outcomes in patients with aortic stenosis. *Eur Heart J* 2014;35:2312–21.
- Baker JO, Tyther R, Liebetrau C, *et al.* Cardiac myosin-binding protein C: a potential early biomarker of myocardial injury. *Basic Res Cardiol* 2015;110:23.
- Jacquet S, Yin X, Sicard P, *et al.* Identification of cardiac myosin-binding protein C as a candidate biomarker of myocardial infarction by proteomics analysis. *Mol Cell Proteomics* 2009;8:2687–99.
- Marjot J, Liebetrau C, Goodson RJ, *et al.* The development and application of a high-sensitivity immunoassay for cardiac myosin-binding protein C. *Transl Res* 2016;170:17–25.
- Aye TT, Scholten A, Taouatas N, *et al.* Proteome-wide protein concentrations in the human heart. *Mol Biosyst* 2010;6:1917–27.
- Barefield D, Sadayappan S. Phosphorylation and function of cardiac myosin binding protein-C in health and disease. *J Mol Cell Cardiol* 2010;48:866–75.
- Marjot J, Kaier TE, Martin ED, *et al.* Quantifying the release of biomarkers of myocardial necrosis from cardiac myocytes and intact myocardium. *Clin Chem* 2017;63:990–6.
- Van Driest SL, Ommen SR, Tajik AJ, *et al.* Sarcomeric genotyping in hypertrophic cardiomyopathy. *Mayo Clin Proc* 2005;80:463–9.
- Gupta MK, Robbins J. Post-translational control of cardiac hemodynamics through myosin binding protein C. *Pflugers Arch* 2014;466:231–6.
- Razzaque MA, Gupta M, Osinska H, *et al.* An endogenously produced fragment of cardiac myosin-binding protein C is pathogenic and can lead to heart failure. *Circ Res* 2013;113:553–61.
- Kaier TE, Twerenbold R, Puelacher C, *et al.* Direct comparison of cardiac myosin-binding protein c with cardiac troponins for the early diagnosis of acute myocardial infarction. *Circulation* 2017;136:1495–508.
- Shah AS, Chin CW, Vassiliou V, *et al.* Left ventricular hypertrophy with strain and aortic stenosis. *Circulation* 2014;130:1607–16.
- Chin CW, Messika-Zeitoun D, Shah AS, *et al.* A clinical risk score of myocardial fibrosis predicts adverse outcomes in aortic stenosis. *Eur Heart J* 2016;37:713–23.
- Chin CW, Semple S, Malley T, *et al.* Optimization and comparison of myocardial T1 techniques at 3T in patients with aortic stenosis. *Eur Heart J Cardiovasc Imaging* 2014;15:556–65.
- Singh A, Horsfield MA, Bekele S, *et al.* Myocardial T1 and extracellular volume fraction measurement in asymptomatic patients with aortic stenosis: reproducibility and comparison with age-matched controls. *Eur Heart J Cardiovasc Imaging* 2015;16:763–70.
- Baumgartner H, Hung J, Bermejo J, *et al.* Echocardiographic assessment of valve stenosis: EAE/ASE recommendations for clinical practice. *Eur J Echocardiogr* 2009;10:1–25.
- Cowell SJ, Newby DE, Prescott RJ, *et al.* A randomized trial of intensive lipid-lowering therapy in calcific aortic stenosis. *N Engl J Med* 2005;352:2389–97.
- Houslay ES, Cowell SJ, Prescott RJ, *et al.* Progressive coronary calcification despite intensive lipid-lowering treatment: a randomised controlled trial. *Heart* 2006;92:1207–12.
- O'Hanlon R, Grasso A, Roughton M, *et al.* Prognostic significance of myocardial fibrosis in hypertrophic cardiomyopathy. *J Am Coll Cardiol* 2010;56:867–74.
- Vigliano CA, Cabeza Meckert PM, Diez M, *et al.* Cardiomyocyte hypertrophy, oncosis, and autophagic vacuolization predict mortality in idiopathic dilated cardiomyopathy with advanced heart failure. *J Am Coll Cardiol* 2011;57:1523–31.
- Saito T, Hojo Y, Hirose M, *et al.* High-sensitivity troponin T is a prognostic marker for patients with aortic stenosis after valve replacement surgery. *J Cardiol* 2013;61:342–7.
- Kupari M, Eriksson S, Turto H, *et al.* Leakage of cardiac troponin I in aortic valve stenosis. *J Intern Med* 2005;258:231–7.
- Quarto C, Dweck MR, Murigu T, *et al.* Late gadolinium enhancement as a potential marker of increased perioperative risk in aortic valve replacement. *Interact Cardiovasc Thorac Surg* 2012;15:45–50.
- Treibel TA, López B, González A, *et al.* Reappraising myocardial fibrosis in severe aortic stenosis: an invasive and non-invasive study in 133 patients. *Eur Heart J* 2017;12:1.
- Apple FS, Collinson PO. IFCC Task force on clinical applications of cardiac biomarkers. Analytical characteristics of high-sensitivity cardiac troponin assays. *Clinical Chemistry* 2012;58:54–61.