CORRESPONDENCE

Response to: 'Congenital tricuspid valve disease can masquerade as primary idiopathic TR' by Rajpal

We received the letter by Dr Rajpal addressing the spectrum of congenital tricuspid valve (TV) abnormalities and highlighting congenital tricuspid disease as an under-recognised cause of isolated tricuspid regurgitation (TR) in adults. We agree wholeheartedly that there is a wide spectrum of congenital tricuspid valve disease, ranging from Ebstein's anomaly to tricuspid valve dysplasia which encompasses both primary leaflet malformations and chordal abnormalities. While Ebstein's anomaly is a well-recognised cause of primary tricuspid regurgitation, isolated tricuspid valve dysplasia is less frequently encountered in adults without other associated congenital abnormalities.¹ The rarity of isolated TV dysplasia, coupled with the technical difficulties of imaging the tricuspid valve, makes this a challenging diagnosis. In

our own practice we have encountered several patients with severe isolated TR who were referred to surgery and were found intra-operatively to have previously unrecognised leaflet dysplasia. A low index of suspicion must be maintained in relatively young adults presenting with isolated tricuspid regurgitation in the absence of other potential causes, particularly as surgery has proven efficacy in treating these patients.^{2,3}

Erin Amanda Fender, Chad J Zack, Rick A Nishimura

Mayo Clinic, Rochester, Minnesota, USA

Correspondence to Dr Rick A Nishimura, Mayo Clinic, Rochester MN 55905, USA; nishimura.rick@mayo.edu

Competing interests None declared.

Provenance and peer review Not commissioned; internally peer reviewed.

OPEN ACCESS

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work

non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

To cite Fender EA, Zack CJ, Nishimura RA. *Heart* 2018;**104**:1308.

Published Online First 24 February 2018

Heart 2018;**104**:1308.

doi:10.1136/heartjnl-2018-313078

REFERENCES

- 1 Fender EA, Zack CJ, Nishimura RA. Isolated tricuspid regurgitation: outcomes and therapeutic interventions. *Heart* 2018;**104**:798–806.
- 2 Said SM, Dearani JA, Burkhart HM, *et al*. Management of tricuspid regurgitation in congenital heart disease: is survival better with valve repair? *J Thorac Cardiovasc Surg* 2014;**147**:412–9.
- 3 Nishimura RA, Otto CM, Bonow RO, *et al*. ACC/AHA Task Force Members. 2014 AHA/ACC Guideline for the Management of Patients With Valvular Heart Disease: executive summary: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation* 2014;**129**:2440–92.