

12

ANTI-CANCER TYROSINE KINASE INHIBITORS INCREASE OXIDATIVE STRESS IN PRIMARY CARDIAC FIBROBLASTS

CJ McMullen, C McCluskey, SJ Kim, S Laovithayanggoon, M MacDonald, M Safar, R Wood, MR Cunningham, S Currie. *Strathclyde Institute of Pharmacy and Biomedical Sciences, The University of Strathclyde, Glasgow, UK*

10.1136/heartjnl-2018-SCF.22

Anti-cancer tyrosine kinase inhibitors (TKI) are known to exert cardiotoxicity that can result in heart failure. Here we have used primary adult rat cardiac fibroblasts (CFs) to examine whether imatinib and sunitinib (both TKIs) increase oxidative stress and cause increased oxidation and activation of calcium/calmodulin dependent protein kinase II (CaMKII). Activation of CaMKII is known to be important for cellular function but excessive activation is linked to cardiac pathology.

CFs, isolated from adult Wistar rat hearts, were maintained in culture and treated over 24 hour with the TKIs sunitinib and imatinib (0.1–20 μ M) in serum-free medium (DMEM). Cell phenotype and growth was monitored using a Nikon Eclipse (TE300) inverted microscope and ROS assays performed using a specially designed assay kit (Abcam, UK) which uses DCFDA to measure hydroxyl, peroxy and other ROS activity. Quantitative immunoblotting was used to assess ox-CaMKII protein levels.

Treatment with both TKIs caused a dose-dependent decrease in cell viability. Sunitinib exerted more toxic effects at lower concentrations than imatinib. Decreased cell viability was evident at concentrations as low as 1 μ M for sunitinib with dramatic cell loss (>50%) evident at 10–20 μ M. Significantly increased ROS levels (using the DCFDA-based assay) were observed in the presence of both imatinib (4956 \pm 102 vs 1984 \pm 358 (imatinib vs vehicle (a.u.), n=3, p<0.05) and sunitinib (15186 \pm 3856 vs 1984 \pm 358 (sunitinib vs vehicle (a.u.), n=3, p<0.01). Quantitative immunoblotting suggested that ox-CaMKII levels were increased following treatment with either 1 μ M imatinib (1.4-fold (ox-CaMKII/GAPDH)) or 1 μ M sunitinib (1.2-fold (ox-CaMKII/GAPDH)) however at higher concentrations (particularly for sunitinib) increased cell loss and resultant loss of total protein was observed.

In conclusion, both imatinib and sunitinib cause increased oxidative stress in adult primary CFs with resultant activation of CaMKII. These effects are evident at lower concentrations of sunitinib than for imatinib and reflect the increased cardiotoxicity of this TKI reported in the patient.

13

STEM-CELL DERIVED MYOGENIC PROGENY ENRICH FOR VASCULAR SMOOTH MUSCLE CELL EPIGENETIC MARKS AT THE MYOSIN HEAVY CHAIN 11 PROMOTER *IN VITRO*

¹Roya Kakimjavadi, ¹Denise Burtenshaw, ¹Emma Fitzpatrick, ¹Mariana Di Luca, ¹Eoin Corcoran, ¹Gillian Casey, ²Eileen M Redmond, ¹Paul A Cahill. ¹*Vascular Biology and Therapeutics Laboratory, School of Biotechnology Faculty of Science and Health, Dublin City University, Dublin 9, Ireland;* ²*Department of Surgery, University of Rochester Medical Centre, Rochester, NY, USA*

10.1136/heartjnl-2018-SCF.23

Epigenetics plays a critical role in SMCs differentiation where histone proteins associated with the myosin heavy chain 11 (*Myh11*) promoter are post-translationally modified while promoter sequences are de-methylated by the DNA-modifying

enzyme ten-eleven translocation-2 (TET2). Di-methylation of lysine 4 on histone 3 (H3K4me2) is generally associated with euchromatin and active gene expression whereas, tri-methylation of lysine 27 on histone 3 (H3K27me3) is generally associated with formation of heterochromatin and gene suppression. Our aim was to determine the level of H3K4me2 and H3K27me3 enrichment on *Myh11* promoter of differentiated SMCs and compare that signature to undifferentiated resident vascular stem cells and their stem cell-derived myogenic progeny. Murine Sca1⁺ stem cells (APCs), rat medial Sox10⁺ multipotent vascular stem cells (MVSCs) and CD44⁺ bone-marrow derived mesenchymal stem cells (MSCs) were all examined for H3K4me2 and H3K27me3 enrichment on *Myh11* promoter, before and after myogenic differentiation and compared to freshly isolated aortic differentiated SMCs and sub-cultured 'de-differentiated SMCs' *in vitro* by Chromatin Immunoprecipitation (ChIP) assay. Differentiated rat and murine SMCs were highly enriched for H3K4me2 mark at the *Myh11* promoter compared to the H3K27me3 mark, concomitant with enhanced TET-2 binding. Phenotypical 'de-differentiated' sub-cultured rat and murine SMC were also enriched for H3K4Me2 mark when compared to H3K27me3, but to a much lesser extent. In contrast, all three stem cell populations were highly enriched for H3K27me3. The levels of enrichment of H3K27me3 significantly reduced following myogenic differentiation with TGF- β 1 concomitant with a significant enrichment of H3K4me2 mark and TET-2 binding to levels that mimicked the level of enrichment in de-differentiated SMCs at the same locus. We conclude that stem-cell derived myogenic progeny enrich for the H3K4me2 mark at the *Myh11* promoter and thus may be responsible for the presence of this mark within arteriosclerotic cells following injury.

14

GENERATION OF χ -SECRETASE INHIBITOR-LOADED PLGA-Fe₃O₄ MAGNETIC NANOPARTICLES

Roa Bashmail, Niamh McKenna, Ciaran O'Shea, Roya Hakimjavadi, Claire Molony, Dorota Kozłowska, Paul A Cahill. *Vascular Biology and Therapeutics Laboratory, School of Biotechnology Faculty of Science and Health*

10.1136/heartjnl-2018-SCF.24

Stem cell derived myogenic progeny play important roles in the pathophysiological processes of various vascular disorders, such as arteriosclerosis, atherosclerosis, and in-stent restenosis. Targeting these cells is an attractive therapeutic strategy for treating vascular remodelling. However, while polymer-coated DES have significantly reduced the incident of in-stent restenosis, current DESs lack the fundamental capacity for (i) adjustment of the drug dose and release kinetics and the (ii) ability to replenish the stent with a new drug on depletion. This limitation can be overcome by a strategy combining magnetic targeting via a uniform field-induced magnetization effect and a biocompatible magnetic nanoparticle (MNP) formulation designed for efficient entrapment and delivery of specific drugs that target the resident vascular stem cell source of the SMC. Magnetic nanoparticles (MNP's) containing magnetite (Fe₃O₄) were fabricated, polymer coated with poly (DL-lactide-co-glycolide) polyvinyl alcohol [PLGA-PVA] and loaded with a χ -secretase inhibitor (GSI) of Notch signalling, DAPT using an oil in water emulsification technique. The free GSI's and GSI-loaded MNP's were assessed for drug release, the efficacy at