



OPEN ACCESS

ORIGINAL RESEARCH ARTICLE

Valve area and the risk of overestimating aortic stenosis

Ana González-Mansilla,^{1,2,3,4} Pablo Martínez-Legazpi,^{1,2,3,4} Andrea Prieto,^{1,2,3,4} Elena Gomá,^{1,2,3,4} Pilar Haurigot,^{1,2,3,4} Candelas Pérez del Villar,^{1,2,3,4} Víctor Cuadrado,⁵ Antonia Delgado-Montero,^{1,2,3,4} Raquel Prieto,^{1,2,3,4} Teresa Mombiela,^{1,2,3,4} Esther Pérez-David,^{1,2,3,4} Elena Rodríguez González,^{1,2,3,4} Yolanda Benito,^{1,2,3,4} Raquel Yotti,^{1,2,3,4} Manuel Pérez-Vallina,⁵ Francisco Fernández-Avilés,^{1,2,3,4} Javier Bermejo^{1,2,3,4}

¹Department of Cardiology, Hospital General Universitario Gregorio Marañón, Madrid, Spain

²Facultad de Medicina, Universidad Complutense de Madrid

³Instituto de Investigación Sanitaria Gregorio Marañón

⁴CIBERCV

⁵Information and Communications Technology Department, Hospital General Universitario Gregorio Marañón, Madrid, Spain

Correspondence to

Dr Javier Bermejo, Department of Cardiology, Hospital General Universitario Gregorio Marañón, Madrid, Spain; javier.bermejo@salud.madrid.org

AG-M and PM-L contributed equally.

AP and EG contributed equally.

Received 15 November 2018

Revised 15 January 2019

Accepted 21 January 2019

Published Online First

16 February 2019

ABSTRACT

Objective To obtain reference values of aortic valve area (AVA) in a large population and to infer the risk of overestimating aortic stenosis (AS) when focusing on flow-corrected indices of severity.

Methods We prospectively measured indices of AS in all consecutive echocardiograms performed in a large referral cardiac imaging laboratory for 1 year. We specifically analysed the distribution of AVA, indexed AVA and velocity ratio (Vratio) in patients with and without AS, the latter defined as the coexistence of valvular outflow obstruction (Vmax ≥ 2.5 m/s) and morphological findings of valve degeneration.

Results 16 156 echocardiograms were analysed, 14 669 of which did not show valvular obstruction (peak jet velocity < 2.5 m/s). In the latter group, AVA was $2.6 \pm 0.7 \text{ cm}^2$ in 8190 studies with normal valves and $2.3 \pm 0.7 \text{ cm}^2$ in 6479 studies with aortic sclerosis (AScl). There was a relatively wide overlap between values of AVA, indexed AVA and velocity ratio between studies of patients with AScl and AS. Values of AVA $\leq 1.0 \text{ cm}^2$ were found in 0.5% of studies with normal valves and 1.8% of studies with AScl. These proportions were 3.1% and 9.3% for AVA $\leq 1.5 \text{ cm}^2$, respectively. Vratio ≤ 0.25 were found in 0.1% of patients without obstruction. Risk factors for a small AVA in patients without obstruction were AScl, female sex, small body surface area, low ejection fraction and mitral regurgitation.

Conclusions Normal values of continuity-equation derived AVA are smaller than previously considered. AVA values below cutoffs of moderate or severe AS can be found in patients without the disease. Flow-corrected indices may overestimate AS in patients with low gradients, particularly in the presence of well-identified risk factors.

severity may be a more difficult task. When Vmax, mean pressure gradient (MPG) and effective aortic valve area (AVA) show consistent values below or beyond established cutoffs, grading is simple.² However, in >30% of patients, these three indices yield discordant grades of severity.³ Reasons for discrepancy may be measurement errors, abnormal flow states, inconsistent thresholds or a combination of these.^{1–5} In most cases of discrepancy, an AVA $\leq 1 \text{ cm}^2$ prevails over low values of MPG and Vmax to define AS as 'severe' and guide clinical decisions. Thus, AVA is frequently considered the most important index to grade AS. Although reference values have never been obtained in a large population, the area of the normal aortic valve is believed to be 3–4 cm^2 ,⁶ and values $< 2.0 \text{ cm}^2$ are believed to cause VOO. We hypothesised that exploring AVA values in patients without VOO would be useful to unmask the risk of overestimating AS in patients who do suffer the disease. We designed a prospective study to measure AVA in a large series of consecutive echocardiographic examinations focusing on patients without VOO. Additional aims were (1) to identify the risk factors related to a small AVA in patients without VOO, and (2) to address whether alternative flow-corrected metrics such as the body-surface indexed AVA (AVAi) and the velocity ratio (Vratio or dimensionless index) are more specific to identify patients with VOO.

METHODS

Study design

We obtained quantitative AS data in a large series of unselected consecutive echocardiograms from a tertiary referral hospital. From 1 July 2015 to 30 June 2016, sonographers and physicians of the adult echocardiography laboratory from our institution were instructed to obtain measurements of AS in all transthoracic studies, irrespective of their indication. All data were analysed anonymously, and the study was performed following current recommendations of the Declaration of Helsinki. The study was approved by the Institutional Ethics Committee which exempted the need for patient informed consent, as it required no modification in standard clinical practice.



► <http://dx.doi.org/10.1136/heartjnl-2019-314734>



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: González-Mansilla A, Martínez-Legazpi P, Prieto A, et al. Heart 2019;105:911–919.




Figure 1 Study population.

Image acquisition and analysis

Ultrasound recordings were obtained according to current guidelines.² Special care was taken to obtain left ventricular outflow tract (LVOT) images from parasternal long-axis view using a zoom to identify the clearest and largest LVOT annulus recordings. Doppler spectrograms were obtained from apical five-chamber views, using multifrequency transducers and colour Doppler, to optimise alignment with the flow. The pulsed-wave Doppler sample volume was located 0.5 to 1 cm proximal to the aortic valve plane. All measurements were performed using EchoPac (BT08; GE Healthcare). LVOT diameter was measured from mid systolic frames at the hinge point of insertion of the aortic cusps from inner-to-inner edges.² Stroke volume (SV) was calculated as the product of LVOT cross-sectional area (derived from the former diameter assuming a circular shape) and the LVOT time-velocity integral (TVI). AVA was calculated as SV/aortic TVI; AVAi was calculated as AVA/patient's body surface area (Du Bois and Du Bois' formula). The Vratio was calculated as LVOT TVI/Aortic TVI. Measurements were averaged from three consecutive beats in regular (sinus or paced) rhythms and five beats in patients with atrial fibrillation. Image quality was graded as 0 (reliable Doppler measurements unavailable), 1 (poor image quality but obtainable measurements), 2 and 3 (good and excellent image quality, respectively). Studies with 0 image quality or performed by in-training personnel were dismissed. Cardiologists with level-III expertise in cardiac ultrasound read and signed reports from all examinations and re-measured all studies showing an AVA ≤ 1.5 cm².

VOO was defined as Vmax ≥ 2.5 m/s.² AS was defined as morphological findings of valve degeneration (B-mode signs of valve fibrosis and/or calcification) and the presence of VOO.² Aortic valves from studies without VOO were classified either as normal or as aortic sclerosis (AScl) based on the absence or the presence of valve degeneration.⁷ Studies with prosthetic aortic valves or with subvalvular (late-peaking LVOT velocity ≥ 1.8 m/s) or supravalvular obstruction (anatomical signs of supravalvular stenosis and Vmax ≥ 2.0 m/s) were excluded from analysis. Custom software written in R was used to forward data from the clinical report system to an anonymised database without any user interaction. Reproducibility of AVA measurements in patients with AS in our laboratory has been previously reported.⁸ However, because reproducibility may vary in other populations, we underwent a measurement reproducibility analysis in 191 randomly selected examinations without VOO (two level-III blinded cardiologists, 3 weeks apart; online supplementary table 1).

Statistical analysis

Data are presented as mean ± SD deviation except otherwise indicated. One-way analyses of variance (followed by Tukey contrasts) and χ² (followed by Holm-adjusted Fisher pairwise contrasts) tests were used to compare studies with normal, sclerotic or stenotic aortic valves. Intraclass correlation and Bland-Altman analyses were used for assessing agreement. Empirical, normal and empirical cumulative distribution functions were calculated for all AS indices. The 95% CI of the latter were calculated using the Kolmogorov-Smirnov's D statistic. Bivariate density plots were used to visualise the overlap between quantitative variables. False positive rates for identifying AS in patients without VOO were calculated for different cutoffs of severity, and their 95% CIs were computed assuming a binomial distribution. Uni and multivariate logistic regression analyses were used to assess the factors related to small AVA values in patients without VOO. For multivariate modelling, the full study database was randomly split into learning (n=9656 studies) and testing (n=4974) independent datasets. Variable selection was performed by backward variable selection based on the Alkaike's information criterion, entering clinically relevant variables and using a cross-validation algorithm to avoid overfitting. Beta coefficients, ORs (and their 95% CIs) and diagnostic cutoff point (based on Youden's J statistic) of this model were calculated. Its predictive capability was then assessed on the testing dataset by means of the ROC curve, its area under the curve (AUC), 95% CI, as well as its sensitivity, specificity, positive and negative predictive values (based on 2000 bootstrap replicates). Due to their derivation formulae, there is inherent strong collinearity between AS indices, SV and the aortic TVI. Therefore, the full multivariate modelling strategy was performed twice, with and without entering these factors. All analyses were performed in R (V3.4.0). P values <0.05 were considered significant.

RESULTS

Study population

From an original pool of 29 514 studies, 16 156 echocardiograms from 13 920 patients were finally analysed (figure 1). No VOO was found in 14 669 (8 190 and 6 479 showing normal and sclerotic valves, respectively), whereas AS was found in 1 487. Demographic, clinical and echocardiographic data are shown in table 1 and figure 2.

Table 1 Clinical and echocardiographic data of the studied population

	No outflow obstruction		Outflow obstruction		P value
	Normal	Aortic valve sclerosis	Aortic valve stenosis	Total	
Studies, n	8190	6479	1487	16156	
Patients, n	7055	5643	1222	13920	
Age, years	57±18	77±12*	79±12*†	66±19	<0.001
Sex, male n (%)	3046 (52)	2330 (50)	490 (47)†	5866 (51)	0.01
Body surface area, m²	1.81±0.21	1.76±0.21*	1.75±0.19*	1.79±0.21	<0.001
Sinus rhythm, n (%)	7269 (92)	4512 (73)*	1019 (72)*	12 800 (83)	<0.001
Study indication, n (%)					<0.001
Known or suspected heart failure	2713 (25)	1845 (23)*	213 (14)*†	4771 (23)	
Known or suspected rhythm disturbance	1271 (12)	492 (6)*	24 (2)*†	1787 (9)	
Known or suspected ischaemic heart disease	618 (6)	271 (3)*	20 (1)*†	909 (4)	
Known or suspected cardioembolism	317 (3)	428 (5)*	23 (2)*†	768 (4)	
Systemic hypertension	386 (4)	232 (3)*	15 (1)*†	633 (3)	
Known or suspected valvular heart disease	62 (1)	101 (1)*	32 (2)*	195 (1)	
Pulmonary hypertension	63 (1)	65 (1)	14 (1)	142 (1)	
Other/Unspecified	5451 (50)	4681 (58)*	1147 (77)*†	11 279 (55)	
LV septal thickness, cm	1.0±0.2	1.0±0.2*	1.2±0.2*†	1.0±0.2	<0.001
Left atrial diameter, cm	3.5±0.6	3.9±0.8*	4.2±0.8*†	3.7±0.8	<0.001
LV end diastolic volume, mL	109±47	125±56*	135±57*	117±52	<0.001
LV end systolic volume, mL	53±37	71±47*	72±43*	61±43	<0.001
LV ejection fraction, %	59±9	55±11*	56±9*†	57±10	<0.001
LV mass index, g/m²	81±22	91±34*	104±50*†	86±29	<0.001
LVOT diameter, cm	2.1±0.2	2.1±0.2	2.1±0.2	2.1±0.2	0.55
LVOT time–velocity integral, cm	21±5	21±6	22±6*†	22±6	<0.001
Stroke volume, mL	69±20	69±20	73±25*†	69±22	<0.001
Peak aortic jet velocity, cm/s	122±46	123±61	337±84*†	142±84	<0.001
Aortic time–velocity integral, cm	28±7	31±8*	75±23*†	33±17	<0.001
Aortic regurgitation, n (%)					<0.001
0–I	8025 (98)	5822 (90)*	1271 (85)*†		
II	96 (1)	427 (7)*	89 (6)*		
III	47 (1)	153 (2)*	21 (1)*†		
IV	22 (0)	77 (1)*	107 (7)*†		
Mitral regurgitation, n (%)					<0.001
0–I	5957 (73)	3204 (49)*	669 (45)*†	9830 (61)	
II	1818 (22)	2144 (33)*	430 (29)*†	4392 (27)	
III	337 (4)	865 (13)*	223 (15)*	1425 (9)	
IV	78 (1)	266 (4)*	165 (11)*†	509 (3)	
E wave velocity, cm/s	75±24	81±34*	95±45*†	78±31	<0.001
A wave velocity, cm/s	70±22	85±26*	100±35*†	78±27	<0.001
E/e' ratio	5±3	8±4*	10±5*†	7±3	<0.001
Systolic pulmonary artery pressure, mm Hg	30±12	38±15*	43±16*†	36±15	<0.001

Data presented as mean±SD, except otherwise indicated. P values obtained by one-way analysis of variance or χ^2 tests, where appropriate. The bold text in this table refers to categories.

*P<0.05 vs normal valves.

†P<0.05 vs sclerotic valves by Tukey or Fisher exact (Holm correction) pairwise contrasts, where appropriate.

LV, left ventricle; LVOT, left ventricular outflow tract.

Indices of AS in patients with and without VOO

AVA was $2.6\pm0.7\text{ cm}^2$ in studies with normal valves and $2.3\pm0.7\text{ cm}^2$ in studies with AScl (table 2, figure 3 and online supplementary figure 1). AVAi and Vratio were $1.4\pm0.4\text{ cm}^2/\text{m}^2$ and 0.8 ± 0.1 , respectively, in studies with normal valves, and $1.3\pm0.4\text{ cm}^2/\text{m}^2$ and 0.7 ± 0.2 in studies with AScl. These reference values of AVA, AVAi and Vratio remained almost unchanged when subgroups of studies with low LV ejection fraction, significant regurgitation, irregular rhythm or poor image quality (grade 1) were excluded (table 2). Values of AVA, AVAi and Vratio overlapped between studies with AScl and AS (figure 3

and online supplementary figure 1). AVA values $\leq1.5\text{ cm}^2$ were found in 5.8% (95% CI 5.5% to 6.2%) of studies without VOO (table 2). A total 0.5% (0.3 to 0.6) of studies with normal valves showed an AVA $\leq1.0\text{ cm}^2$ and 3.1% (2.7 to 3.5) showed an AVA $\leq1.5\text{ cm}^2$. In patients with AScl, these proportions were 1.8% (1.5 to 2.1) and 9.3% (8.6 to 10.0), respectively. Values of Vratio ≤0.5 were found in 3.6% (95% CI 3.2% to 4.0%) of studies with normal valves and in 14.6% (95% CI 13.7% to 15.4%) of studies with sclerotic valves. Only 0.1% of the studies without VOO showed a Vratio ≤0.25 (table 2).




Figure 2 Echocardiographic data. Boxplots of the echocardiographic data of patients with normal (green), sclerotic (blue) and stenotic valves (red). Panel A: LVOT diameter (cm). Panel B: LVOT TVI (cm). Panel C: peak aortic jet velocity (cm/s). Panel D: aortic TVI (cm). LVOT, left ventricular outflow tract; TVI, time–velocity integral.

Predictors of small AVA in the absence of outflow obstruction
 Studies without VOO and $\text{AVA} \leq 1.5 \text{ cm}^2$ showed smaller LVOT diameters, smaller LVOT TVIs, smaller SVi and larger aortic TVIs than those with $\text{AVA} > 1.5 \text{ cm}^2$ (table 3 and online supplementary figure 2). Risk factors for an $\text{AVA} \leq 1.5 \text{ cm}^2$ in the absence of VOO by multivariate analysis were: female sex, AScl, low LV ejection fraction, age, a small body surface and the degree of mitral regurgitation (figure 4). Overall predictive capability of the model was good, as demonstrated in the independent validation dataset (figure 4 Insert): $\text{AUC}=0.75$ (95% CI 0.73 to 0.78), specificity=0.92 (0.90 to 0.93), sensitivity=0.99 (0.91 to 1.00), positive predictive value=0.42 (0.39 to 0.45) and negative predictive value=1.00 (1.00 to 1.00). When SV and aortic TVI were forced in the multivariate model, mitral regurgitation and body surface area were excluded as independent predictors (online supplementary table 2). Risk factors for an $\text{AVA} \leq 1.5 \text{ cm}^2$ were also predictors of an $\text{AVAi} \leq 0.85 \text{ cm}^2/\text{m}^2$ and a $\text{Vratio} < 0.5$ in the absence of outflow obstruction (online supplementary table 2). Representative examples of studies with a small AVA in the absence of AS are shown in figure 5 and supplemental videos 1–3.

DISCUSSION

To our knowledge, this is the first study providing reference values for AVA, AVAi and Vratio. ‘Normal’ AVA values obtained by the continuity equation are in average 1 cm^2 smaller than usually described.⁶ By focusing on echocardiographic studies from patients without the disease, we show that flow-corrected indices may overestimate AS. Importantly, AVA values consistent with moderate or severe AS can be identified in as many of 3% of patients with normal valves and 9% of patients with AScl. These patients were not diagnosed of AS because their Vmax was $< 2.5 \text{ m/s}$. However, if eventually Vmax were to increase beyond this threshold, these patients would automatically shift

to a moderate or severe (low gradient) AS category based on AVA. Thus, AVA poorly accounts for mild degrees of AS and the cut-off criteria established for moderate and severe AS may overestimate the real haemodynamic relevance of the disease. The Vmax and the pressure gradient should not be overlooked to grade AS, particularly in the less severe degrees of severity.

Haemodynamic criteria to grade AS

The unreliability of focusing excessively on AVA to grade AS has been recently emphasised,⁹ based on the results of a large retrospective analysis of patients with the disease.¹⁰ The present study focuses on patients without the disease to provide further insight into the limitations of flow-corrected indices obtained by Doppler echocardiography. Beyond measurement errors, several factors may account for the observed small values of flow-corrected indices observed even in the absence of VOO. AVA, AVAi and Vratio rely on several physical assumptions which have not been fully validated in the absence of VOO. The peak velocity—as measured using continuous-wave Doppler—overestimates average velocity along the full cross-section of the transvalvular jet if its flow profile is not completely flat. By this and other mechanisms, flow-dependence^{8 11} and pressure recovery¹² may sometimes yield misleadingly small AVAs. An alternative interpretation of our findings is that the $\text{Vmax} > 2.5 \text{ m/s}$ criterion lacks sensitivity for the diagnosis of VOO, all patients with AScl and small AVA values suffering AS irrespective of their Vmax values. However, from a physiological perspective, a haemodynamically significant outflow obstruction that generates a relevant convective acceleration is highly unlikely with lower transvalvular velocities.^{13–15}

Because the most challenging grading situation is PLGAS, current guidelines emphasise combining several techniques in this scenario.² Cross-sectional ultrasound and CT must ensure calcification and limited leaflet excursion to grade AS as severe, but there is a high overlap between the morphology of sclerotic and stenotic valves. Cardiac catheterisation is useful to confirm haemodynamic measurements, but AVA/MPG discrepancies are also common using this technique.⁵ Finally, concurrent abnormalities in the vascular tree may be responsible for the increased afterload to the LV in addition to AS.^{11 16} Thus, despite this large armamentarium, decision-making in low-gradient AS remains difficult in some patients. The present study shows that the major source of small AVAs in patients without AS is a low SV. Hence, misclassification of AS will most frequently combine a small AVA, a low gradient and a low-flow situation. Furthermore, clinical conditions previously described as highly prevalent in PLGAS¹⁷—female sex, age and low body surface area—are identified in our study as the main risk factors for showing small AVAs in patients without VOO.

A post hoc analysis of a prospective clinical trial has shown that outcomes of patients with AS are best predicted in terms of their MPG.¹⁸ PLGAS may therefore represent a milder degree of severity than high gradient AS, particularly when transvalvular flow is normal.¹⁹ Nevertheless, several observational studies have shown that outcomes of symptomatic patients are improved by valve replacement in low-flow PLGAS.^{17 20–22} Although PLGAS is currently a IIa-level recommendation of valve replacement,¹⁴ the benefit of therapy is clearly lower than in high-gradient AS. In fact, a low MPG at the time of treatment is a consistent predictor of lack of clinical improvement after valve replacement.^{23 24} The limited specificity of current grading criteria could be responsible for driving some patients with milder degrees of AS towards valve replacement. Our findings emphasise that special care needs to be taken to follow

Table 2 Summary statistics of indices of AS in patients without valvular outflow obstruction

	Normal aortic valve			Aortic valve sclerosis						
	All studies	LVEF >35%	MR ≤2	Regular rhythm	Image quality ≥2	All studies	LVEF >35%	MR ≤2	Regular rhythm	Image quality ≥2
Studies, n	8190	7685	7773	7735	7012	6479	5756	5319	5032	4997
AVAi (cm^2/m^2), mean±SD	2.6±0.7	2.6±0.7	2.6±0.7	2.6±0.7	2.5±0.7	2.3±0.7	2.3±0.7	2.4±0.7	2.4±0.7	2.3±0.7
AVAi ≤1.5 cm^2/m^2 , n (%)	256 (3.1)	219 (2.8)	219 (2.8)	216 (2.8)	217 (3.1)	600 (9.3)	495 (8.6)	425 (8.0)	350 (7.0)	504 (10.1)
(95% CI)	(2.7 to 3.5)	(2.5 to 3.2)	(2.4 to 3.2)	(2.4 to 3.2)	(2.7 to 3.5)	(8.6 to 10.0)	(7.9 to 9.3)	(7.3 to 8.7)	(6.3 to 7.7)	(9.3 to 10.9)
AVAi ≤1 cm^2/m^2 , n (%)	38 (0.5)	25 (0.3)	31 (0.4)	32 (0.4)	32 (0.5)	117 (1.8)	87 (1.5)	79 (1.5)	52 (1.0)	98 (2.0)
(95% CI)	(0.3 to 0.6)	(0.2 to 0.5)	(0.3 to 0.5)	(0.3 to 0.6)	(0.3 to 0.6)	(1.5 to 2.1)	(1.2 to 1.8)	(1.2 to 1.8)	(0.8 to 1.3)	(1.6 to 2.3)
AVAi (cm^2/m^2), mean±SD	1.4±0.4	1.4±0.4	1.4±0.4	1.4±0.4	1.4±0.4	1.3±0.4	1.3±0.4	1.3±0.4	1.3±0.4	1.4±0.4
AVAi ≤0.85 cm^2/m^2 , n (%)	332 (4.1)	289 (3.8)	287 (3.7)	294 (3.8)	285 (4.1)	683 (10.6)	545 (9.5)	492 (9.2)	410 (8.1)	561 (11.2)
(95% CI)	(3.6 to 4.5)	(3.3 to 4.2)	(3.3 to 4.1)	(3.4 to 4.2)	(3.6 to 4.5)	(9.6 to 11.4)	(8.5 to 10.0)	(7.4 to 8.9)	(7.4 to 8.9)	(10.4 to 12.1)
AVAi ≤0.6 cm^2/m^2 , n (%)	37 (0.5)	32 (0.4)	30 (0.4)	32 (0.4)	32 (0.5)	101 (1.5)	102 (1.9)	119 (2.4)	119 (2.4)	168 (15.4)
(95% CI)	(0.3 to 0.6)	(0.3 to 0.6)	(0.2 to 0.5)	(0.3 to 0.6)	(0.3 to 0.6)	(1.45 to 2.3)	(1.45 to 2.3)	(1.5 to 2.2)	(1.5 to 2.2)	(2.0 to 2.8)
Vratio, mean±SD	0.8±0.1	0.8±0.1	0.8±0.1	0.8±0.1	0.8±0.1	0.8±0.1	0.8±0.1	0.7±0.2	0.7±0.2	0.7±0.2
Vratio ≤0.50, n (%)	294 (3.6)	229 (3.0)	246 (3.2)	252 (3.3)	254 (3.6)	939 (14.6)	740 (12.9)	672 (12.6)	573 (11.4)	768 (15.4)
(95% CI)	(3.2 to 4.0)	(2.6 to 3.4)	(2.8 to 3.6)	(2.9 to 3.7)	(3.2 to 4.1)	(13.7 to 15.4)	(12.0 to 13.7)	(10.5 to 12.3)	(10.5 to 12.3)	(11.1 to 13.5)
Vratio ≤0.25, n (%)	11 (0.1)	3 (0.0)	9 (0.1)	8 (0.1)	10 (0.1)	8 (0.1)	5 (0.1)	4 (0.1)	6 (0.1)	6 (0.1 to 0.2)
(95% CI)	(0.1 to 0.2)	(0.0 to 0.1)	(0.0 to 0.2)	(0.0 to 0.2)	(0.0 to 0.2)	(0.0 to 0.2)	(0.0 to 0.2)	(0.0 to 0.2)	(0.0 to 0.2)	(0.0 to 0.2)

Data presented as mean±SD, and number of patients and their proportion (%), and 95% CI showing values below the established cut-offs for moderate and severe AS.

AS, aortic valve stenosis; AVAi, indexed aortic valve area; LVEF, left ventricular ejection fraction; MR, mitral regurgitation; Vratio, velocity ratio.




Figure 3 The distribution of grading indices of AS. Empirical cumulative distribution functions for AVA, AVAi and Vratio. Vertical dotted lines account for current established cutoffs of AS. AS, aortic valve stenosis; AVAi, indexed aortic valve area; Vratio, velocity ratio.

current recommendations of an integrated stepwise approach in low-gradient AS.² The risk factors for misdiagnosis we have identified in the present study—small body surface area, female gender, advanced age, impaired systolic function and significant mitral regurgitation—raise a call for reassessment before indicating valve replacement in these situations.

Our study suggests that there is a wide continuum in the natural history of aortic valve degeneration spanning from a normal aortic valve to severe high-gradient AS. How early in this process patients may benefit from valve replacement is currently a matter of reassessment, particularly considering that the risks of percutaneous therapy are becoming lower. Meanwhile,

Table 3 Clinical and echocardiographic data of studies without valvular outflow obstruction according to AVA > or ≤ 1.5 cm²

	AVA > 1.5 cm ²	AVA ≤ 1.5 cm ²	P value
Patients, n	11 980	710	
Studies, n	13 774	856	
Aortic valve type, n (%)			<0.001
Normal	7932 (58)	256 (30)	
Sclerosis	5842 (42)	600 (70)	
Age, years	65±19	76±17	<0.001
Sex male, n (%)	6422 (54)	188 (27)	<0.001
Body surface area, m ²	1.79±0.21	1.63±0.18	<0.001
Peak aortic jet velocity, cm/s	139±32	169±34	<0.001
Stroke volume (mL)	71±21	41±11	<0.001
Aortic time–velocity integral, cm	29±7	34±9	<0.001
LVOT diameter, cm	2.1±0.2	1.9±0.2	<0.001
LVOT time–velocity integral, cm	21±5	18±5	<0.001
LV end diastolic volume, mL	114±50	126±57	0.02
LV end systolic volume, mL	59±41	76±51	<0.001
LV ejection fraction, %	58±10	54±12	<0.001
LV mass index, g/m ²	85±27	91±29	0.02
Relative wall thickness	0.4±0.1	0.5±0.2	0.18
Mitral regurgitation, n (%)			<0.001
0–I	8715 (63)	425 (50)	
II	3733 (27)	219 (26)	
III	1055 (8)	142 (17)	
IV	271 (2)	70 (8)	

Data presented as mean±SD, except otherwise indicated.

AVA, aortic valve area; LVOT, left ventricular outflow tract.

physicians must be aware that when examined individually, the contradictory information yield by AVA, Vmax and MPG is not exclusively caused by the haemodynamic definitions of AS. Outcome trials prospectively addressing the impact of therapy in earlier stages of the disease need to be performed, not only in terms of symptomatic status,²⁵ but also of lower degrees of haemodynamic significance.

Limitations

The scope of this work was to assess the limitations of conventional transthoracic echocardiography metrics of AS as currently recommended in clinical practice. A different reference method such as CT would have provided interesting results on studies showing small AVA values. A relatively high proportion of studies were unsuitable for analysis. However, the large volume of data allowed for highly powered statistical analyses, allowing for reliable index distribution estimates and narrow CIs. We measured the LVOT diameter at the annulus level instead of 0.5 to 1.0 cm proximal to the valvular plane (at the point where the pulsed-wave Doppler sample volume is placed in the LVOT).²⁶ Which location is best for calculating AVA is a matter of debate,^{2,27} but the funnel-shape of the LVOT is likely to yield lower values at the annulus.²⁶ In patients with AS, combining Doppler data with LVOT cross-sectional area measured by transoesophageal echocardiography²⁸ or CT²⁹ may increase the accuracy of AVA measurements and yields slightly higher values than those obtained exclusively by transthoracic echocardiography. However, outcome studies have not unequivocally demonstrated improved clinical efficacy of these hybrid methods,²⁹ and transthoracic echocardiography is still the reference technique for clinical decision-making.¹ We did not use dedicated continuous-wave (Pedoff) transducers, and we focused exclusively on




Figure 4 Predictors of small AVA values in studies without VOO. Multivariate logistic regression predictors of an AVA $\leq 1.5 \text{ cm}^2$ in studies without VOO. ORs and 95% CI and cutoffs are shown for the model obtained in the learning dataset ($n=9656$) without entering SV and aortic TVI in the model. Insert: ROC curves, cutoffs and AUC obtained in the testing dataset ($n=4974$) for this model (blue) and for the one described in the online supplementary table 1, obtained forcing SV and aortic TVI into the model (green). AUC, area under the curve; AVA, aortic valve area; ROC, receiver operating characteristic; TVI, time–velocity integral; VOO, valvular outflow obstruction.




Figure 5 Representative examples of small AVA values obtained in studies without VOO. The top row shows zoom images of the LVOT and measurements of the aortic annulus. The middle row shows pulsed-wave Doppler spectrograms of the LVOT velocities, whereas the bottom row shows continuous-wave Doppler spectrograms of the aortic valve jet. Results are displayed as measured beat / average results of 3–5 consecutive beats. Panel A and online supplemental video 1 are from a 69-year-old woman studied for suspected coronary artery disease. Her echocardiogram was normal except for an impaired relaxation filling pattern. The aortic valve was morphologically normal. Panel B and online supplemental video 2 are from a 67-year-old woman studied before chemotherapy. Her aortic valve was morphologically normal. Panel C and online supplemental video 3 are from a 75-year-old man with chronic liver disease. Main echocardiographic findings were normal except for a 2+ mitral regurgitation, a systolic pulmonary artery pressure of 40 mm Hg and a sclerotic aortic valve. AVA, aortic valve area; LVOT, left ventricular outflow tract; VOO, valvular outflow obstruction.

Valvular heart disease

the apical five-chamber view. However, reported values of AVA, AVAi and Vratio would have been even lower if these additional approaches had been used. Due to the moderate reproducibility of AS indices, the 'normal' values herein reported need to be interpreted with caution. However, reproducibility is as a limitation of the continuity equation method per se. Although AScI is an independent risk factor for future cardiovascular events in population studies, this risk is not associated with the progression to AS.⁷ Only prospective studies will clarify the clinical relevance of small AVA in patients with Vmax <2.5 m/s and help to define whether this threshold should be lowered in patients with low-flow states.

CONCLUSIONS

Normal values of continuity-equation derived AVA are smaller than previously considered. Values of AVA consistent with the definitions moderate or even severe AS can be found in patients without outflow obstruction. Until their prognostic significance is clarified, the isolated finding of a small AVA value needs to be interpreted with caution. Women with advanced age, a small body surface area, a low ejection fraction, AScI and/or significant degrees of mitral regurgitation are at the highest risk of overestimating AS when focusing excessively on flow-corrected indices.

Key messages

What is already known about this subject?

- Grading aortic valve stenosis is based on peak transvalvular jet velocity (Vmax), mean pressure gradient (MPG) and effective aortic valve area (AVA). In case of discrepancy, the flow-corrected criteria usually prevail to establish severity. Establishing the diagnosis of aortic valve stenosis (AS) requires the existence of valvular outflow obstruction, defined as a Vmax ≥ 2.5 m/s.

What does this study add?

- For the first time, continuity-equation-derived AVA normal values are estimated from a large population. Normal values of AVA are smaller than previously considered. We found that AVA values consistent with moderate or even severe AS can be identified in patients without the disease. AVA may overestimate the severity of AS, particularly in small and elderly women with low ejection fraction, concomitant mitral regurgitation and aortic sclerosis.

How might this impact on clinical practice?

- Small AVA values may overestimate AS. In the presence of low Vmax and MPG values, flow-corrected indices of AS must be interpreted with caution.

Acknowledgements The authors are in debt with all the personnel of the Laboratory of Cardiac Imaging from our institution for their contribution to this study.

Contributors All authors have approved this resubmission. All of the authors have made substantial contributions in the conception or design or analysis and interpretation of data. All the authors have participated drafting the manuscript or revising it critically for important intellectual content. All the authors have finally approved the submitted manuscript. JB, AG-M and PM-L: responsible for the overall content as guarantors.

Funding CPV has been funded by the Juan Rodés Fellowship (JR15/00039) (to CPV) from the Instituto de Salud Carlos III. PML was supported by the Juan de la Cierva Incorporación fellowship (IJCI-2014-19507) from the Ministerio de Economía y Competitividad, Spain and the CIBERCV.

Disclaimer The manuscript has not been published and is not being considered for publication elsewhere.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- 1 Baumgartner H, Falk V, Bax JJ, et al. ESC Scientific Document Group. 2017 ESC/EACTS Guidelines for the management of valvular heart disease. *Eur Heart J* 2017;38:2739–91.
- 2 Baumgartner H, Hung J, Bermejo J, et al. Recommendations on the echocardiographic assessment of aortic valve stenosis: a focused update from the European association of cardiovascular imaging and the American society of echocardiography. *Eur Heart J Cardiovasc Imaging* 2017;18:254–75.
- 3 Minners J, Allgeier M, Gohlke-Baerwolf C, et al. Inconsistencies of echocardiographic criteria for the grading of aortic valve stenosis. *Eur Heart J* 2008;29:1043–8.
- 4 Nishimura RA, Otto CM, Bonow RO, et al. 2017 AHA/ACC Focused Update of the 2014 AHA/ACC Guideline for the management of patients with valvular heart disease: a report of the american college of cardiology/american heart association task force on clinical practice guidelines. *J Am Coll Cardiol* 2017;70:252–89.
- 5 Minners J, Allgeier M, Gohlke-Baerwolf C, et al. Inconsistent grading of aortic valve stenosis by current guidelines: haemodynamic studies in patients with apparently normal left ventricular function. *Heart* 2010;96:1463–8.
- 6 Gaasch WH. Aortic valve area in aortic stenosis. 2016 <https://www.uptodate.com/contents/aortic-valve-area-in-aortic-stenosis>.
- 7 Otto CM, Lind BK, Kitzman DW, et al. Association of aortic-valve sclerosis with cardiovascular mortality and morbidity in the elderly. *N Engl J Med* 1999;341:142–7.
- 8 Bermejo J, García-Fernández MA, Torrecilla EG, et al. Effects of dobutamine on doppler echocardiographic indexes of aortic stenosis. *J Am Coll Cardiol* 1996;28:1206–13.
- 9 Baumgartner H. Should we forget about valve area when assessing aortic stenosis? *Heart* 2019;105:92–3.
- 10 Bradley SM, Foag K, Monteagudo K, et al. Use of routinely captured echocardiographic data in the diagnosis of severe aortic stenosis. *Heart* 2019;105:112–6.
- 11 Pérez Del Villar C, Yotti R, Espinosa MÁ, et al. The functional significance of paradoxical low-gradient aortic valve stenosis: hemodynamic findings during cardiopulmonary exercise testing. *JACC Cardiovasc Imaging* 2017;10:29–39.
- 12 Garcia D, Pibarot P, Dumesnil JG, et al. Assessment of aortic valve stenosis severity: a new index based on the energy loss concept. *Circulation* 2000;101:765–71.
- 13 Yotti R, Bermejo J, Antoranz JC, et al. Noninvasive assessment of ejection intraventricular pressure gradients. *J Am Coll Cardiol* 2004;43:1654–62.
- 14 Yotti R, Bermejo J, Desco MM, et al. Doppler-derived ejection intraventricular pressure gradients provide a reliable assessment of left ventricular systolic chamber function. *Circulation* 2005;112:1771–9.
- 15 Bermejo J, Antoranz JC, Burwash IG, et al. In-vivo analysis of the instantaneous transvalvular pressure difference-flow relationship in aortic valve stenosis: implications of unsteady fluid-dynamics for the clinical assessment of disease severity. *J Heart Valve Dis* 2002;11:557–66.
- 16 Yotti R, Bermejo J, Gutiérrez-Ibañes E, et al. Systemic vascular load in calcific degenerative aortic valve stenosis: insight from percutaneous valve replacement. *J Am Coll Cardiol* 2015;65:423–33.
- 17 Hachicha Z, Dumesnil JG, Bogaty P, et al. Paradoxical low-flow, low-gradient severe aortic stenosis despite preserved ejection fraction is associated with higher afterload and reduced survival. *Circulation* 2007;115:2856–64.
- 18 Jander N, Minners J, Holme I, et al. Outcome of patients with low-gradient "severe" aortic stenosis and preserved ejection fraction. *Circulation* 2011;123:887–95.
- 19 Bleakley C, Monaghan MJ. Assessment of normal-flow aortic stenosis: delving too deep? *Circ Cardiovasc Imaging* 2017;10:e007293.
- 20 Clavel MA, Berthelot-Richer M, Le Ven F, et al. Impact of classic and paradoxical low flow on survival after aortic valve replacement for severe aortic stenosis. *J Am Coll Cardiol* 2015;65:645–53.
- 21 Clavel MA, Dumesnil JG, Capoulade R, et al. Outcome of patients with aortic stenosis, small valve area, and low-flow, low-gradient despite preserved left ventricular ejection fraction. *J Am Coll Cardiol* 2012;60:1259–67.
- 22 Clavel MA, Ennezat PV, Maréchaux S, et al. Stress echocardiography to assess stenosis severity and predict outcome in patients with paradoxical low-flow, low-gradient aortic stenosis and preserved LVEF. *JACC Cardiovasc Imaging* 2013;6:175–83.
- 23 Arnold SV, Reynolds MR, Lei Y, et al. Predictors of poor outcomes after transcatheter aortic valve replacement: results from the PARTNER (Placement of Aortic Transcatheter Valve) trial. *Circulation* 2014;129:2682–90.

- 24 Baron SJ, Arnold SV, Herrmann HC, et al. Impact of ejection fraction and aortic valve gradient on outcomes of transcatheter aortic valve replacement. *J Am Coll Cardiol* 2016;67:2349–58.
- 25 Généreux P, Stone GW, O’Gara PT, et al. Natural history, diagnostic approaches, and therapeutic strategies for patients with asymptomatic severe aortic stenosis. *J Am Coll Cardiol* 2016;67:2263–88.
- 26 Caballero L, Saura D, Oliva-Sandoval MJ, et al. Three-dimensional morphology of the left ventricular outflow tract: impact on grading aortic stenosis severity. *J Am Soc Echocardiogr* 2017;30:28–35.
- 27 Hahn RT, Pibarot P. Accurate Measurement of left ventricular outflow tract diameter: comment on the updated recommendations for the echocardiographic assessment of aortic valve stenosis. *J Am Soc Echocardiogr* 2017;30:1038–41.
- 28 Khaw AV, von Bardeleben RS, Strasser C, et al. Direct measurement of left ventricular outflow tract by transthoracic real-time 3D-echocardiography increases accuracy in assessment of aortic valve stenosis. *Int J Cardiol* 2009;136:64–71.
- 29 Clavel MA, Malouf J, Messika-Zeitoun D, et al. Aortic valve area calculation in aortic stenosis by CT and Doppler echocardiography. *JACC Cardiovasc Imaging* 2015;8:248–57.