OPEN ACCESS

ORIGINAL RESEARCH ARTICLE

Higher left ventricular mass–wall stress–heart rate product and outcome in aortic valve stenosis

Eva Gerdtts ,¹ Sahrai Saeed,² Helga Midtbø,² Anne Rossebø,³ John Boyd Chambers,⁴ Eigir Einarsen,⁵ Edda Bahlmann ,⁶ Richard Devereux⁷

¹Department of Clinical Science, University of Bergen, Bergen, Norway

²Department of Heart Disease, Haukeland Universitetssjukehus, Bergen, Norway

³Cardiology Department, Oslo University Hospital, Ullevål, Norway

⁴Guys and St Thomas Hospital, London, UK

⁵Department of Clinical Science, University of Bergen, Bergen, Hordaland, Norway

⁶Cardiology, Asklepios Clinic St. Georg, Hamburg, Germany

⁷NewYork-Presbyterian Hospital/Weill Cornell Medical Center, New York, New York, USA

Correspondence to

Professor Eva Gerdtts, Department of Clinical Science, University of Bergen, Bergen 5020, Norway; Eva.Gerdtts@med.uib.no

Received 7 January 2019

Revised 8 May 2019

Accepted 9 May 2019

Published Online First

1 June 2019

ABSTRACT

Objective Whether increased myocardial oxygen demand could help explain the association of left ventricular (LV) hypertrophy with higher adverse event rate in patients with aortic valve stenosis (AS) is unknown.

Methods Data from 1522 patients with asymptomatic mostly moderate AS participating in the Simvastatin-Ezetimibe in AS study followed for a median of 4.3 years was used. High LV mass–wall stress–heart rate product was identified as >upper 95% CI limit in normal subjects. The association of higher LV mass–wall stress–heart rate product with major cardiovascular (CV) events, combined CV death and hospitalised heart failure and all-cause mortality was tested in Cox regression analyses, and reported as HR and 95% CI.

Results High LV mass–wall stress–heart rate product was found in 19% at baseline, and associated with male sex, higher body mass index, hypertension, LV hypertrophy, more severe AS and lower LV ejection fraction (all $p < 0.01$). Adjusting for these confounders in time-varying Cox regression analysis, 1 SD higher LV mass–wall stress–heart rate product was associated with higher HR of major CV events (HR 1.16(95% CI 1.06 to 1.29)), combined CV death and hospitalised heart failure (HR 1.29(95% CI 1.09 to 1.54)) and all-cause mortality (HR 1.34(95% CI 1.13 to 1.58), all $p < 0.01$).

Conclusion In patients with initially mild–moderate AS, higher LV mass–wall stress–heart rate product was associated with higher mortality and heart failure hospitalisation. Our results suggest that higher myocardial oxygen demand is contributing to the higher adverse event rate reported in AS patients with LV hypertrophy.

Trial registration number NCT000092677; Post-results.

and elevated LV ventricular wall stress.^{3–7} Experimental studies on hypertension in dogs subjected to standardised coronary artery ligation demonstrated that LV hypertrophy was associated with increased myocardial infarct size and higher rates of ventricular arrhythmias.^{8–9} However, a subsequent study suggested increased myocardial oxygen demand rather than the level of LV hypertrophy as the principal determinant of myocardial infarct size.¹⁰ In line with these experimental findings, we recently demonstrated that higher myocardial oxygen demand, measured indirectly by echocardiography as LV mass–wall stress–heart rate product, was indeed associated with higher rates of myocardial infarction and CV mortality in hypertensive patients with LV hypertrophy.¹¹

Traditionally, LV hypertrophy has been regarded as an adaptive response that keeps the LV wall stress close to normal in aortic valve stenosis (AS). However, in older subjects with AS, hypertension often coexists.¹² Recent publications have demonstrated that both LV adaptation to AS and prognosis are influenced by concomitant hypertension.^{12–13} Furthermore, presence of LV hypertrophy or a disproportionately increased LV mass in relation to actual haemodynamic load in AS was independent of hypertension associated with higher rates of heart failure hospitalisation and mortality.¹⁴ Whether higher myocardial oxygen demand may contribute to the observed impaired prognosis in AS patients with LV hypertrophy is unknown. The aim of the present analysis was to test the association of higher LV mass–wall stress–heart rate product, as an indirect measure of myocardial oxygen demand, with CV events and prognosis in AS patients participating in the Simvastatin-Ezetimibe in AS (SEAS) study.

INTRODUCTION

Left ventricular (LV) hypertrophy is a common manifestation of subclinical cardiovascular (CV) disease that strongly predicts increased risk for clinical CV disease including myocardial infarction, stroke, heart failure and death both in the general population and in patients with hypertension.^{1–2} The association of LV hypertrophy with higher CV morbidity and mortality is multifactorial and may involve reduced myocardial flow reserve, increased myocardial oxygen demand, myocardial fibrosis and scarring, as well as neuroendocrine activation, capillary rarefaction, increased production of proinflammatory cytokines, cellular dysfunction

METHODS

Study population

The SEAS study was a randomised, double-blind, placebo-controlled study performed during the years 2002–2008 which assessed the effect of combined simvastatin and ezetimibe on AS progression and CV morbidity and mortality in 1873 patients with initially asymptomatic, mild-to-moderate AS.^{13–15} Of these, the core lab received baseline echocardiograms on 1788. LV mass could be assessed in 1730 patients at baseline, and in 1656 of these also on at least 1 follow-up echocardiogram. Due to lack of end-echocardiography blood pressure measurements, the circumferential end-systolic

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Gerdtts E, Saeed S, Midtbø H, et al. *Heart* 2019;105:1629–1634.

stress (CESS) could only be calculated in 1623 patients at baseline, leaving 1522 patient with data on CESS, heart rate and LV mass measured both at the baseline and at least 1 follow-up echocardiogram eligible for the current study population. Sex, age and peak aortic jet velocity did not differ significantly between eligible and non-eligible patients (all $p>0.3$), but non-eligible patients had a lower likelihood of suffering a primary study endpoint ($p<0.001$).

All patients signed informed consent prior to study enrolment.

Echocardiography

Echocardiography was performed following a standardised study protocol at baseline and then annually and before valve surgery.^{14 15} All echocardiographic analysis was performed at the SEAS Echocardiography Core Laboratory, and 94% of final reading was done by the same experienced reader. The time range between the baseline and final echocardiogram was 2.7–84.4 months.

Quantitative echocardiographic evaluation of the LV and AS severity was performed following current guidelines.^{16 17} LV hypertrophy was considered present if LV mass/height^{2.7} >46.7 g/m^{2.7} in women and 49.2 g/m^{2.7} in men, respectively.¹⁴ Concentric LV geometry was considered present if relative wall thickness >0.420 .¹⁶ CESS was calculated by an invasively validated method,¹⁸ taking mean aortic valve gradient into account. Heart rate was measured on the echocardiograms as the average over 3 beats in patients in sinus rhythm and over 15 beats in patients with atrial fibrillation. Myocardial oxygen demand was estimated from the LV mass $\times$ CESS $\times$ heart rate product (LV mass–wall stress–heart rate product).¹¹ Midwall shortening and stress-corrected midwall shortening were calculated using previously validated equations.¹⁹ Stress-corrected LV myocardial oxygen demand was estimated from LV mass–wall stress–heart rate product.¹¹ High LV mass–wall stress–heart rate product was identified by the upper 95% CI limit calculated in a previously collected normotensive New York population ($>2.13 \times 10^6$ g kdyne/cm² bpm).²⁰ Aortic valve area adjusted for pressure recovery in the aortic root, the energy loss, was calculated as previously published and used as the primary measure of AS severity, to avoid overestimation of severity in patients with milder degree of AS.²¹ LV stroke volume was calculated by Doppler and indexed for body surface area. LV diastolic function

was assessed by peak early and atrial transmitral filling velocities, their ratio and left atrial anterior–posterior diameter.

Study endpoints

The primary endpoint was major CV events, a composite of aortic valve events (combined aortic valve replacement, CV death and hospitalisation for heart failure due to progression of AS) and ischaemic CV events (combined coronary artery bypass grafting, non-fatal myocardial infarction, hospitalisation for unstable angina pectoris, percutaneous coronary intervention and non-haemorrhagic stroke).¹⁵ All-cause mortality was a tertiary study endpoint. We also assessed hospitalisation for heart failure, combined CV death and hospitalisation for heart failure and combined all-cause mortality and hospitalisation for heart failure in the present analysis. All endpoints in the SEAS study were adjudicated by an independent committee.¹⁵

Statistical analyses

Statistical analyses were done using IBM SPSS V.25. Data are presented as mean $\pm$ SD for continuous variables and as percentages for categorical variables. Groups of patients with high and normal LV mass–wall stress–heart rate product were compared using unpaired Student's *t*-test and χ^2 test, as appropriate. Clinically relevant covariables of high LV mass–wall stress–heart rate product identified in univariable logistic regression analysis were included in the multivariable model and reported as OR and 95% CI.^{12–15} Hosmer-Lemeshow test was used to assess the goodness-of-fit for the model. The association of high versus normal baseline LV mass–wall stress–heart rate product with outcome was assessed by Kaplan-Meier plots and log rank test. The association of 1 SD higher in-study LV mass–wall stress–heart rate with HRs of study endpoints was assessed in unadjusted and adjusted time-varying Cox regression models and reported as HRs and 95% CI. In the secondary model, prevalent atrial fibrillation and use of beta-blocker treatment at baseline and time-varying left atrial diameter, stroke volume index and serum creatinine were added. The incremental predictive value of LV mass–wall stress–heart rate product over its composites was assessed by comparison of global χ^2 values in sequential Cox models including LV mass, CESS and heart rate as individual variables and the LV mass–wall stress–heart rate product,

Table 1 Baseline characteristics of the total study population and in groups with high and normal LV mass–wall stress–heart rate product

Variable	All (n=1522)	High LV mass–wall stress–heart rate product (n=289)	Normal LV mass–wall stress–heart rate product (n=1233)	P value
Age (years)	67 $\pm$ 10	68 $\pm$ 9	67 $\pm$ 10	0.272
Male (%)	61.1	74.1	58.1	<0.001
Height (m)	1.70 $\pm$ 0.09	1.73 $\pm$ 0.09	1.70 $\pm$ 0.09	<0.001
Weight (kg)	78.0 $\pm$ 14.3	84.6 $\pm$ 15.3	76.4 $\pm$ 13.6	<0.001
Body surface area (m ²)	1.89 $\pm$ 0.20	1.98 $\pm$ 0.20	1.87 $\pm$ 0.19	<0.001
Body mass index (kg/m ²)	26.8 $\pm$ 4.2	28.2 $\pm$ 4.5	26.4 $\pm$ 4.1	<0.001
Obesity (%)	19.9	30.3	17.4	<0.001
Systolic BP (mm Hg)	148 $\pm$ 20	155 $\pm$ 20	146 $\pm$ 20	<0.001
Diastolic BP (mm Hg)	83 $\pm$ 10	85 $\pm$ 11	82 $\pm$ 10	<0.001
Heart rate (bpm)	66 $\pm$ 11	71 $\pm$ 13	64 $\pm$ 10	
Hypertension (%)	83.3	92.1	81.2	<0.001
Atrial fibrillation (%)	9.7	14.9	8.4	0.001
Serum creatinine (μ mol/L)	93 $\pm$ 16	95 $\pm$ 17	93 $\pm$ 15	0.017
Haemoglobin (g/L)	139 $\pm$ 10	139 $\pm$ 12	140 $\pm$ 13	0.042

BP, blood pressure; bpm, beats per minute; LV, left ventricular.

Table 2 Echocardiographic findings at baseline in the total study population and in groups of patients with high or normal LV mass–wall stress–heart rate product

Variable	All (n=1522)	High LV mass–wall stress–heart rate product (n=289)	Normal LV mass–wall stress–heart rate product (n=1233)	P value
LV end-diastolic diameter (cm)	5.03±0.63	5.55±0.59	4.91±0.57	<0.001
LV end-systolic diameter (cm)	3.19±0.56	3.75±0.51	3.06±0.48	<0.001
Interventricular septal thickness (cm)	1.15±0.28	1.32±0.30	1.11±0.25	<0.001
Posterior wall thickness (cm)	0.89±0.19	0.98±0.19	0.86±0.18	<0.001
LV ejection fraction (%)	67±6	62±6	68±6	<0.001
Relative wall thickness	0.36±0.09	0.35±0.08	0.36±0.09	0.716
LV mass index (g/m ^{2.7})	45.5±14.4	59.9±15.6	42.1±11.8	<0.001
LV hypertrophy (%)	34.4	74.6	24.9	<0.001
Concentric LV geometry (%)	18.7	18.7	18.7	0.990
Midwall shortening (%)	17.0±3.3	15.3±2.9	17.4±3.2	<0.001
Circumferential end-systolic stress (dyne/cm ²)	129±35	158±38	122±31	
Stress-corrected midwall shortening (%)	97±19	91±18	99±19	<0.001
LV mass–wall stress–heart rate product (g kdynes/cm ² bpm)	1.6×10 ⁶ ±0.7×10 ⁶	2.8×10 ⁶ ±0.7×10 ⁶	1.3×10 ⁶ ±0.4×10 ⁶	
Peak aortic jet velocity (m/s)	3.08±0.54	3.20±0.56	3.05±0.53	<0.001
Peak aortic valve gradient (mm Hg)	39±14	42±14	38±13	<0.001
Mean aortic valve gradient (mm Hg)	23±9	25±9	22±8	<0.001
Energy loss (cm ²)	1.50±0.66	1.54±0.65	1.49±0.67	0.210
Energy loss index (cm ² /m ²)	0.90±0.46	0.84±0.36	0.91±0.48	0.009
Severe AS (%)	19.9	20.4	17.8	0.329
Stroke volume index (mL/m ²)	45±13	46±13	45±13	0.188
Left atrial diameter (cm)	3.74±0.66	3.97±0.68	3.69±0.64	<0.001
Peak early mitral filling velocity (m/s)	0.75±0.23	0.73±0.23	0.75±0.23	0.146
Peak atrial mitral filling velocity (m/s)	0.83±0.25	0.86±0.25	0.83±0.25	0.045
Peak early/peak atrial mitral filling velocity ratio	0.93±0.35	0.89±0.39	0.94±0.34	0.050
Deceleration time (ms)	227±76	223±78	228±76	0.305

AS, aortic valve stenosis; LV, left ventricular.

respectively. A two-tailed $p < 0.05$ was considered statistically significant in all analyses.

RESULTS

A high LV mass–wall stress–heart rate product was found in 19% of the study population at baseline. The group with high LV mass–wall stress–heart rate product included more men and patients with obesity and hypertension than those with normal LV mass–wall stress–heart rate product (table 1), and also had higher prevalence of atrial fibrillation and LV hypertrophy, lower LV systolic function and more severe AS, while age and LV diastolic filling was comparable in the two groups (table 2). In multivariable logistic regression analysis, presence of hypertension, male sex, lower LV ejection fraction and peak early/peak atrial transmitral velocity ratio, and higher body mass index and

peak aortic jet velocity were all independently associated with presence of high LV mass–wall stress–heart rate product (all $p < 0.001$) (table 3).

Prognostic significance of LV mass–wall stress–heart rate product

During a median of 4.3-year follow-up, the group with high LV mass–wall stress–heart rate product at baseline had higher rates of major CV events and higher all-cause mortality compared with those with normal LV mass–wall stress–heart rate product at baseline (both $p < 0.01$) (figure 1). In unadjusted time-varying Cox regression, 1 SD higher in-study LV mass–wall stress–heart rate product was associated with a 23% higher HR of major CV events, a 50% higher HR of CV death, a 33% higher HR of all-cause mortality and a 60% higher HR of heart failure hospitalisation (all $p < 0.001$) (table 4). The associations of high LV mass–wall stress–heart rate product remained significantly associated with mortality and combined mortality and hospitalised heart failure after adjustment for sex, hypertension, age and time-varying body mass index, LV hypertrophy, LV ejection fraction and peak aortic jet velocity, while the association with hospitalised heart failure was explained by the covariables (table 4, model 1). Adding prevalent atrial fibrillation and beta-blocker use at baseline, and time-varying left atrial diameter, Doppler stroke volume index and serum creatinine as covariables in a second model did not change the results (table 4, model 2). The yield (global χ^2 value) of the primary Cox model did not differ significantly when LV mass, CESS and heart rate at baseline

Table 3 Covariables of high LV mass–wall stress–heart rate product

Variable	OR (95% CI)	P value
Male sex	1.90 (1.35 to 2.68)	<0.001
Hypertension	3.40 (1.99 to 5.82)	<0.001
Body mass index (kg/m ²)	1.07 (1.03 to 1.11)	0.001
LV ejection fraction (%)	0.87 (0.85 to 0.89)	<0.001
Peak aortic jet velocity (m/s)	2.05 (1.53 to 2.73)	0.002
Left atrial diameter (cm)	1.61 (1.24 to 2.90)	<0.001
Peak early/atrial mitral velocity ratio	0.60 (0.38 to 0.96)	0.035

Multivariable logistic regression analysis in the total study population.
LV, left ventricular.

Figure 1 Association of high LV mass-wall stress-heart rate product with HRs of major cardiovascular events (A) and all-cause mortality (B) in the study population. Kaplan-Meier plots.

were included as individual variables compared with when the LV mass-wall stress-heart rate product at baseline was included ($p=0.876$), demonstrating that the prognostic information of the LV mass-wall stress-heart rate product was fully explained by the composites.

DISCUSSION

LV mass-wall stress-heart rate product and outcome

The present hypothesis-generating analysis from the large, prospective SEAS study extend previous evidence in hypertensive patients with LV hypertrophy,^{3 11} suggesting increased LV myocardial oxygen demand as a major demand-side predisposition to myocardial ischaemia and subsequent CV events in AS. Although concomitant coronary artery disease is an important cause of myocardial ischaemia, it is not a prerequisite.²² In LV hypertrophy caused by hypertension or severe AS, myocardial ischaemia has been demonstrated in absence of stenotic coronary artery disease.^{22–24} Although LV mass-wall stress-heart rate product was typically higher in patients with LV hypertrophy and the majority of these patients had concomitant hypertension, the associations with higher HRs of all-cause mortality and major CV events were independent of presence of LV hypertrophy, as well as independent of concomitant hypertension, higher body mass index and AS severity, all well-documented prognosticators in AS.^{13–15 21}

LV mass-wall stress-heart rate product and LV systolic dysfunction

The transition from compensated LV hypertrophy to congestive heart failure in AS is driven by progressive cardiomyocyte death and myocardial fibrosis.²⁵ Using contrast echocardiography, Galiuto *et al* found that myocardial hypoperfusion was associated with cardiomyocyte apoptosis in myocardial biopsies

from patients with severe AS.²⁴ In a study using late gadolinium enhancement cardiac MRI, Dweck *et al* found that among patients with moderate and severe AS, fibrosis was particularly present in the LV midwall in patients with higher LV mass, and associated with impaired prognosis.⁴ In line with these observations, patients with high LV mass-wall stress-heart rate product in the present study population had significantly lower LV midwall shortening, probably reflecting more advanced myocardial disease in this group.

LV mass-wall stress-heart rate product: relation to symptoms and CV risk factors

Previous studies have found concomitant hypertension in AS to be associated with higher combined valvuloarterial LV load and to influence LV adaptation and prognosis.^{13 26} Earlier development of AS-related symptoms has also been suggested in patients with combined AS and hypertension.²⁷ The presence of LV hypertrophy was independently associated with onset of cardinal symptoms in a prospective study of 622 patients with AS by Pellikka *et al*.²⁸ In a study of 58 patients with severe AS, Hansson *et al* used ¹¹C-acetate positron emission tomography and cardiac magnetic resonance for measurement of myocardial oxygen consumption and myocardial external efficiency, respectively.²⁹ However, myocardial oxygen consumption and myocardial external efficiency did not discriminate between asymptomatic and symptomatic patients. The present results add to this knowledge by demonstrating that concomitant hypertension was more common in AS patients with high LV mass-wall stress-heart rate product, suggesting that high myocardial oxygen demand in combined hypertension and AS may contribute to the observed earlier onset of heart failure symptoms and higher mortality in these patients.^{13 14 27} In line with previous findings in

Table 4 Associations of in-study LV mass-wall stress-heart rate product with HRs of study outcomes

Study outcome	Unadjusted HR (95% CI)	P value	Adjusted model 1* HR (95% CI)	P value	Adjusted model 2† HR (95% CI)	P value
Primary endpoint (n=456)	1.23 (1.14 to 1.32)	<0.001	1.16 (1.06 to 1.29)	0.002	1.16 (1.05 to 1.29)	0.002
Heart failure hospitalisation (n=53)	1.60 (1.37 to 1.87)	<0.001	1.18 (0.94 to 1.50)	0.162	n.a.	
CV death (n=63)	1.50 (1.31 to 1.77)	<0.001	1.42 (1.14 to 1.79)	0.002	n.a.	
Combined CV death and hospitalised HF (n=105)	1.49 (1.34 to 1.66)	<0.001	1.29 (1.09 to 1.54)	0.004	1.33 (1.11 to 1.60)	0.002
All-cause mortality (n=121)	1.33 (1.17 to 1.51)	<0.001	1.34 (1.13 to 1.58)	0.001	1.36 (1.15 to 1.62)	<0.001
Combined all-cause mortality and hospitalised HF (n=153)	1.40 (1.26 to 1.56)	<0.001	1.29 (1.12 to 1.49)	<0.001	1.32 (1.14 to 1.53)	<0.001
Non-fatal myocardial infarction (n=26)	1.40 (1.07 to 1.83)	0.016	n.a.		n.a.	

Unadjusted and adjusted time-varying Cox regression models. Results of 1 SD higher in-study LV mass-wall stress-heart rate are presented as HR and 95% CI.

*Adjusted for sex, baseline hypertension and age, and time-varying body mass index, LV hypertrophy, LV ejection fraction and peak jet velocity.

†Adjusted also for prevalent atrial fibrillation use of beta-blocker treatment at baseline and time-varying left atrial diameter, stroke volume index and serum creatinine.

CV, cardiovascular; HF, heart failure; LV, left ventricular; n.a., not applicable.

hypertension, use of beta-blocker treatment in AS was associated with lower HRs for these adverse outcomes.^{11 30}

Study limitations

The SEAS study excluded patients with diabetes, renal failure and known coronary artery disease.¹⁵ Thus, the results should be applied with caution in less selected groups of AS patients. The prospective SEAS study was performed during the years 2002–2008, and the imaging protocol did not include mitral annulus velocities by spectral tissue Doppler or recording of tricuspid regurgitation by continuous wave Doppler, as recommended by current guidelines on assessment of LV diastolic function. The assessment of the associations of LV mass–wall stress–heart rate product with LV diastolic dysfunction was therefore limited. Finally, coronary artery disease was not systematically assessed in the SEAS study. In particular, diagnosis of myocardial ischaemia by cardiac MRI, positron emission tomography, contrast stress echocardiography or exercise testing was not performed. The association of LV mass–wall stress–heart rate product as an indirect measure of myocardial oxygen demand could therefore not be compared with direct assessment of myocardial oxygen consumption by positron emission tomography.²⁹

CONCLUSION

In mostly hypertensive AS patients without known CV disease or diabetes participating in the SEAS study, higher LV mass–wall stress–heart rate product, an indirect measure of myocardial oxygen demand, was associated with higher all-cause mortality and higher HR for CV events, independent of well-known prognosticators in AS including stenosis severity and presence

of LV hypertrophy and hypertension. Whether higher myocardial oxygen demand assessed by LV mass–wall stress–heart rate product contributes to impaired prognosis through induction of chronic myocardial ischaemia should be tested in future studies using modern non-invasive imaging methods for assessment of myocardial perfusion.

Contributors EG and RD contributed to the conception and design of the work, analysis and interpretation of data and drafting and critical revision of the work. SS, HM, AR, JBC, EE and EB contributed to data collection and critical revision of the work. All authors gave final approval of the version published and agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Funding The SEAS echocardiography core laboratory was supported by the MSP Singapore Company, LLC, Singapore, a partnership between Merck & Co., and the Schering-Plough Corporation during study conduct from 2002 to 2008.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The SEAS study was approved by ethical committees in all study centres.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Eva Gerds <http://orcid.org/0000-0003-4109-2311>

Edda Bahlmann <http://orcid.org/0000-0002-4242-0535>

REFERENCES

- Devereux RB, Alderman MH. Role of preclinical cardiovascular disease in the evolution from risk factor exposure to development of morbid events. *Circulation* 1993;88:1444–55.
- Levy D, Garrison RJ, Savage DD, *et al.* Prognostic implications of echocardiographically determined left ventricular mass in the Framingham Heart Study. *N Engl J Med* 1990;322:1561–6.
- Devereux RB, Roman MJ, Palmieri V, *et al.* Left ventricular wall stresses and wall stress–mass–heart rate products in hypertensive patients with electrocardiographic left ventricular hypertrophy. *J Hypertens* 2000;18:1129–38.
- Dweck MR, Joshi S, Murigu T, *et al.* Midwall fibrosis is an independent predictor of mortality in patients with aortic stenosis. *J Am Coll Cardiol* 2011;58:1271–9.
- Liao Y, Cooper RS, McGee DL, *et al.* The relative effects of left ventricular hypertrophy, coronary artery disease, and ventricular dysfunction on survival among black adults. *JAMA* 1995;273:1592–7.
- Galderisi M, de Simone G, D'Errico A, *et al.* Independent association of coronary flow reserve with left ventricular relaxation and filling pressure in arterial hypertension. *Am J Hypertens* 2008;21:1040–6.
- Irvine T, Kenny A. Aortic stenosis and angina with normal coronary arteries: the role of coronary flow abnormalities. *Heart* 1997;78:213–4.
- Koyanagi S, Eastham CL, Harrison DG, *et al.* Increased size of myocardial infarction in dogs with chronic hypertension and left ventricular hypertrophy. *Circ Res* 1982;50:55–62.
- Koyanagi S, Eastham C, Marcus ML. Effects of chronic hypertension and left ventricular hypertrophy on the incidence of sudden cardiac death after coronary artery occlusion in conscious dogs. *Circulation* 1982;65:1192–7.
- Inou T, Lamberth WC, Koyanagi S, *et al.* Relative importance of hypertension after coronary occlusion in chronic hypertensive dogs with LVH. *Am J Physiol* 1987;253(Pt 2):H1148–H1158.
- Devereux RB, Bang CN, Roman MJ, *et al.* Left ventricular wall stress–mass–heart rate product and cardiovascular events in treated hypertensive patients: LIFE study. *Hypertension* 2015;66:945–53.
- Saeed S, Gerds E. Managing complications of hypertension in aortic valve stenosis patients. *Expert Rev Cardiovasc Ther* 2018;16:897–907.
- Rieck AE, Cramariuc D, Boman K, *et al.* Hypertension in aortic stenosis: implications for left ventricular structure and cardiovascular events. *Hypertension* 2012;60:90–7.
- Gerds E, Rossebø AB, Pedersen TR, *et al.* Relation of left ventricular mass to prognosis in initially asymptomatic mild to moderate aortic valve stenosis. *Circ Cardiovasc Imaging* 2015;8:e003644.

Key messages

What is already known on this subject?

- Presence of hypertension in aortic valve stenosis (AS) is associated with higher left ventricular (LV) mass and wall stress and predicts higher risk of ischaemic cardiovascular events and higher mortality. Increased LV mass, wall stress and heart rate all lead to increased LV myocardial oxygen demand in AS, which may predispose to myocardial ischaemia even in the absence of coronary artery disease. In hypertension, reduction of LV mass–wall stress–heart rate product by atenolol treatment was associated with improved prognosis.

What might this study add?

- The present study demonstrates that higher LV mass–wall stress–heart rate product, an indirect measure of myocardial oxygen demand, was associated with earlier onset of heart failure symptoms and increased mortality, independent of presence of other prognosticators in AS including LV hypertrophy, hypertension, stenosis severity, sex, age, body mass index and LV ejection fraction.

How might this impact on clinical practice?

- AS patients with high LV mass–wall stress–heart rate product are at increased risk for death and heart failure hospitalisation. Based on studies in hypertension, treatment with a beta-blocker may be initiated in asymptomatic AS patients with hypertension to reduce LV mass–wall stress–heart rate product and thereby myocardial oxygen demand to reduce risk for adverse events.

- 15 Rossebø AB, Pedersen TR, Boman K, *et al.* Intensive lipid lowering with simvastatin and ezetimibe in aortic stenosis. *N Engl J Med* 2008;359:1343–56.
- 16 Lang RM, Bierig M, Devereux RB, *et al.* Recommendations for chamber quantification: a report from the American Society of Echocardiography's Guidelines and Standards Committee and the Chamber Quantification Writing Group, developed in conjunction with the European Association of Echocardiography, a branch of the European Society of Cardiology. *J Am Soc Echocardiogr* 2005;18:1440–63.
- 17 Baumgartner H, Falk V, Bax JJ, *et al.* ESC/EACTS Guidelines for the management of valvular heart disease. The Task Force for the Management of Valvular Heart Disease of the European Society of Cardiology (ESC) and the European Association for Cardio-Thoracic Surgery (EACTS). *Eur Heart J* 2017;2017:2739–91.
- 18 Gaasch WH, Zile MR, Hoshino PK, *et al.* Stress-shortening relations and myocardial blood flow in compensated and failing canine hearts with pressure-overload hypertrophy. *Circulation* 1989;79:872–83.
- 19 de Simone G, Devereux RB, Koren MJ, *et al.* Midwall left ventricular mechanics. An independent predictor of cardiovascular risk in arterial hypertension. *Circulation* 1996;93:259–65.
- 20 Devereux RB, de Simone G, Pickering TG, *et al.* Relation of left ventricular midwall function to cardiovascular risk factors and arterial structure and function. *Hypertension* 1998;31:929–36.
- 21 Bahlmann E, Gerds E, Cramariuc D, *et al.* Prognostic value of energy loss index in asymptomatic aortic stenosis. *Circulation* 2013;127:1149–56.
- 22 Stanton T, Dunn FG. Hypertension, Left Ventricular Hypertrophy, and Myocardial Ischemia. *Med Clin North Am* 2017;101:29–41.
- 23 Lønnebakken MT, Rieck AE, Gerds E. Contrast stress echocardiography in hypertensive heart disease. *Cardiovasc Ultrasound* 2011;9:33.
- 24 Galiuto L, Lotrionte M, Crea F, *et al.* Impaired coronary and myocardial flow in severe aortic stenosis is associated with increased apoptosis: a transthoracic Doppler and myocardial contrast echocardiography study. *Heart* 2006;92:208–12.
- 25 Hein S, Arnon E, Kostin S, *et al.* Progression from compensated hypertrophy to failure in the pressure-overloaded human heart: structural deterioration and compensatory mechanisms. *Circulation* 2003;107:984–91.
- 26 Rieck AE, Gerds E, Lønnebakken MT, *et al.* Global left ventricular load in asymptomatic aortic stenosis: covariates and prognostic implication (the SEAS trial). *Cardiovasc Ultrasound* 2012;10:43.
- 27 Antonini-Canterin F, Huang G, Cervesato E, *et al.* Symptomatic aortic stenosis: does systemic hypertension play an additional role? *Hypertension* 2003;41:1268–72.
- 28 Pellikka PA, Sarano ME, Nishimura RA, *et al.* Outcome of 622 adults with asymptomatic, hemodynamically significant aortic stenosis during prolonged follow-up. *Circulation* 2005;111:3290–5.
- 29 Hansson NH, Sörensen J, Harms HJ, *et al.* Myocardial oxygen consumption and efficiency in aortic valve stenosis patients with and without heart failure. *J Am Heart Assoc* 2017;6:pii: e004810.
- 30 Bang CN, Greve AM, Rossebø AB, *et al.* Antihypertensive treatment with β -blockade in patients with asymptomatic aortic stenosis and association with cardiovascular events. *J Am Heart Assoc* 2017;6:pii: e006709.