Abstract 134 Table 1 Multivariate regression analysis using predefined clinical characteristics

	Odds Ratio (95% CI)	P value
Age at diagnosis (years)	0.97 (0.94, 0.99)	0.007
FH of SCD or HCM	2.18 (1.07, 4.42)	0.031
Reverse Curve	2.85 (1.45, 5.57)	0.002

Abstract 134 Table 2 The likelihood of an informative test using a predictive model

No. Factors	Likelihood of positive result (%)
3/3	75.0
2/3	63.8
1/3	29.9
0/3	20.6

Factors: Age <50, FH SCD/HCM, reverse curve septal morphology

(2.8%), ACTC1=2 (2.8%), TNNT2=2 (2.8%), MYL2=1 (1.4%), MYL3=1 (1.4%) and TPM1=1 (1.4%).

Conclusion Reverse curve morphology of the septum (OR 2.85, CI 1.45–5.57), FH of SCD or HCM (OR 2.18, CI 1.07–4.42) and younger age at testing (OR 0.97, CI 0.94–0.99), are predictors of informative genetic testing in HCM, especially when two or more of these variables occur simultaneously (Table 1).

Conflict of Interest None

135 REAL WORLD OUTCOMES FOR “INTERMEDIATE-HIGH” MORTALITY RISK PATIENTS PRESENTING WITH SUBMASSIVE PULMONARY EMBOLISM IN A TERTIARY CARDIOTHORACIC CENTRE

¹Paul Brennan*, ²Jaimie Coburn, ²Dean Carolan, ²Mark Spence, ³Timothy Warke, ³Kathy Cullen, ³Julian Leggett, ²Nicola Johnston, ²Damian McCall, ²Ganesh Manoharan. ¹Royal Victoria Hospital Belfast; ²Cardiology; ³Respiratory

10.1136/heartjnl-2019-BCS.132

Background Thrombolysis is an established recommended therapy for patients presenting with acute pulmonary embolism (PE) and shock. The management of “intermediate-high” risk patients is not as clear and the use of thrombolysis splits opinion and as such it remains a “consideration” for these patients in both the ESC and BTS guidelines. Both efficacy and safety of thrombolysis in this patient group have been variable in the major studies.

Aim sWe aimed to report real-world outcomes for all patients coded with a diagnosis of “intermediate-high” PE (SBP >90mmHg without a drop in SBP of >40mmHg, but evidence of right ventricular (RV) strain and/or cardiac biomarker rise) in our hospital between January 2016 and December 2017.

Methods Baseline characteristics, clinical and echocardiographic outcomes were obtained, via clinical note review and Northern Ireland Electronic Care Record.

Results 41 patients (mean age 65 ±14 years, female sex 59%) met the diagnosis criteria. 25% patients had an active/prior cancer diagnosis, 30% had a prior VTE/PE and the lead symptom at presentation was dyspnoea in 89% patients. Cardiac arrest was the initial presentation for 2 patients. All patients had a cardiac biomarker rise (mean troponin T-hs 121.67 ±304 ng/L, mean NTpro-BNP 1666 ±2313 ng/L). 75% patients had a TTE performed acutely and RV systolic dysfunction/strain were seen in 63% patients, while 46% patients had an RVSP>40mmHg (mean RVSP 51 ±22mmHg).

16% (n=3, all had severe RV systolic dysfunction & RVSP>40mmHg) patients received IV thrombolysis (Alteplase) and no patients had percutaneous therapy. 2 patients had an absolute contraindication to lysis agents. All of the lysed patients survived to discharge while 5% (n=2) of the non-lysed patients died as inpatients. No patients in either group required vasopressors, inotropes or mechanical ventilation. Major bleeding occurred only in the lysis group (33%, n=1) while minor bleeding occurred only in the non-lysed group (8%, n=3, all non-intracranial). Non-haemorrhagic stroke wasn't seen and there were no allergic reactions to Alteplase.

At 30 days, 1 lysed patient had an extracranial major bleed, and mortality, only seen in non-lysed patients, was 8%.

1-year mortality for lysed and non-lysed patients was 0% and 16% (all non-PE/cardiac-related) respectively. 39% patients had a recurrent VTE/PE event on anticoagulation. 23 patients (3 lysed, 20 non-lysed) had a repeat TTE performed (median time to diagnosis 120days) in whom evidence of possible chronic thromboembolic hypertension (CTEPH, (RVSP>40mmHg)) and/or RV systolic dysfunction was present in 67% (n=2) of lysed and 20% (n=4) non-lysed patients respectively.

Conclusion sIV PE lysis was administered safely and effectively in a small group of selected PE patients with severe RV systolic dysfunction and biomarker rise. The lack of cardiac mortality at 1 year in those not lysed, alongside the modest freedom from potential CTEPH, is reassuring. Further study is required to correctly allow for prediction of patients that will benefit from IV thrombolysis for submassive PE.

Conflict of Interest None

136 SAME-DAY ADMISSION FACILITATED BY A NURSE LED PATHWAY REDUCES HOSPITAL LENGTH OF STAY FOR TRANSFEMORAL TRANSCATHETER AORTIC VALVE IMPLANTATION

¹Gemma MCCALMONT, ²Paul Williams, ³Douglas Muir, ⁴Bilal Bawamia, ⁵Muhammad Arshad. ¹South Tees Hospitals NHS Foundation Trust; ²South Tees Hospitals NHS Foundation Trust; ³South Tees Hospitals NHS Foundation Trust; ⁴South Tees Hospitals NHS Foundation Trust; ⁵South Tees Hospitals NHS Foundation Trust

10.1136/heartjnl-2019-BCS.133

Background There is a growing interest in reducing the hospital length of stay for patients undergoing transcatheter aortic valve intervention (TAVI), with particular focus on patients