

Supplemental Table 2 Baseline characteristics of MINOCA patients requiring PCI

	<i>MINOCA patients with PCI</i> (n= 40)
Age (years)	56.3±10.1
Males (%)	31 (77.5)
<i>Presentation (%)</i> :	
STEMI	33 (82.5)
NSTEMI	7 (17.5)
Heart failure at presentation (%)	0 (0)
Grace Score (points)	112.8±22.7
Heart rate at presentation (bpm)	75±16
Systolic BP at presentation (mmHg)	137±20
Diastolic BP at presentation (mmHg)	84±12
Left ventricular ejection fraction (%)	55±16
Diabetes mellitus (%)	10 (25.0)
Hypertension (%)	23 (57.5)
Dyslipidaemia (%)	12 (30.0)
Family history of premature CAD (%)	8 (20.0)
Smokers (%)*	12 (30.0)
Previous MI (%)	2 (5.0)
Previous PCI (%)	4 (10.0)
Previous Stroke (%)	0 (0)
Peripheral arterial disease (%)	0 (0)
Cancer (%)	2 (5.0)
<i>Antithrombotics prior to randomization (%)</i> :	
Aspirin	12 (30.0)
Clopidogrel	4 (10.0)
Unfractionated heparin	40 (100.0)
LMWH	32 (80.0)
Fondaparinux	3 (7.5)
Bivalirudin	3 (7.5)
GP IIb/IIIa inhibitors	12 (30.0)
Fibrinolytics†	2 (28.6)
<i>Antiplatelets after randomization (%)</i> :	
Higher-dose ASA	24 (60.0)
Lower-dose ASA	16 (40.0)
Double-dose clopidogrel	22 (55.0)
Standard dose clopidogrel	18 (45.0)
<i>Baseline laboratories</i> :	
Platelets (x 10 ⁹ /L)	229.0 (197.0; 251.0)
Haemoglobin (g/dL)	14.4±1.6
WBC (x 10 ⁹ /L)	10.7±5.9
Creatinine clearance (mL/min)	82.2±17.9

Creatinine ($\mu\text{mol/L}$)	87.2 \pm 15.5
Troponin I ($\mu\text{g/L}$)	1.0 (0.3; 3.0)
Troponin T ($\mu\text{g/L}$)	0.3 (0.2; 1.8)
CK-MB (U/L)	44.3 (21.2; 88.8)
CK-MB (%)	11.5 (9.0; 14.0)
Total CK (U/L)	294.0 (169.0; 723.0)
<i>Stents (%)</i> :	
Bare metal stents (only)	27 \pm 1.7
Drug eluting stents	13 \pm 0.8
Visible angiographic thrombus before PCI (%)	17 (42.5)

Data are mean (standard deviation), median (interquartile range) or number (percentage), as appropriate. BP = Blood pressure; CAD = Indicates coronary artery disease; GP IIb/IIIa inhibitors = Glycoprotein IIb/IIIa inhibitors; LMWH = Low molecular weight heparin; MI = Myocardial infarction; MINOCA = Myocardial infarction with no obstructive coronary artery disease; NSTEMI = Non-ST-segment elevation myocardial infarction; PCI = Percutaneous coronary intervention; STEMI = ST-segment elevation myocardial infarction; WBC = White blood cell count.

* Smokers were classified as current smokers.

† As initial reperfusion strategy.