

Supplemental Table 1: Crude 30-day survival according to bystander interventions and initial shockable rhythm

	All patients (21,480)				
	Income quartile 1 5,370	Income quartile 2 5,370	Income quartile 3 5,370	Income quartile 4 5,370	p-value
30-day survival					
30-day survival overall, n (%)	226 (4.2)	342 (6.4)	479 (8.9)	1,043 (19.4)	<0.001
30-day survival in patients receiving bystander CPR, n (%)	147 (7.8)	231 (9.4)	349 (14.6)	857 (27.7)	<0.001
30-day survival in patients not receiving bystander CPR, n (%)	79 (2.3)	111 (3.8)	130 (4.4)	186 (8.2)	<0.001
30-day survival in shockable patients, n (%)	172 (15.8)	262 (21.8)	393 (27.1)	850 (40.0)	<0.001
30-day survival in shockable patients with bystander CPR and bystander defibrillation, n (%)	6 (25.0)	15 (31.9)	28 (50.9)	66 (61.7)	<0.001

30-day survival in shockable patients with bystander CPR, but no defibrillation, n (%)	101 (19.7)	161 (25.3)	241 (32.7)	585 (43.4)	<0.001
30-day survival in shockable patients with no bystander interventions, n (%)	54 (10.5)	73 (15.5)	92 (15.7)	136 (24.4)	<0.001
CPR = cardiopulmonary resuscitation; OHCA = out-of-hospital cardiac arrest					