have a post-ROSC ECG demonstrating ST elevation/LBBB and absence of epinephrine administration during resuscitation with shorter low flow times. Extent of CAD was similar between the four groups. From patients with LVEF $\geq$ 40%, patients in Group C had substantially lower SYNTAX scores than compared with Group D (0.5 vs 13.5, $p<0.001$). However, both Group B and C (RWMA) groups had highest rates of culprit lesions compared with matched global groups which was reflected in higher PCI rates (figure 2 and table 3). The primary endpoint of 12-month mortality was lowest in Group D and highest in the Group A group. A similar effect was observed for poor neurological outcome and 30-day mortality. Patients with regional LVSD had significantly improved survival at 12 months compared with those with global LVSD (70.5% vs 48.3%, $p<0.001$) vs 51). Those in Group D had

Abstract 42 Figure 2

Abstract 42 Table 1 Baseline Characteristics by Presence of RWMAs and Initial LV Systolic Function

	LVEF < 40%			LVEF $\geq$ 40%		
	RWMAs N = 29	No RWMAs N=67	P value	RWMAs N = 110	No RWMAs N=60	P value
	3	4		1	2	
Age	61 (54-66)	67 (59-75)	0.009	60 (51-69)	62 (48-70)	0.982
Male (%)	25 (86.2%)	53 (79.1%)	0.413	87 (79.1%)	38 (63.3%)	0.026
Witnessed	23 (79.3%)	58 (86.6%)	0.369	91 (82.7%)	50 (83.3%)	0.920
Residence	17 (58.6%)	35 (52.2%)	0.564	53 (48.2%)	33 (55.0%)	0.395
Location						
Bystander	20 (69.0%)	48 (71.6%)	0.791	85 (77.3%)	48 (80.1%)	0.680
CPR						
Low Flow	25.5 (11.5-36.25)	28 (17.5-40)	0.238	19 (9-28)	25 (17.5-33.5)	0.008
VF/VT	22 (75.9%)	54 (80.6%)	0.600	93 (84.5%)	47 (78.3%)	0.310
Epinephrine use	17 (58.6%)	57 (85.1%)	0.005	55 (50.9%)	40 (66.7%)	0.049
pH	7.24 (7.17-7.32)	7.21 (7.05-7.29)	0.553	7.28 (7.20-7.35)	7.22 (7.1-7.32)	0.017
Lactate	4.6 (2.7-6.6)	4.8 (2.5-9.3)	0.828	3.1 (1.7-5.8)	4.8 (2.3-7.4)	0.031

HTN, hypertension; CPR, cardiopulmonary resuscitation; VF/VT, ventricular fibrillation/ventricular tachycardia

Abstract 42 Table 2 Admission and Outcome Metrics by Presence of Regional Wall Motion Abnormalities (RWMAs) and LV Systolic Function on Initial Transthoracic Echocardiography

	LVEF < 40%			LVEF $\geq$ 40%		
	RWMAs N=29	No RWMAs N= 67	P Value	RWMAs N=110	No RWMAs N=60	P Value
ST Elevation/LBBB	26 (89.7%)	47 (70.1%)	0.039	83 (75.5%)	25 (41.7%)	<0.001
LVEF (median, IQR)	30 (30-35)	30 (30-30)	0.002	45 (40-50)	55 (45-55)	<0.001
Culprit lesion	25 (86.2%)	32 (47.8%)	<0.001	91 (82.7%)	26 (43.3%)	<0.001
PCI	25 (86.2%)	28 (41.8%)	<0.001	88 (80%)	25 (41.7%)	<0.001
SYNTAX score	19 (9-26.5)	15.5 (0-28)	0.332	13.5 (7-21.5)	0.5 (0-12)	<0.001
SCAI C-E	18 (62.1%)	44 (66.7%)	0.665	35 (32.4%)	23 (38.3%)	0.439
12 month mortality	14 (48.3%)	39 (58.2%)	0.369	27 (24.5%)	27 (45%)	0.006
30 day mortality	14 (48.3%)	37 (55.2%)	0.531	25 (22.7%)	26 (43.3%)	0.005
Cardiac Death	5/14 (35.7%)	10/41 (24.4%)	0.411	1/29 (3.4%)	2/28 (7.1%)	0.532
Poor Neurological Outcome	14 (48.3%)	46 (68.7%)	0.058	37 (33.6%)	31 (51.7%)	0.022

highest survival at 12 months, while this was similar for Groups B and C and lowest in Group A (figure 1). Cardiac aetiology death was significantly higher in those with LVEF<40% compared to those with LVEF $\geq$ 40% (70.5% vs 48.3%, $p<0.001$).

Conclusions Patients with significant LVEF <40% have higher rates of cardiogenic shock and mortality which was driven by cardiac aetiology death, while presence of RWMAs are associated with a higher rate of culprit coronary lesions and improved outcome

Conflict of Interest None

43

ST-ELEVATION MYOCARDIAL INFARCTION IN SUB-SAHARAN AFRICA: THE NEED FOR BETTER DATA

Jonathan Hudson. London School of Hygiene & Tropical Medicine, London, UK

10.1136/heartjnl-2021-BCS.43

Introduction There is growing evidence that cardiovascular disease (CVD) prevalence is increasing in Sub-Saharan Africa (SSA) (1). Historically the prevalence of MI in SSA has been thought to be low (2). However, this may be changing as parts of SSA undergo urbanization, and the rates of CVD increase(1), there is a growing awareness of the need to have health systems in place that can manage a time-sensitive cardiovascular emergency like STEMI. The acute management of STEMI presents significant challenges to health systems in SSA. A systematic review was conducted to explore the

outcomes and rates of emergency coronary reperfusion therapy in adult patients with STEMI in SSA.

Methods A literature search of two databases (MEDLINE & EMBASE) was performed up to March 1st, 2021 around the keywords 'ST-segment elevation myocardial infarction' and 'Sub-saharan Africa'. Studies were included if they were performed in SSA and reported the in-hospital mortality and emergency management of patients with STEMI. The quality of the studies was appraised using a structured checklist. All analyses were performed using STATA. The Freeman-Turkey double arcsine transformation was used to pool the data by random-effects meta-analysis. This produced a pooled prevalence of in-hospital mortality in patients with STEMI. Heterogeneity between the studies was assessed using the I² measure.

Results Study selection: A total of 12 studies from six different countries were included in the final analysis. 1096 records were retrieved from the database search and screened. 30 full texts were retrieved for consideration. STEMI outcomes: A total of 2279 patients with STEMI were included across 12 studies (table 1 & table 2). The pooled in-hospital mortality was 12.8% (95% CI 8.41, 17.92) (figure 1). When stratified by income level of the country (figure 2), STEMI in-hospital

mortality decreases as income level increases (Low-income countries (LIC) = 17.79% [95% CI 9.16, 28.39]), lower middle-income countries (LMIC) = 11.28% [95% CI 6.91, 16.50], upper middle income (UMIC) = 5.4% [95% CI 3.62, 7.97]). The pooled proportion of STEMI patients receiving emergency coronary reperfusion was 29.54% (CI 11.65, 51.11) (Fig2). Thrombolysis was the most common reperfusion therapy used. Primary PCI was rarely used as the method of coronary reperfusion with only 87 patients receiving it across all studies. Significant delays in presentation to hospital may be contributing to poor outcomes (table 2). In Burkina Faso, the average time from onset of symptoms to admission was 48hrs with 82% arriving >12hrs after symptoms onset. Two studies from Ethiopia reported an average time from onset of symptoms to admission of >90hrs.

Conclusion This systematic review demonstrates high variability in the management and outcomes of patients with STEMI in SSA and has three key findings. First, the pooled in-hospital mortality of patients with STEMI in SSA is high, at 12.8%. Second, a low proportion of patients with STEMI receive emergency coronary reperfusion. Thirdly, for patients with STEMI, there exist long delays between symptom onset and

Abstract 43 Table 1 Baseline characteristics of studies included. CS= cross sectional study. L1 low income, LMI low middle Income, UMI upper middle income

Author	Country	Income Level of country	Design of study	Type of hospital	Department	PCI capacity?	From	To	Mean age	% Men	Total number of patients in study	no. of patients with STEMI in the study
Desta et al. 2020 (3)	Ethiopia	LI	CS	Tertiary	Cardiology	Yes	2013	2018	59	72.2	151	110
Beyers et al. 2020 (4)	South Africa	UMI	CS	Tertiary	Cardiology	Yes	2017	2018	56	66.1	447	426
Yao et al. 2019 (5)	Ivory Coast	LMI	Prospective	Tertiary	ICU	Yes	2012	2015	57	82.6	329	329
Bogale et al. 2019 (6)	Ethiopia	LI	CS	Tertiary	Cardiology	No	2012	2014	56.3	75.8	124	90
Kabore 2019 (7)	Burkina Faso	LI	Prospective	Tertiary	Cardiology	No	2010	2015	57.6	77.5	111	98
Varwani 2019 (8)	Kenya	LMI	CS	Tertiary	Cardiology	Yes	2012	2013	58.7	81.2	230	101
Bahiru 2018 (9)	Kenya	LMI	CS	Tertiary	Cardiology	Yes	2013	2016	57	65	196	112
Pessinaba 2018 (10)	Togo	LI	Prospective	Tertiary	Cardiology	No	2014	2017	60	65.7	67	48
N'Guetta et al. 2016 (11)	Ivory Coast	LMI	Prospective	Tertiary	Cardiology	Yes	2010	2013	55.4	79.5	425	304
Wachira 2013 (12)	Kenya	LMI	CS	Tertiary	cardiology	Yes	2012	2013	59.7	82	45	45
Shavadia et al. 2012 (13)	Kenya	LMI	Prospective	Tertiary	ICU	Yes	2008	2010	63	82	111	62
Yameogo 2012 (14)	Burkina Faso	LI	Prospective	Tertiary	Cardiology	No	2010	2011	56	88.4	43	43

Abstract 43 Table 2 In-hospital mortality of patients with STEMI, coronary reperfusion rates, and delays

Study & Country	In Hospital STEMI mortality n/N (%)	Coronary Reperfusion therapy			Delays in presentation	
		No. patients receiving thrombolysis	No. patients receiving primary PCI	No. of patients receiving emergency coronary reperfusion n/N (%)	Reported delay	Metric used to report delay
Desta 2020 Ethiopia (3)	25/110 (22.7)	7	6	13/110 (11.82)	95.85hrs	Average time from onset of symptoms to admission
Beyers 2020 South Africa (4)	23/426 (5.4)	318	0	318/426 (74.6)	5.5hrs	Median time between symptoms onset and fibrinolysis
Yao 2019 Ivory Coast (5)	47/329 (14.2)	10	17	27/329 (8.2)	20.0hrs	Median time between onset of symptoms and admission
Bogale 2019 Ethiopia (6)	32/90 (35.6)	0	0	0/90 (0)	91.2hrs	Median time between onset of symptoms and admission
Kabore 2019 Burkina Faso (7)	9/98 (9.2)	8	0	8/98 (8.2)	82.9%	% of patients presenting >12hrs from symptom onset
Varwani 2019 Kenya (8)	15/101 (14.9)	49	19	68/101 (67.3)	>12hrs	Median time between symptoms onset and admission
Bahiru 2018 Kenya (9)	23/112 (20.4)	2	0	2/112 (1.8)	Not reported	
Pessinaba 2018 Togo (10)	6/48 (12.5)	7	0	7/48 (14.6)	81.9hrs	Average time from onset of symptoms to admission
N'Guetta 2016 Ivory Coast (11)	34/304 (11.2)	25	20	45/304 (14.8)	44.7hrs	Average time from onset of symptoms to admission
Wachira 2013 Kenya (12)	0/45 (0)	28	17	45/45 (100)	22%	% of patients presenting >12hrs from symptom onset
Shavadia 2012 Kenya (13)	6/62 (9.7)	34	8	42/62 (67.7)	12.9hrs	Average time from onset of symptoms to admission
Yameogo 2012 Burkina Faso (14)	5/43 (11.6)	2	0	2/43 (4.7)	48hrs	Average time from onset of symptoms to admission

Abstract 43 Figure 1 In hospital mortality of patients with STEMI stratified by income level of country. Effect Size (ES) is the % of patients with STEMI who died in hospital

Abstract 43 Figure 2 Meta-analysis of the proportion of patients with STEMI who received emergency coronary reperfusion (either PCI or thrombolysis) stratified by country income level

presentation to hospital. This is the first meta-analysis attempting to pool the in-hospital mortality and emergency treatment of patients with STEMI in SSA. The studies included were of moderate to poor quality and suffered from small sample sizes. As the burden of IHD increases in SSA, health systems must adapt to be able to produce better outcomes for patients presenting with STEMI.

Conflict of Interest None

44

IDENTIFYING LOW-RISK CHEST PAIN IN THE EMERGENCY DEPARTMENT WITHOUT TROPONIN TESTING: A VALIDATION STUDY OF THE HE-MACS AND HEAR RISK SCORES

Fraser Todd, James Duff, Ed Carlton. *North Bristol Trust, Bristol, UK*

10.1136/heartjnl-2021-BCS.44

Introduction Patients presenting to Emergency Departments with chest pain of possible cardiac origin represent a substantial and challenging cohort to risk stratify. Scores such as HE-MACS (History and Electrocardiogram-only Manchester Acute Coronary Syndromes decision aid) and HEAR (History, ECG, Age, Risk factors) have been developed to stratify risk without the need for troponin testing. Validation of these scores remains limited at present.

Methods We performed a post-hoc analysis of the LoDED (Limit of Detection and ECG discharge strategy) randomised-controlled trial dataset. Data collected during the trial, and prior to troponin results, were used to calculate HEAR and HE-MACS scores. Previously published thresholds of <4% for HE-MACS and <2 for HEAR defined very low risk patients. The primary outcome of MACE (major adverse cardiac events) at 30 days was used to assess diagnostic accuracy.

Results 629 patients were included, with a 7% (42/629) incidence of MACE within 30 days. HE-MACS and HEAR scores identified 85/629 and 181/629 patients predicted to be at very low risk of MACE. Within these cohorts, MACE occurred in 0/85 and 1/181 patients respectively. Sensitivity of HE-MACS and HEAR were 100% (95% CI: 91.6-100%) and 97.6% (95% CI: 87.7-99.9%) respectively. Receiver operating characteristic (ROC) curves demonstrated area under curve (AUC) of 0.80 [95%CI: 0.74-0.85] for HE-MACS and 0.76 [95% CI: 0.69-0.82] for HEAR.

Conclusion HEAR and HE-MACS show potential as rule out tools for acute myocardial infarction without the need for troponin testing. However, prospective studies are required to further validate these scores for clinical implementation.

Conflict of Interest None declared

45

COMPLETE REVASCULARISATION IS ASSOCIATED WITH IMPROVED SURVIVAL AFTER OUT OF HOSPITAL CARDIAC ARREST

Ritesh Kanyal, Nilesh Pareek, Dhruv Sarma, Apurva Bharucha, Rafal Dworakowski, Narbeh Melikian, Ian Webb, Ajay Shah, Philip McCarthy, Jonathan Byrne. *King's College Hospital, London, UK*

10.1136/heartjnl-2021-BCS.45

Introduction Coronary artery disease (CAD) is common in patients with Out of Hospital Cardiac Arrest (OOHCA) but the clinical relevance of burden of CAD and evidence for revascularisation strategies in relation to outcomes and modes of death remains unclear. This study sought to assess the extent of CAD as defined by the SYNTAX score and prognostic value of complete compared with incomplete revascularisation by evaluating the SYNTAX revascularisation index (SRI) in patients with OOHCA.

Methods 619 patients with OOHCA were admitted at our centre between 1st May 2012 and 31st December 2017. 237 were excluded for having a non-cardiac aetiology or prior neurological disability. 398 patients were included into the study and of these 272 (xx%) had early coronary angiography and were included in the final analysis. The baseline SYNTAX score (bSS) and residual SYNTAX score (rSS) were determined from the coronary angiograms by a blinded cardiologist. Patients were subdivided into 4 subgroups according to quartiles of the baseline syntax score (bSS) of 0, Group A: 1-10, Group B: 11-20 and Group C: ≥21. Complete