Abstract 81 Figure 1

Conclusion This training website seeks to provide a single source of access to the key knowledge needed for cardiac physiologists and registrars training in TTE. Our survey suggests that junior trainees will find it to be a useful adjunct to their experiential learning. The project is to continue to create content and we hope to open the website as a free, national open-access learning resource.

Conflict of Interest None

82

OUT OF SIGHT BUT NOT OUT OF MIND. A DISTRICT HOSPITAL CARDIOLOGY TELECLINIC SETUP... PATIENTS', SUSTAINABILITY AND ENVIRONMENTAL BENEFITS

¹Zhihong Yao, ²Ahmed Farag, ²Pauline Mullineux, ²Wendy Matthews, ²Mark Carimichael, ²Chris Barlow. ¹Scunthorpe General Hospital, Scunthorpe, UK; ²Warrington Hospital

10.1136/heartjnl-2021-BCS.82

Introduction The ever-greater prevalence of cardiovascular disease has placed increasing pressure on cardiology services in the UK to achieve higher efficacy and improved quality of care. Telemedicine is emerging as a cost effective way to enhance patient's care. With the COVID19 pandemic, virtual-consultation had been become more widely used. The effectiveness and safety of tele-health intervention in cardiovascular condition in the UK remains unclear.

Aim To describe our early experience and outcomes setting up virtual teleconsultation for stable follow up cardiology patients in a district general hospital.

Method Since April 2019, a single cardiologist (AF) started enrolling selected stable cardiology follow up patients onto the virtual follow-up. Following the initial standard consultation, the results of the investigations would be reviewed by the attending cardiologist and subsequent consultation would be carried out via pre-arranged telephone appointment if patients are agreeable. Patient's outcomes and satisfaction survey were collated, sustainability and social benefits and environmental impact of the pilot were analysed.

Result A total of 116 patients were enrolled into the pilot. Fifty eight percent were males with an average age of 67 years (range 26-93). Conditions enrolled were: stable

ischaemic heart disease (put numbers and percentage 42%), Stable valvular heart disease (26%), arrhythmias (13%), heart failure (9%) and others (9%).

Sixty patients (52%) were discharged and 42% (49) patients had further routine tele-clinic follow-up. There were 6% (7) patients failed to attend the telephone consultation. Patients feedback was overwhelmingly positive with the main themes being: Convenience, avoiding car parking congestion, avoiding traffic / transportation issues as well as avoiding overall visit costs. The main concerns raised in patients using tele-consultation were related: 1) Mobile phone connectivity 2) Privacy issues and 3) Logistic of completing pre-requisite investigation prior appointment. Figure 1 demonstrates the clinic's sustainability benefits. The did not attend (DNA) rates were significantly less compared to standard clinic (5.8% vs 8.2%). This pilot saved the trust c.a £3000 on consumables and logistics support and maximised outpatient clinic capacity utilisation.

Abstract 82 Figure 1 The sustainability benefits of virtual clinic

Conclusion Telemedicine had been shown to be a safe and effective method in the care of stable cardiology patients. The social, economic and environmental benefit of tele-consultation raised exciting opportunity for future service designs. Collaboration between hospital specialist, management team, community health providers and communication service suppliers are vital to ensure the sustainable implementation of telemedicine in cardiology.

Conflict of Interest none

Cardiac rhythm management

83

AUTONOMIC AND VASCULAR FUNCTION CHARACTERISTICS IN PATIENTS WITH ATRIAL HIGH RATE EPISODES

¹Ahsan Khan, ²Reem Alsharari, ³Rehan Junejo, ⁴Neil Thomas, ⁵James Fisher, ⁶G. Lip. ¹University of Birmingham, Birmingham, UK; ²Institute of Cardiovascular Sciences, University of Birmingham; ³Manchester Metropolitan University; ⁴Institute of Applied Health Research, University of Birmingham; ⁵University of Auckland; ⁶Liverpool Centre for Cardiovascular Science, University of Liverpool and Liverpool Heart & Chest

10.1136/heartjnl-2021-BCS.83

Introduction Patients who experience frequent atrial high rate episodes (AHREs), recorded on implantable cardiac devices, are at an increased risk of stroke and systemic embolism. Increased duration of AHRE is related to higher risk of thromboembolic event. The underlying pathophysiology is unclear; however, autonomic and/or endothelial dysfunction may contribute. We tested the hypothesis that patients with high AHRE burden have worse autonomic and

vascular function compared to patients with low AHRE burden.

Methods We studied 44 patients split in two groups: high AHRE burden (longest AHRE ≥ 24 hours; $n = 22$) and low AHRE burden (longest AHRE < 24 hours; $n = 22$). Resting time- and frequency-domain measures of HRV were determined to index cardiac autonomic function. High-resolution ultrasound was used to measure brachial artery diameter at rest and during reactive hyperaemia (endothelium-dependent flow mediated dilation (FMD)).

Results FMD was higher in the low AHRE burden group compared to high AHRE group (5.5% [95% confidence interval (CI):3.4–7.6] vs 3.1% [95% CI:1.9–6.2]; $p=0.04$) (table 1). Mean heart rate ($p<0.001$) and AHRE burden ($p=0.02$) were independent predictors of FMD on multivariate analysis. Time-domain, frequency-domain and non-linear indices of HRV were not significantly different between the groups. BMI was an independent predictor of HRV on stepwise multivariate analysis ($p=0.002$).

Conclusion Endothelium-dependent FMD is impaired in patients with high AHRE burden, while HRV derived indices of autonomic function were not affected by AHRE burden. Endothelial dysfunction may play an important role in the adverse outcomes seen in patients who experience frequent AHRE.

Conflict of Interest None

84

IMPACT OF THE INTRODUCTION OF AN INTEGRATED MULTI-PROFESSIONAL PATHWAY FOR ATRIAL FIBRILLATION ON LONG-TERM OUTCOMES

Saadia Aslam, Clare Vickers, Wendy Veevers, Layla Farrar, Elizabeth Hartley, Karthik Viswanathan. Calderdale and Huddersfield NHS Foundation Trust, Halifax, UK

10.1136/heartjnl-2021-BCS.84

Background An integrated structured approach to atrial fibrillation (AF) management is recommended by ESC.¹ However, limited data is available on its 'real-world' impact on anticoagulation uptake and subsequent AF-related hospitalisation and stroke rates.

Purpose To evaluate the long-term impact of the introduction of a community-wide integrated AF service based in a secondary care hospital.

Methods In September 2017 we implemented a new integrated pathway for patients with newly diagnosed 'symptomatic' atrial arrhythmias across two regions (population 450,000) in England in collaboration with primary care. All patients were seen in a one-stop multi-professional clinic (Arrhythmia nurse and Electrophysiologist with ECG and echocardiogram) within 2 weeks. They underwent standardised screening for modifiable risk factors, counselling regarding diagnosis/ anticoagulation and received an individualised AF management plan as well as access to a nurse-led telephone helpline. We followed up the first 126 consecutive patients for a period of 12 months.

Results Baseline characteristics are shown in table 1, showing a high incidence of modifiable risk factors (previously unidentified in the majority). After review in clinic, 30-day AF-related readmission rates to hospital were low (1.6%, $n=2$) as compared to historical local data (5.8%). This was sustained at 3 months (4.7%, $n=6$) and at 12 months (7.9%, $n=10$). 99% of 83 of eligible patients (CHA2DS2-VASc score of >2

Abstract 83 Table 1 Differences in autonomic and vascular function in patients with AHREs

	Low AHRE group (n = 22)	High AHRE group (n = 22)	P
Age, years	74.5 [67.5 – 81]	72 [60.8 – 82.8]	0.56
Sex (males/females)	15/7	19/3	0.15
CHA2DS2-VASc score	3 $\pm$ 2	3 $\pm$ 2	0.93
BMI (kg/m ²)	29.6 [26.9 – 32.9]	27.7 [24.8 – 36.7]	0.70
Heart rate (bpm)	69 [61 – 75]	68 [60 – 80]	0.97
Systolic BP (mm/Hg)	152 $\pm$ 18	135 $\pm$ 17	0.003
Ejection fraction (%)	59 [55 – 62]	55 [40 – 58]	0.009
Baseline diameter (mm)	4.6 [4.3 – 5.0] ^a	4.8 [4.5 – 5.2] ^a	0.36
Absolute FMD change (mm)	0.2 [0.2 – 0.4] ^b	0.1 [0.1 – 0.3] ^b	0.07
FMD (%)	5.5 [3.4 – 7.6] ^b	3.1 [1.9 – 6.2] ^b	0.04
Shear rate stimulus [sec.-1]	5592 [4174 – 9815] ^b	4173 [2458 – 10547] ^b	0.41
SDNN (ms)	41 [28 – 55] ^a	60 [39 – 80] ^a	0.12
rMSSD (ms)	40 [27 – 54] ^a	58 [39 – 77] ^a	0.10
pNN50 (%)	5 [2 – 23] ^b	17 [2 – 60] ^b	0.12
LF (ms ²)	575 [137 – 1013] ^a	665 [264 – 1593] ^a	0.83
HF (ms ²)	160 [66 – 687] ^b	112 [42 – 1839] ^b	0.98
LF normalised/HF normalised	0.8 [0.5 – 1.0] ^b	0.9 [0.5 – 1.4] ^b	0.84
SD1 (ms)	29 [19 – 38] ^a	41 [28 – 54] ^a	0.11
SD2 (ms)	49 [31 – 67] ^a	67 [47 – 87] ^a	0.15

Descriptive data are presented as numbers (with percentages). Normally distributed data are expressed as mean [95% confidence intervals (CI)]. Identified by superscript a. Non-normally distributed data are displayed as median [95%CI]. Identified by superscript b. Statistical differences were tested using independent t-test (for parametric data) or Mann-Whitney U test (for non-parametric data). Significance $p \leq 0.05$