optimize therapy. In 2 cases, there was a rupture of the shockwave balloon, and this was believed to be due to inadequate balloon preparation, the procedure continued with a replacement shockwave balloon without any complication. Baseline characteristics are described in table 1.

Discussion During this study, 2 episodes of hypotension were recorded during the procedures, one recovered immediately with the challenge of normal saline, the second patient who was admitted as ACS (NSTEMI), had a heavy calcified left main disease that required bifurcation stenting, Intra-aortic balloon was required for 24 hours. The patient was discharged

home after 7 days. Two years' follow-up showed 4 cases required to repeat coronary angiography; One case atypical chest pain with the non-obstructive disease, the second case had new lesions in different coronary arteries. The third; with prior history of CABG and stent to the native RCA with ISR had IVUS guided NC ballooning. The last one also prior to CABG, had severe ISR of his native LM/LCX even after high-pressure balloon and DEB treatment. The patient was referred for Laser ablation ± Re-do CABG. Unfortunately, due to comorbidities this patient was not fit for these procedures and passed away after two months.

Conclusion Initial experience with intravascular lithotripsy has demonstrated good angiographic outcomes without significant complications even after stent implantation. S-IVL has a minimal learning curve and has been quickly adopted as a safe and successful alternative to Rotablation. If shockwaves fail Rotablation still remains an option (2 of our cases). Patients with a previous history of CABG had a lower success rate reflective of the chronicity and complexity of the disease.

Abstract 24 Table 1 Procedure approach

Procedure approach, N 37	Number of Cases (%)
Arterial approach, n (%)	R Radial 35 (94.5%) R Femoral 2 (5.4%)
Catheter size	6 Fr 27 (72.9%) 7Fr 10 (27%)
Guiding catheter	
Voda L	17 (out of 27 for L+ system)
CLS	9
EBU	1
AL1	7 (out of 10 for R+ system)
FR 4	3
JR 4 Guide	1
Treated Vessels:	
-LAD	18 (48.6%)
-RCA	10 (27%)
-LM/LAD	5 (13.5%)
-LCX	2 (5.4%)
-LM/LCX	1(2,7%)
-LM/LAD/LCX	1(2,7%)
Predilation, n (%)	33/37 (89.1%)
	2.0: 3/37 2.5: 16/37 3.0: 8/37 3.5: 6/37 N: 4/37
Maximal Predilation balloon Diameter mm	
Adjunctive devices	
Guideliner Support	6 (16.2%)
Rotational Atherectomy	2(5.4%)
Shockwave Balloon	
	3.0x12 (18 cases) 3.5x12 (11 cases) 4.0x12 (5 case) 2.5x12 (3 case)
Shockwave Balloon diameter	
Average Pulse	50 +/- 12
Stenting	35/37 (94.5%)
	3.0: 12/35 3.5: 18/35 4.0: 5/35
Maximal stent diameter mm	
Post dilation, n (%)	37 (100%)
	3.0: 2/37 3.5: 16/37 4.0: 11/37 4.5: 8/37
Maximal post-dilation balloon diameter, mm	

25 CT CORONARY ANGIOGRAPHY AND COVID-19: INPATIENT USE IN ACUTE CHEST PAIN SERVICE

M Cronin, P Wheen, R Armstrong, R Kumar, A McMahon, M White, N Sheehy, G McMahon, R Murphy. *St. Lukes Hospital Kilkenny, Ireland*

10.1136/heartjnl-2021-ICS.25

Introduction CT coronary angiography (CTCA) is a well-validated clinical tool in the evaluation of chest pain. In January 2020 a care pathway was agreed between the cardiology, radiology and emergency departments in our facility to implement a strategy of CTCA as a first line inpatient examination for selected patients with chest pain of low to moderate risk of significant coronary artery disease. Impact on length of stay and clinical outcomes including subsequent invasive angiography and revascularisation were evaluated, including major adverse cardiac events at 30 days.

Methods Study design was retrospective. Patients referred for cardiology review between October 2019 and May 2020 with chest pain and/or dyspnoea were broken into three cohorts: a baseline cohort, a cohort with increased CTCA availability and a cohort with increased CTCA availability, but after the national lockdown due to COVID-19 in Ireland. These patients were identified via the Hospital Inpatient Enquiry dataset and correlated with chart review. Statistical analysis

Abstract 25 Figure 1 Number of CTCA (% of performance in overall cohort)

was via SPSS. Ethical approval was provided by the research and ethics committee in St. James Hospital, Dublin.

Results Between October 2019 and May 2020 513 patients (35.3% female) presented between cohorts 1 (n=179), 2 (n=182), and 3 (n=153). Overall population characteristics, investigations and outcomes are available in table 1. This same information is available for those who underwent CTCA in table 2. Daily rate of ED referral to cardiology increased from 2.32 to 2.46 and 2.73 patients. Overall length of stay between cohorts 1 and 3 decreased from 2.66 days median to 1.71 days ($p = 0.014$). 73 inpatient CTCA were performed over the 8 month period, increasing from 7.8% overall in cohort 1% to 20.4% in cohort 3 (see figure 1). A history of

Abstract 25 Table 1 Overall characteristics

	14/10/19 – 30/12/19	13/1/20 – 27/3/20	27/3/20 – 22/5/20
Total	179	182	153
Female	59 (33%)	71 (39.9%)	51 (33.3%)
Presenting complaint	153 Chest pain (85.5%) 26 Dyspnoea (14.5%)	151 Chest Pain (83%) 6 Dyspnoea (3%) 25 Chest pain/Dyspnoea (14%)	131 Chest Pain (85.6%) 8 Dyspnoea (5.2%) 14 Chest pain/Dyspnoea (9.2%)
Age	62.6 (55.7 – 71.8)	59 (50 – 70)	59 (50 – 70)
Length of stay (days)	2.66 (0.96 – 4.92)	2.16 (1.2 – 5.1)	1.71 (1 – 3.3)
Family history	84 (46.9%)	41 (22.5%)	40 (26.1%)
Hypertension	100 (55.9%)	85 (46.7%)	60 (39.2%)
Diabetes Mellitus	31 (17.3%)	20 (11%)	12 (7.8%)
Smoker	49 Current (27.4%) 66 Ex (36.9%) 64 Never (35.7%)	46 Current (25.3%) 61 Ex (33.5%) 75 Never (41.2%)	38 Current (25%) 53 Ex (34.9%) 62 Never (40.1%)
Dyslipidaemia	94 (52.5%)	97 (53.3%)	71 (46.4%)
History of cardiovascular/cerebrovascular disease	74 (41.3%)	67 (36.8%)	50 (32.7%)
Systolic blood pressure	126 (110 – 114)	136 (121 – 153)	144 (125 – 156)
Diastolic blood pressure	74 (66 – 81.5)	81 (72 – 89)	82 (74 – 89)
Heart rate	70 (64 – 79)	76 (66 – 86)	72 (65 – 84)
Weight (kilogram)	78 (69.9 – 90.9)*	80 (70 – 89.4)*	80.5 (73.8 – 93.5)*
New ECG ischaemia	63 (35.2%)	36 (19.8%)	22 (14.3%)
Troponin-T (ng/L)	15 (0 – 44)	0 (0 – 21.5)	0 (0 – 20)
eGFR (ml/min/1.73m²)	84 (67 – 99)	90 (75 – 107)	89 (72 – 103)
CTCA amount	14 (7.8%)	28 (15.4%)	31 (20.3%)
Exercise stress test	43 (24%)	47 (25.8%)	0 (0%)
Invasive Coronary Angiography	82 (45.8%)	71 (39%)	52 (33.9%)
Revascularisation performed	53 (29.6%)	29 (15.9%)	25 (16.3%)
Death at 1 month	6 (3.4%)	1 (0.5%)	2 (1.3%)
Myocardial infarction at 1 month	0	1	1

Outpatient functional/anatomical imaging	Total 23 (12.8%) 9	Total 28 (15.4%) 10	Total 26 (16.9%)
	Invasive coronary angiogram (39%)	Stress perfusion cardiac magnetic resonance imaging (35.7%)	11 CT coronary angiography (42.3%)
	4 Stress perfusion cardiac magnetic resonance imaging (17.4%)	6 Invasive coronary angiogram (21.4%)	9 Invasive coronary angiogram (34.6%)
	4 Dobutamine stress echocardiography (17.4%)	5 CT coronary angiography (18%)	4 Stress perfusion cardiac magnetic resonance imaging (15.4%)
	3 Single-photo emission computed tomography (13.1%)	4 Single-photo emission computed tomography (14%)	1 Dobutamine stress echocardiography (3.85%)
	3 CT coronary angiography (13.1%)	3 Dobutamine stress echocardiography (10.9%)	1 Single-photo emission computed tomography (3.85%)

Weight: Not available = 36 in cohort 1, 39 in cohort 2, 50 in cohort 3

Abstract 25 Figure 2 Overall/CTCA length of stay median/IQR (days)

Abstract 25 Table 2 CTCA cohort characteristics

	14/10/19 – 30/12/19	13/1/20 – 27/3/20	27/3/20 – 22/5/20	14/10/19 – 27/3/20
Total	14 (7.8%)	28 (15.4%)	31 (20.4%)	73
Female	5 (35.7%)	14 (50%)	12 (38.7%)	31 (42.5%)
Age	61.1 (57.8 – 67.4)	50.5 (44.8 – 56.5)	54 (50 – 60.5)	55 (49 – 60)
Length of stay (days)	4.2 (2.16 – 5.5)	1.98 (1.4 – 2.4)	2.5 (1.9 – 4.5)	2.2 (1.7 – 4.3)
Family History	5 (36%)	8 (28.6%)	10 (32.3%)	23 (31.5%)
Hypertension	6 (42.9%)	4 (14.3%)	10 (32.3%)	20 (27.4%)
Diabetes Mellitus	2 (14.3%)	1 (3.6%)	0 (0%)	3 (4.1%)
Smoker	Current = 4 (28.6%) Ex = 6 (42.8%) Never = 4 (28.6%)	Current = 9 (32.1%) Ex = 7 (25%) Never = 12 (42.9%)	Current = 9 (29%) Ex = 11 (35.5%) Never = 11 (35.5%)	Current = 22 (30.1%) Ex = 24 (32.9%) Never = 27 (37%)
Dyslipidaemia	5 (35.7%)	12 (42.9%)	10 (32.3%)	27 (37%)
History of cardiovascular/cerebrovascular disease	0 (0%)	3 (10.7%)	0 (0%)	3 (4.1%)
Weight (kilogram)	74 (64 – 80.8)*	72 (65.8 – 78.8)*	80.2 (74.5 – 88.2)*	76 (69 – 83.5)*

New ECG ischaemia	1 (7.1%)	4 (14.3%)	2 (6.4%)	7 (9.6%)
Troponin-T (ng/L)	0 (0 – 0)	0 (0 – 0)	0 (0 – 0)	0 (0 – 0)
Exercise stress test	6 (42.9%)	7 (25%)	0 (0%)	13 (17.8%)
Time to CTCA (hours)	23 (6.8 – 45.3)	20.16 (16.8 – 32.3)	30.1 (0.4 – 92.5)	22 (8.2 – 43.5)
Calcium score	68.45 (1.6 – 103.9)*	0 (0 – 6)	0 (0 – 25)*	0 (0 – 31)*
% CTCA with an abnormal finding	57	35.7	54.8	47.9
Beta-blocker used for rate control	7 Nebivolol (50%) 3 Metoprolol (IV) (21.4%) 2 Metoprolol (PO) (14.3%) 1 Bisoprolol (7.15%) 1 None needed (7.15%)	21 Nebivolol (75%) 4 Metoprolol (PO) (14.3%) 2 Metoprolol (IV) (7.1%) 1 Propranolol (3.6%)	17 Nebivolol (61.6%) 9 Metoprolol (IV) (32.3%) 3 None needed (10%) 1 Bisoprolol (3.3%) 1 Unable to control (3.3%)	45 Nebivolol (61.6%) 14 Metoprolol (IV) (19.2%) 6 Metoprolol (PO) (8.2%) 4 None needed (5.5%) 2 Bisoprolol (2.7%) 1 Propranolol (1.3%) 1 Unable to control (1.3%)
Invasive coronary angiography	5 (35.7%)	5 (17.8%)	7 (22.6%)	17 (23.3%)
Revascularisation performed	3 (21.4%)	3 (10.7%)	2 (6.5%)	8 (10.9%)
Alive at 1 month	14	28	31	73 (100%)
MI at 1 month	0	0	0	0

Weight: Not available = 3 in cohort 1, 4 in cohort 2, 5 in cohort 3
Calcium score: Not available = 4 in cohort 1, 4 in cohort 3

cerebrovascular/cardiovascular disease was significantly associated with CTCA use ($p = 0.00097$) in cohort 1. Age ($p = 0.0017$), hypertension ($p = 0.001$) and history of cerebrovascular/cardiovascular disease ($p = 0.0015$) were associated with CTCA use in cohort 2. Diabetes ($p = 0.037$) and history of cerebrovascular/cardiovascular disease ($p = 0.000246$) were associated with CTCA use in cohort 3. Overall length of stay for the patients undergoing CTCA decreased from a median of 4.2 days in cohort 1 to 2.5 days in cohort 3 (see figure 2),

with no increase in 30 days adverse outcomes. Invasive coronary angiography rate dropped from 45.8% to 39% in cohort 2, and to 34.2% in cohort 3 ($p = 0.0286$). Revascularization was required in 29.6% in cohort 1, 15.9% in cohort 2, and 16.3% in cohort 3 ($p = 0.00448$).

Conclusions This study demonstrated an increase in inpatient CTCA provision. The doubling of proportion of patients investigated by CTCA from 7.8% to 15.4% was associated with a reduction in length of stay from 4.2 days to 1.98 days. The improved efficiency in processing these intermediate rather than high risk patients was maintained during the COVID-19 pandemic, with a continued lower length of stay. A corresponding reduction in the rates of invasive coronary angiography and revascularization was seen, without an increase in 30-day major adverse cardiac events. We envisage our findings to be referenced when encouraging the development of CTCA use within national healthcare systems, particularly in light of the increasing trend towards shortening length of hospital admissions within the context of an international pandemic.

26

CARDIAC DEVICE REMOTE MONITORING – FREQUENCY OF CLINICALLY IMPORTANT EVENT IN RELATION TO INTERVENTION

¹LR Hamel, ²A Mittal, ²A Rehman, ²J O'Neill, ²J Galvin, ²E Keelan, ²R O'Mahony, ²T O'Neill.
¹Royal College of Surgeons in Ireland – Medical University of Bahrain, Bahrain; ²Connolly Hospital Blanchardstown, Dublin, Ireland

10.1136/heartjnl-2021-ICS.26

Introduction Remote Cardiac Device Monitoring (RCDM) has been an advancing technology in the field for many years. Assessing the efficacy, safety and logistics of the RCDM ensures patient safety. This study set out to examine the time from clinically relevant events to an intervention in view of the potential risk of either missed events or delay in discovery of such events. There are no national standardized operating procedures concerning RCDM in Ireland, which puts responsibility on cardiac teams to ensure in-house protocols and stringent strategies are in place to provide best practice and patient safety.

Abstract 26 Figure 1 Total number of transmissions to average time to intervention in days in all three companies