

## Correction: *Heart failure medication dosage and survival in women and men seen at outpatient clinics*

Bots SH, Onland-Moret NC, Tulevski II, *et al.* Heart failure medication dosage and survival in women and men seen at outpatient clinics. *Heart* 2021;107:1748–55.

This article has been corrected since it was first published to correct . The rows were misaligned in the 'Ejection fraction' section; please see below for the corrected portion of the table.

	Women	Men
Ejection fraction (n, %)		
<40%	162 (28.9)	291 (47.3)
40%–49%	81 (15.3)	123 (20.0)
>50%	272 (48.5)	155 (25.2)
Not recorded	41 (7.3)	46 (7.5)



### OPEN ACCESS

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

*Heart* 2022;108:e4. doi:10.1136/heartjnl-2021-319229corr1

